1

Daubenmire Data Sheet

Sampling Date and Year:

of tot at standing 50m of took as sprowted to sprowted

V051 Site:

Data Collected by: KELBEY HOWN & ROYCE on this date: 6/27/16

| Woody Bare Leaf Litter Species Ground Litter Depth | 20 | , 98 | 200 | Wo 450 | S 860 4cm | П | + | | | |
|---|-----|------|-----|--------|-----------|----|---|---|---|--|
| Milkweed | 0 | 0 | 0 | 0 | 0 | P | | | | |
| Forbs | T. | 0 | 3 | 80 | 9 | 0 | | | | |
| WSG | 0 | 0 | 0 | 0 | 0 | 0 | | | | |
| 982 | 200 | 9 | 9) | 91 | 38 | 38 | | | | |
| | - | 2 | က | 4 | 5 | 9 | 7 | œ | 6 | |

| | ١ |
|----------|---|
| | ŀ |
| | ı |
| | ı |
| 0 | ١ |
| - | ı |
| 0 | ı |
| 1 | ı |
| | ı |
| a | ı |
| ڡ | ١ |
| 0 | ١ |
| α | ١ |

| 10 | | | | |
|----|---|---|---|---|
| | | | | |
| 0 | | | | |
| œ | | | | |
| 7 | | | | |
| • | N | 2 | 4 | M |
| 2 | 3 | 5 | 2 | 9 |
| 4 | 1 | | 7 | 1 |
| m | Н | 3 | 7 | 8 |
| 2 | 7 | 8 | 2 | 4 |
| 1 | 7 | | 1 | |
| | Z | ш | S | * |

Bieveneter = 50m transect

98 - 95%-100% 86 - 75%-95% 38 - 25%-50% 63 - 50%-75% 16 - 5%-25% Categories: $\underline{0}$ - absent $\underline{1}$ - trace $\underline{3}$ - trace-5%

CSG - cool season grasses (including sedges)

Forbs - forbs not including milkweed WSG - warm season grasses

Milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material), Litter Depth - to nearest .25 cm