Daubenmire Datasheet Date:

Observer:

Recorder:

Team lead:

Round:

Transect:

vos1

tvos1a 2

50m	-

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	3	Ø	63	Ø	Ø	Ø	Ø	Ø	3	810	6.0
10 m	10	16	86	Ø	Ø	Ø,	Ø_	Ø,	38	63	2.0
20 m	10	38	63	Ø	Ø	Ø	Ø	Ø	16	63	3.0
30 m	3	16	43	8	8	Ø	Ď,	Ø	10	63	3.0
40 m	3	16	38	Ø	Ø	0	Ø	S.	38	63	4.0
50 m	\wedge \wedge \wedge		\sim								
60 m											
70 m						$\ell = \lambda I$					
80 m							and the second				
90 m		V									

Robel Pole:

	O m	10 m	20 m	30 m	40 m	50 m 60 m 70 m 80 m 90 m
N	1	2	2	2	3	
Ε	5	1		5	3	Commence of the commence of th
s	3	3	3	2	3	
W	1	2	2	3	(0	

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm