Date: Recorder: Observer: Daub-imire Datasheet Team lead: 14-00 Site: Round: Transect: tbcr1a 50m bcr1

	CS6	WS6	Forbs	Milkweed	Common	Swamp	Butterfly	Woody	Bare	Leaf	Litter
0	20	2	2 2)	ramer #	ramet #	ramet #	Species	Ground	Litter	Depth (cm)
3	10	. 0	58	C	C	0	O	O.	0	6	_
10 m	16	_	208	0	0	\supset))	2	12.2	
20 8	0/1		1.0) () (C		C	600	
20 =	- 8		22	C	C	0	0	O		00000	J 0
30 m	100		600	0	0)	<u>ي</u>))	N	200	- (
40 E	N	N	102	})) () (3	. (Do	_
0	((C		16	S	· S
300									,		(
60 m											
70 m											
10											
80 m											
90 m											
Robel Pole:	ole:										

	z	m	S	8
m 0	W	S	F	5
10 m	7	U	0)	4
20 m	W	(1)	7	6
30 m	S	ଠୀ	N	Λ
40 m	0	S	H	(N-
50 m		4		
60 m				
70 m				
80 m				
90 m				

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm