Daubenmire Datasheet

Date: 81/3/18

Observer:

Recorder:

Team lead: North

Site:

Transect: _ pre1b

tpre1b

Round:

100m

ω

	CS6	WS6	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare	Leaf Litter	Litter Depth (cm)
O #	0		28	0		0	0	0	V		0.25
10 m	0	0	98	0	Ø	Q	9	0	27		-
20 m	<u></u>	0	63	0	O	0	0	0	\sim		0.5
30 m	3	0	56		0	Ø	0	Ø.	W	_	_
40 m	16	0	63	O	Ø	0	0	9		v	_
50 m	3	0	63	0	0		Ø	0		~	್ಷಾ
60 m	_	0	98	0		0		0.		8	0.5
70 m		0	98	0		0	0.	0		W	0.25
80 m	5		63	0	0	0	0		***************************************	~	0.5
90 m	3	0	5	0	Q	C	0	E	3	7	27.0

	0 3	10 m	20 m	30 m	40 m		50 m		50 m
Z	6	12	8	2	-		13	(3)	8 8
Е	9)	15	2	0	8	WILDOWS	(3	13 16	13 16 16
S	3	1)	8	8	9		0	4	5 + 8
8	31		,		,		2))

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground – includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm