Daubenmire Datasheet

Date: 8-12-19

Observer: \_\_\_\_\_\_\_\_\_

Recorder: Name

Team lead: \_\_\_

Si	te	:
----	----	---

nor1

Transect:

tnor1b

Round:

3 50m

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	63	38	3	0				0	0	38	1.60
10 m	16	4	63	0		_		0	38	10	0,25
20 m	3	0	63	0	-	_		0	(03	3	0,25
30 m	80	0	160	9				0	10	63	/.00
40 m	63	0	38	0				0	16	63	1-00
50 m							35		7		
60 m					1						
70 m											
80 m											
90 m											

Robel Pole:

	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	5	\(	14	10	16					
E	2	12	16	15	14					
5	10	12	13		16					
W	à	12	14	14	14					

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm