Daurenmire Datasheet

Date: 6-13-19

Recorder: Seth Observer: Sex

Team lead: Jesh

Site: Transect: tprd2a prd2

100m

Round:

	CS6	WS6	Forbs	Milkweed	Common	Swamp	Butterfly ramet #	Woody	Bare	Leaf	Litter Depth (cm)
0 m	63	0	91	0	0	0	0	0	0	88	1.75
10 m	63	0	16	0	0	0	0	0	0	88	5,0
20 m	38	0	38	0	0	0	0	0	0	88	4.0
30 m	16	0	38	0	0	0	0	0	0	98	3,5
40 m	16	16	16	0	0	0	0	0	_	98	5.0
50 m	38	0	16	0	0	0	0	0	0	98	3, 0
60 m	38	0	38	0	0	0	0	0	-	98	3.0
70 m	9)	~	38	0	0	0	0	0	_	98	0.5
80 m	63	w	16	0	0	0	0	0	w	8.6	3.0
90 m	16	0	38	0	0	0	0	0	S	38	0.1
Robel Pole:	onle:										

50 m	60 m 70
	50 m

80 m

90 m

does conada rye full over + form a thick mat in hormant season?

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100% probardyrass 0-5m

WSG - warm season grasses

€

S m

0

2

w

\_

2

7

I

T

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm