Daubenmire Datasheet Date: Observer:

Recorder:

Team lead:

nkn1 Site: tnkn1a Transect:

Round:

2

100m

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	38	0	38	0					86	3	0,25
10 m	98	0	0	\bigcirc				0	16	86	0.75
20 m	98	0	3	0				0	16	86	1.0
30 m	86	0	16	0		1		0	3	86	1.0
40 m	86	0	16	0					38	63	0,5
50 m	16	0	86	0		X		6	0	98	2.0
60 m	38	0	30	0	1			\circ	38	63	1.0
70 m	38	(2)	63	0				0	16	86	1-5
80 m	0	0	98	0				0	16	86	1.5
90 m	38		63	0				0	3	98	1.5

Robel Pole:

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	(0	8	3	6	8	5	10	8	16	7
Ε	4	1	3	5		3	13	7	13	7
s	1	6	5	6		4	11	9	13	6
W	3	1	3	7	8	3	10	10	13	10

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm