1

* only did (50m) Temsects So no # 7-10 needed

Sampling Date and Year:

Daubenmir ata Sheet

Site: Van

Jackie A, Bub. H. Data Collected by: Sach New

Litter	Depth	200		boun	CENT	200		O . C	N	S C C C	T			
Leaf	Litter	2	0	300	20	63	100	3	0.0					
Bare	Ground	M	a)(0	~	0	0	a			4.		
Woody	Species	0	E	3)	0	Q)	4					
							N.							
Milkweed	(.0)(C) -)	0			4	up)	
Forbs	0	0	2	160		3	200	2	7					
WSG	E	5.	2	9	110	2)	28	10	20					
952	(9.3	\$:	9	0)/	11.0	3	0)	00	20					
	-		2	m	4	- 1	S.	9		/ /	Ø		6	10

0
0
0
_
0
0
0
-
X
_

10				
6				
80				
1	Not			
/			÷	
9	5	6	×	0
വ	0	0	0	
4	1	0)	2	00
က	9	9	7	
2	(0)	9	1	2
1	10	5	7	7
5	Z	В	S	*

<u>63</u> - 50%-75% <u>86</u> - 75%-95% <u>98</u> - 95%-100% 38 - 25%-50% 16 - 5%-25% 3 - trace-5% CSG - cool season grasses (including sedges) Categories: 0 - absent 1 - trace

WSG - warm season grasses

Forbs - forbs not including milkweed

Milkweed

Wood - woody cover

Leaf Litter (horizontal dead material), Litter Depth - to nearest .25 cm Bare Ground - includes rocks, lichens, moss, branches, and cow pies