Date: Daubenmire Datasheet

Observer:

Recorder: Team lead:

> Site: tcre1a cre1

Transect: 1

Round: 100m

				Ma		711	9		I			T
	Robel Pole:	90 m	80 m	>70 m	60 m	50 m	40 m	30 m	20 m	10 m	0	
O E	ole:	Q	16	100	16	1(0	0)8	28	38	16	160	CS6
10 m	,	Ø	'Q	8	8	Ø	Q	Q	Ø	w	Q	WS6
20 m		Q	\propto	\varnothing	A)	\propto	~	<i>∞</i>		.\	_	
30 m		866	0	6	0	9	86	0	₹.	XX	S	Forbs
40 E		Ø	X	R	Q	D	Q	Q	Q)	Q	Ø	Milkweed
50	1	S/S	2	R	Q	D	Q	Q	Q	Q	Q	ramet #
60 m 70 m	Parish	SP	8/8	27	3/2	×)	(d	Q	2 8	XX	-	Swamp ramet #
200	0	S/K	8/8	25	X	SK	2	S/O	X R	2/2	2	Butterfly ramet #
3	0	2/2	1	2 8	Q	3/3	S R	2,2	20	28	Pocies	Woody
	O	56	160	70	JC	NO	V	12	N ~	751	OI OUNG	Bare
	80	200		S C	26	Q E	56	500	600	- 6	Liller	Leaf
	7,00		00.7	9.0	7.00	5.00	0, 10	000	0.50	000	рерти (ст)	Litter

10 m				-	20
U)	W	N	A	Л	4
(((C	(-/
1	Л	7)	4	2	>
(((,	1	C
N	7	4	7	_) [
(-			-	(
J		1	0	N	N
1 17 1	01000		- P U7 W	- PUNW	~ P C N CV + C + C N CV + C + C N CV +

CSG - cool season grasses (including sedges) trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed Wood - woody cover

Bare Ground – includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm