Date: Recorder: Observer: Daubenmire Datasheet Site: Round: Transect:

Team lead:

tdun2a dun2 Н

100m

	CS6	WS6	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare	Leaf Litter	Litter Depth (cm)
0	20	Ø	38	\$	· ·	Designation of the Control of the Co)	Ф	(V)	6	
10 m	20	Þ	86	P			1	6	(v	6	ر
20 m	20	P	6	Φ	1	\	1	ф	(V)	(N)	ىر
30 m	200	φ	6	Ф	()		В	16	38	9
	52	Ф	63	P	1)	Ф	16	6	
50 m	200	P	0	B	1	\	1	ф	(h	-6	
60 m	000	Þ	503	Φ	1			þ	6	160	0,5
70 m	5 (P "	50	ф			The second secon	Ф	6	38	CV
80 m	5	φ	26	φ		1		Ф	<u>-</u> 6	3	ى
90 m	6	P	38	Φ				P	16	2	(v
	-										

_	
	RO
	<u>6</u>
	Po
	6

3		30 m	3	3	т 40 m	m 40 m 50 m 60	m 40 m 50 m 60 m
5	1	6	(V)) C>) C>) C>
A 5		7	7 (0	6	6	(0)	60 7 55
7 G		6	(M)	6	6 0 7	7	7
					Л		

Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100% CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm