Daubenmire Datasheet

Date:

Observer:

Recorder:

Team lead:

Site:	nie1		
Transect:	tnie1a		

Round: 50m

rfly	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
	Ø	1(0	86	5.00
(- N.	8	3	86	2.00
8	8	3	86	2.00

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Depth (cm)
0 m	1(0	38	8(0	Ø	Ø	Ø	Ø	Ø	10	86	5.00
10 m	3	10	818	Ø	Ø)	8	Ø	3	000	2.00
20 m	1(0	3	86	3		0	9	Ø	5		0.75
30 m	38	3	63	8	8	0	0	0	10	86	1.00
40 m	10	X	63	0	8	8	0	X	10	900	1.00
50 m∕	\sim	~									
60 m						1	//		\bigcap	/ \ /	
70 m						1					V
80 m	9										
90 m					<u> </u>						

Robel Pole:

bel Po	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	11	7	8	9	Ol	//				
Е	8	13	5	9	8	5				
s	8	(0)	7	19	10) (
W	7	10	7	10	12	_	<i></i>	The same of the sa	<	

Categories: 0 - absent 1 - trace 3 - trace-5% 16 - 5%-25% 38 - 25%-50% 63 - 50%-75% 86 - 75%-95% 98 - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm