Daubenmire Datasheet		Site:
Date:	12/19	Transect:
Observer:	Kusux	Round:
Recorder:		Noung
Team lead:	~	

								445.0			
2 V	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	38	8	103	0	8	9	2	12	1.0	38	Deprii (ciii
10 m	63	8	43	Ø	a	X	8	X	25	00	0.50
20 m	1	0	98	0	X	08	0	X	17	63	0.00
30 m	63	0	1(0	a	X	X	X	100	65	58	0.50
40 m	10	Ø	7/e	X	(X	8	X	10	10	63	0.75
50 m	28	Ò	(03	X	8	2	8		10	80	0.75
60 m	103	8	24	20	20	9	8	8	10	86	0.50
70 m	3	(8	30	9	8	0	0	0	10	8(2	0.50
	8le	X	10	Ø	8	0	0	8	\	810	1.00
80 m	10	8	218	0	8	8	8	9	10.4	Cla	0.50
90 m	80	Ø	100	0	08	0	08	0	10 -	50	0.50
Robel	Pole:	2			- //			X	(0	8 Ce	0 00

Robel Pole:

r.	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	15	13	(0	12	W/	Q	14	11	17	5
Ε	15	13	10	13	13	q	14	17	17	7
s	15	12	15	13	1-7	10	13	12		14
W	15	12	15	13	1.2	10	18		1 (14

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm

byd2 tbyd2a 2 100m