Daubenmire Datasheet

Date:

Observer: Ory

Recorder:

Team lead: _

Site:

Transect:

tbcr2a 2

bcr2

Round:

50m

	CSG	W56	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
	^	0	38	. 0				0		98	0,5
0 m	0							0	D	98	0.25
10 m	3	0	38	0				0	Λ	98	0.25
20 m	3	0	38	0				A	0	98	4
30 m	3	0	63	6				0	0	1.	1.0
40 m	16	0	63	0				0	U	98	1.0
50 m											
60 m) *				1						
70 m										-	
80 m					Armini						
90 m											

Robel Pole:

obel Po	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	20	1	2		(
E	1			1	(
s	1)	1	1,						
W	1	1,46			1					

Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm