* lots of . bd grass Daubenmire Datasheet Date: Observer: _

Round 50m

	CSG	WSG	Forbs	Milkweed	Common ramet #	S
Tec	am lead:	Kelser	1		was s	SPro
Red	corder:	Manne	1		100KS	likt

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter
0 m	3	6	0	05	8	d	0	7	16	a C	Depth (cm)
10 m	16	8	16	Ø	1	X			70	18	1.0
20 m	38	Cr	3	8	0	6			38	63	0.5
30 m	7	8	3	(X	0	//		16	16	78	0.25
40 m	10	B		8			-6	2	8	98	9.6
50 m	AA	AA	AAA	AA	ANT			0	1	98	2.5
60 m	1							0009		Vhr	
70 m					4						
80 m											
90 m											

	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	0	0	6	0	6	*				, , , , , , , , , , , , , , , , , , ,
Ε		6	3	0		5				
s	0	6	0		0	5				
W	1	0.4	Ò	0		3				

Categories: 0 - absent 1 - trace	3 + E%	1/ 50/ 050/				10.	
Categories: $\underline{0}$ - absent $ \underline{1}$ - trace $ \underline{0}$	3 - 1race-5%	<u>16</u> - 5%-25%	<u>38</u> - 25%-50%	<u>63</u> - 50%-75%	<u>86</u> - 75%	6-95% <u>98</u>	- 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm

MOSULINESS SUPPOSE INTESSE

nie1

tnie1a

1

Site:

Trans