Daubenmire Datasheet

Date:

Observer:

Recorder:

Team lead:

nkn2 Site: tnkn2a Transect: 3

Round: 100m

		W66	Forbs	Milkweed	Common	Swamp	Butterfly	Woody	Bare Ground	Leaf Litter	Litter Depth (cm)
* 11	CSG	WSG	Loros		ramet #	ramet #	ramet #	Species	110	28	0.25
O m	16	-3	38	0			1978年		10	38	0.25
10 m	) (	3	38	0					110	38	0.25
20 m	38	3	16	O					38	38	0.25
30 m	10	1(0	10	0					0	38	0.25
40 m	38	10	38	0			Tells /	0	10	20	
50 m	16	10	10	0				0	58	102	0.25
60 m	10	103	/12	0				0	10	03	0.00
70 m	16	6	63	6		1		0	14	63	10.25
80 m	16	16	160	0				0	16	38	0.05
90 m	16	16	63	0				0	16	38	0.20

Robel Pole:

Robel Po	le:				All has		/	70	80 m	90 m
	O m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	90 III	70 111
N	2	1	2	2	2		2	2	2	7
	0	1	3	)		7	2	7	2	A
Е	d			0	1	100	^		0	/
S	2	1	3	2	h 1	1	3	1	1	
		\ add	2			2	1	7	2	2
W	1	100		0	10					

Categories: 0 - absent 1 - trace 3 - trace-5% 16 - 5%-25% 38 - 25%-50% 63 - 50%-75% 86 - 75%-95% 98 - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm