

Daubermire Datasheet

Date: 7-18-18

Observer: Sydney Lizotte-Hall

Recorder: Sydney

Team lead: Sydney & Seth

Site:

cre1
tcre1a

Transect:

2

Round:

100m

* decent amount
of unspaced m.v.
grasses & Eriogonum
around 20-30 meters of
inside center plot.

	CS6	WS6	Forbs	Milkweed	Common ranet #	Swamp ranet #	Butterfly ranet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	16	3	86	0	0	0	0	0	16	63	0.25
10 m	38	14	38	0	0	0	0	0	38	38	0.25
20 m	3	3	63	0	0	0	0	0	16	86	4.0
30 m	16	1	63	0	0	0	0	0	16	63	4.0
40 m	5	1	63	0	0	0	0	0	16	63	2.0
50 m	16	1	63	0	0	0	0	0	16	38	0.25
60 m	1	1	63	0	0	0	0	0	16	86	2.0
70 m	0	0	63	0	0	0	0	0	1	86	1.5
80 m	0	0	86	0	0	0	0	0	16	86	2.0
90 m	1	0	86	0	0	0	0	0	0	86	1.0

Rebel Pole:

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	9	11	10	10	5	8	6	8	12	11
E	8	10	10	8	7	5	6	14	8	8
S	7	12	10	10	5	8	7	10	8	10
W	9	11	10	13	6	9	7	8	7	8

Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

CS6 - cool season grasses (including sedges)

WS6 - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm