Dauberimire Datasheet

Date: 5-18

Team lead:

Recorder: Observer:

Site: nkn2

Transect: tnkn2a

Round: 100m

	CS6	WS6	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody	Bare	Leaf Litter	Litter Depth (cm)
0	16	O	20	0	O	0	0	0	(0)	(23	
10 m	100	O	228	0	0	0	0	0		876	- 1
20 m	38	0	03	0	0	0	0	0	[/0	63	
30 m	38	0	20) -	0	<u></u>	0	0	(0)	00	- 1
40 m	76	0	Q.	S	0	0	0		1.(0	(0)	
50 m	× ×	0	8	0	0	0	0	0	N	00	
60 m	38	C	978	9	0	0	<u>ن</u>	0	160	63	- 1
70 m	6.0	90	16/	0	0	0	0	0	W	20	- 1
80 m	5	\bigcirc	20	0	0	0		0	-	200	
90 m	63	0	603	0	O	0	0	0	0)	20	,
Pohal Pola:											

5	2	
5	2	
٩	a	
		7
•	τ)
(3	
7	0	*
•	•	

	0	10 m	20 m	30 m	40 m	50 m	60 m		70 m	70 m 80 m
Z	5	0	0	0	0				7	01 4
m	12	1.1	70	10	ــــــــــــــــــــــــــــــــــــــ	7	7	1	1	_
S	رد	12	9.	QJ	ال	7	_		6	91
₹	J.	5	5	ď	8	7-1.	7	1	14	

CSG - cool season grasses (including sedges) Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm