Daubenmire Datasheet

Date:

Recorder: Observer:

Team lead:

Site: Transect: tbyd2a byd2

Round:

					Common	Swamp	Butterfly	Woody	Bare	Leaf	Litter
	600	WOG	rorbs	WIIKWeed	ramet #	ramet #	ramet #	Species	Ground	Litter	Depth (cm)
0 m	CN	Ø	8	Ø	Ø	Ø	Ø	Ø	CC	86	0.50
10 m	5	Ø	200	Ø	Ø	Ø	Ŋ	Ø	(N)	W	0,00
20 m	38	Ø	3	Ø	Ø	B	Ø	Ø	63	38	0.50
30 m	16	R	6	Ø	Ø	9	Ø	9	(N)	N ~	0,0
40 m	ZZ	B	6	Ø	9	Þ.	B	B	N N	60	0,50
50 m	25	P	6	Ø	þ	p	Ø	Ø	N	6W	8
60 m	160	R	200	Ø	P	b	ф	Ø	6 N	W X	· ?
70 m	N	P	W	þ	P	Ø	þ	B	0)1	26	0.50
80 m	200	É	22	Ø	B	4	Ø	Ø	63	28	0,25
3	OX	2	160	è,	Q	P)	Z	R	XX	N	4

	z	ш	S	8
0 m	4	(Q	4	4
10 m	4	R	4	7
20 m	2	4	t	4
30 m	٩	Q	9	0
40 m	0	۹	و_	٥
50 m	S	U	4	4
60 m	_0	P	٩	و
70 m	4	4	4	4
80 m	4	4	4	4
90 m	0	0	0	<u></u>

CSG - cool season grasses (including sedges) Categories: <u>0</u> - absent | <u>1</u> - trace | <u>3</u> - trace-5% | <u>16</u> - 5%-25% | <u>38</u> - 25%-50% | <u>63</u> - 50%-75% | <u>86</u> - 75%-95% | <u>98</u> - 95%-100%

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm

Bild rest plans