

Daubenmire Datasheet

Date: 7/15/20
 Observer: Kef
 Recorder:
 Team lead:

Site: Bayd 1
 Transect:
 Round:

	CSG	WSG	Forbs	Milkweed	Common ramet #	Swamp ramet #	Butterfly ramet #	Woody Species	Bare Ground	Leaf Litter	Litter Depth (cm)
0 m	38	0	63	3	1	0	0	0	16	86	2.5
10 m	3	0	86	0	0	0	0	0	16	86	2.0
20 m	38	0	63	16	1	0	1	0	16	86	3.0
30 m	16	0	86	0	0	0	0	0	16	86	2.5
40 m	16	0	86	0	0	0	0	0	16	86	2.5
50 m	86	0	38	0	0	0	0	0	16	86	3.0
60 m	63	0	38	0	0	0	0	0	16	86	3.0
70 m	38	0	63	0	0	0	0	0	16	86	3.0
80 m	38	0	63	3	2	0	0	0	16	86	3.0
90 m	38	0	63	3	1	0	0	0	16	86	3.0

Robel Pole:

	0 m	10 m	20 m	30 m	40 m	50 m	60 m	70 m	80 m	90 m
N	7	10	10	12	12	13	13	13	14	14
E	7	10	10	12	12	13	13	13	14	14
S	7	10	10	12	12	13	13	13	14	14
W	7	10	10	12	12	13	13	13	14	14

Categories: 0 - absent | 1 - trace | 3 - trace-5% | 16 - 5%-25% | 38 - 25%-50% | 63 - 50%-75% | 86 - 75%-95% | 98 - 95%-100%

CSG - cool season grasses (including sedges)

WSG - warm season grasses

Forbs - forbs not including milkweed

Wood - woody cover

Bare Ground - includes rocks, lichens, moss, branches, and cow pies

Leaf Litter (horizontal dead material)

Litter Depth - to nearest .25 cm