

```

setwd("C:\\Users\\it23761582\\Downloads\\IT23761582")
getwd()

data <- read.table("Exercise - Laptopsweights.txt", header = TRUE)
head(data)

weights <- data$weight
weights

#Q1
pop_mean <- mean(weights)
pop_mean

pop_sd <- sd(weights)
pop_sd

#Q2
set.seed(123)
sample_means <- numeric(25)
sample_sds <- numeric(25)

for (i in 1:25) {
  sample_data <- sample(weights, size = 6, replace = TRUE)
  sample_means[i] <- mean(sample_data)
  sample_sds[i] <- sd(sample_data)
  cat("Sample", i, "- Mean:", sample_means[i], " SD:", sample_sds[i], "\n")
}

#Q3
mean_of_sample_means <- mean(sample_means)
mean_of_sample_means

sd_of_sample_means <- sd(sample_means)
sd_of_sample_means

#comparing
popmn <- mean(weights)
popstd <- sd(weights)

samplemn <- mean(sample_means)
samplestd <- sd(sample_means) |

popmn
samplemn
popstd
samplestd

```

```

> setwd("C:\\Users\\it23761582\\Downloads\\IT23761582")
> getwd()
[1] "C:/Users/it23761582/Downloads/IT23761582"
> data <- read.table("Exercise - Laptopsweights.txt", header = TRUE)
> head(data)
  weight.kg.
1      2.46
2      2.45
3      2.47
4      2.71
5      2.46
6      2.05
> weights <- data$weight
> weights
[1] 2.46 2.45 2.47 2.71 2.46 2.05 2.60 2.42 2.43 2.53 2.57 2.85 2.70
[14] 2.53 2.28 2.20 2.57 2.89 2.51 2.47 2.66 2.06 2.41 2.65 2.76 2.43
[27] 2.61 2.57 2.73 2.17 2.67 2.05 1.71 2.32 2.23 2.76 2.70 2.13 2.75
[40] 2.20
> #Q1
> pop_mean <- mean(weights)
> pop_mean
[1] 2.468
> pop_sd <- sd(weights)
> pop_sd
[1] 0.2561069
> #Q2
> set.seed(123)
> sample_means <- numeric(25)
> sample_sds <- numeric(25)
> for (i in 1:25) {
+   sample_data <- sample(weights, size = 6, replace = TRUE)
+   sample_means[i] <- mean(sample_data)
+   sample_sds[i] <- sd(sample_data)
+   cat("Sample", i, "- Mean:", sample_means[i], " SD:", sample_sds[i], "\n")
+ }

```

```

+ }
Sample 1 - Mean: 2.53 SD: 0.1513935
Sample 2 - Mean: 2.573333 SD: 0.1191078
Sample 3 - Mean: 2.473333 SD: 0.1718914
Sample 4 - Mean: 2.591667 SD: 0.1345239
Sample 5 - Mean: 2.456667 SD: 0.2749303
Sample 6 - Mean: 2.401667 SD: 0.254434
Sample 7 - Mean: 2.59 SD: 0.2167026
Sample 8 - Mean: 2.466667 SD: 0.4530195
Sample 9 - Mean: 2.401667 SD: 0.2230172
Sample 10 - Mean: 2.335 SD: 0.3237746
Sample 11 - Mean: 2.586667 SD: 0.1706068
Sample 12 - Mean: 2.378333 SD: 0.3235686
Sample 13 - Mean: 2.381667 SD: 0.2993604
Sample 14 - Mean: 2.465 SD: 0.2314951
Sample 15 - Mean: 2.485 SD: 0.1745566
Sample 16 - Mean: 2.451667 SD: 0.2762909
Sample 17 - Mean: 2.385 SD: 0.2042303
Sample 18 - Mean: 2.338333 SD: 0.2436733
Sample 19 - Mean: 2.428333 SD: 0.2481465
Sample 20 - Mean: 2.551667 SD: 0.2654367
Sample 21 - Mean: 2.538333 SD: 0.1708118
Sample 22 - Mean: 2.466667 SD: 0.2451666
Sample 23 - Mean: 2.47 SD: 0.2405826
Sample 24 - Mean: 2.448333 SD: 0.279243
Sample 25 - Mean: 2.475 SD: 0.2358601
> #Q2
> set.seed(123)
> sample_means <- numeric(25)
> sample_sds <- numeric(25)
> for (i in 1:25) {
+   sample_data <- sample(weights, size = 6, replace = TRUE)
+   sample_means[i] <- mean(sample_data)
+   sample_sds[i] <- sd(sample_data)


```

```

+   cat("Sample", i, "- Mean:", sample_means[i], " SD:", sample_sds[i], "\n")
+ }
Sample 1 - Mean: 2.53 SD: 0.1513935
Sample 2 - Mean: 2.573333 SD: 0.1191078
Sample 3 - Mean: 2.473333 SD: 0.1718914
Sample 4 - Mean: 2.591667 SD: 0.1345239
Sample 5 - Mean: 2.456667 SD: 0.2749303
Sample 6 - Mean: 2.401667 SD: 0.254434
Sample 7 - Mean: 2.59 SD: 0.2167026
Sample 8 - Mean: 2.466667 SD: 0.4530195
Sample 9 - Mean: 2.401667 SD: 0.2230172
Sample 10 - Mean: 2.335 SD: 0.3237746
Sample 11 - Mean: 2.586667 SD: 0.1706068
Sample 12 - Mean: 2.378333 SD: 0.3235686
Sample 13 - Mean: 2.381667 SD: 0.2993604
Sample 14 - Mean: 2.465 SD: 0.2314951
Sample 15 - Mean: 2.485 SD: 0.1745566
Sample 16 - Mean: 2.451667 SD: 0.2762909
Sample 17 - Mean: 2.385 SD: 0.2042303
Sample 18 - Mean: 2.338333 SD: 0.2436733
Sample 19 - Mean: 2.428333 SD: 0.2481465
Sample 20 - Mean: 2.551667 SD: 0.2654367
Sample 21 - Mean: 2.538333 SD: 0.1708118
Sample 22 - Mean: 2.466667 SD: 0.2451666
Sample 23 - Mean: 2.47 SD: 0.2405826
Sample 24 - Mean: 2.448333 SD: 0.279243
Sample 25 - Mean: 2.475 SD: 0.2358601
> #Q3
> mean_of_sample_means <- mean(sample_means)
> mean_of_sample_means
[1] 2.4668
> sd_of_sample_means <- sd(sample_means)
> sd_of_sample_means
[1] 0.07624874

> #Q3
> mean_of_sample_means <- mean(sample_means)
> mean_of_sample_means
[1] 2.4668
> sd_of_sample_means <- sd(sample_means)
> sd_of_sample_means
[1] 0.07624874
> #comparing
> popmn <- mean(weights)
> popstd <- sd(weights)
> samplemn <- mean(sample_means)
> samplestd <- sd(sample_means)
> popmn
[1] 2.468
> samplemn
[1] 2.4668
> popstd
[1] 0.2561069
> samplestd
[1] 0.07624874
> |

```

data	40 obs. of 1 variable	
values		
i	25L	
mean_of_sample_me...	2.4668	
pop_mean	2.468	
pop_sd	0.256106948813907	
popmn	2.468	
popstd	0.256106948813907	
sample_data	num [1:6] 2.57 2.42 2.66 2.45 2.7 2.05	
sample_means	num [1:25] 2.53 2.57 2.47 2.59 2.46 ...	
sample_sds	num [1:25] 0.151 0.119 0.172 0.135 0.275 ...	
samplemn	2.4668	
samplestd	0.0762487401231677	
sd_of_sample_means	0.0762487401231677	
weights	num [1:40] 2.46 2.45 2.47 2.71 2.46 2.05 2.6 2.42 ...	