

IT24100636– Fernando T.M.I.U

IT2120 - Probability and Statistics Lab Sheet 05


01)

```
setwd("C:\\Users\\imasha\\Desktop\\PS_IT24100636")
Delivery_Times <- read.table("Exercise - Lab 05.txt", header = TRUE)
head(Delivery_Times)
```

```
> setwd("C:\\Users\\imasha\\Desktop\\PS_IT24100636")
> Delivery_Times <- read.table("Exercise - Lab 05.txt", header = TRUE)
> head(Delivery_Times)
```

02)

```
#Question 02
breaks <- seq(20, 70, length.out = 10)
hist(Delivery_Times$Delivery_Time,
     breaks = breaks,
     right = FALSE,
     main = "Histogram of Delivery Times",
     xlab = "Delivery Time (minutes)",
     col = "green",
     border = "black")
```



03.

```
#Question 03
# The distribution of delivery time is unimodal and slightly right-skewed.
# Most deliveries occur around 40 minutes, while a few deliveries take much longer,
# creating a tail on the right side of the histogram.
```

04.

```

#Question 04
histogram <- hist(Delivery_Times$Delivery_Time,
                  breaks = seq(20, 70, length.out = 10),
                  right = FALSE,
                  plot = FALSE)

breaks <- histogram$breaks
freq <- histogram$counts

classes <- c()
for(i in 1:(length(breaks) - 1)) {
  classes[i] <- paste0("[", breaks[i], ", ", breaks[i+1], ")")
}

freq_table <- cbind(Class_Interval = classes, Frequency = freq)
print(freq_table)

cum_freq <- cumsum(freq)

cum_freq_with0 <- c(0, cum_freq)

plot(breaks, cum_freq_with0, type = "o",
     main = "Cumulative Frequency Polygon (Ogive)",
     xlab = "Delivery Time (minutes)",
     ylab = "Cumulative Frequency",
     col = "red", pch = 16)

cum_table <- cbind(Upper_Class = breaks, Cum_Freq = cum_freq_with0)
print(cum_table)

```

(Top Level) ⚙

le Terminal Background Jobs

```
Console Terminal x Background Jobs x
R 4.5.1 · C:/Users/vimuk/OneDrive/Desktop/IT24103506/ ↗

> #Question 04
> histogram <- hist(Delivery_Times$Delivery_Time,
+                   breaks = seq(20, 70, length.out = 10),
+                   right = FALSE,
+                   plot = FALSE)
> breaks <- histogram$breaks
> freq <- histogram$counts
> classes <- c()
> for(i in 1:(length(breaks) - 1)) {
+   classes[i] <- paste0("[", breaks[i], ", ", breaks[i+1], ")")
+ }
> freq_table <- cbind(Class_Interval = classes, Frequency = freq)
> print(freq_table)
      Class_Interval      Frequency
[1,] "[20, 25.5555555555556)"      "2"
[2,] "[25.5555555555556, 31.1111111111111)" "3"
[3,] "[31.1111111111111, 36.6666666666667)" "6"
[4,] "[36.6666666666667, 42.2222222222222)" "9"
[5,] "[42.2222222222222, 47.7777777777778)" "6"
[6,] "[47.7777777777778, 53.3333333333333)" "3"
[7,] "[53.3333333333333, 58.8888888888889)" "6"
[8,] "[58.8888888888889, 64.4444444444444)" "3"
[9,] "[64.4444444444444, 70)"      "2"
[10,] "[70, 75.5555555555556)"      "0"
> cum_freq <- cumsum(freq)
> cum_freq_with0 <- c(0, cum_freq)
> plot(breaks, cum_freq_with0, type = "o",
+      main = "Cumulative Frequency Polygon (Ogive)",
+      xlab = "Delivery Time (minutes)",
+      ylab = "Cumulative Frequency",
+      col = "red", pch = 16)
> cum_table <- cbind(Upper_Class = breaks, Cum_Freq = cum_freq_with0)
> print(cum_table)
      Upper_Class Cum_Freq
[1,] 20.00000      0
[2,] 25.55556      2
[3,] 31.11111      5
[4,] 36.66667     11
[5,] 42.22222     20
[6,] 47.77778     26
[7,] 53.33333     29
[8,] 58.88889     35
[9,] 64.44444     38
[10,] 70.00000     40
```

