

## IT24100636 Fernando T.M.I.U.

### IT2120 - Probability and Statistics | Lab Sheet 08

```
setwd("C:\\Users\\imasha\\Desktop\\IT24100636_PS_08")  
  
data<-read.table("Exercise - LaptopsWeights.txt",header = TRUE)  
fix(data)  
attach(data)
```

```
> setwd("C:\\Users\\imasha\\Desktop\\IT24100636_PS_08")  
> data<-read.table("Exercise - LaptopsWeights.txt",header = TRUE)  
> fix(data)  
attach(data)
```

Data Editor						
File Edit Help						
	Weight.kg.	var2	var3	var4	var5	var6
1	2.46					
2	2.45					
3	2.47					
4	2.71					
5	2.46					
6	2.05					
7	2.6					
8	2.42					
9	2.43					
10	2.53					
11	2.57					
12	2.85					
13	2.7					
14	2.53					
15	2.28					
16	2.2					
17	2.57					
18	2.89					
19	2.51					

01)

```
7 #Question01
8 popmn<-mean(weight.kg.)
9 popmn
10 popsd<-sd(weight.kg.)
11 popsd
12
> #Question01
> popmn<-mean(weight.kg.)
> popmn
[1] 2.468
> popsd<-sd(weight.kg.)
> popsd
[1] 0.2561069
```

02)

```
#Question02
samples<-c()
n<-c()
for(i in 1:25){
  s<-sample(weight.kg.,6,replace = TRUE)
  samples<-cbind(samples,s)
  n<-c(n,paste('S',i))
}
colnames(samples)=n
s.means<-apply(samples,2,mean)
s.means
s.sd<-apply(samples,2,sd)
s.sd
```

```

> #Question02
> samples<-c()
> n<-c()
> for(i in 1:25){
+   s<-sample(weight.kg.,6,replace = TRUE)
+   samples<-cbind(samples,s)
+   n<-c(n,paste('S',i))
+ }
> colnames(samples)=n
> s.means<-apply(samples,2,mean)
> s.means
      S 1      S 2      S 3      S 4      S 5      S 6      S 7      S 8      S 9      S 10      S 11
2.478333 2.556667 2.320000 2.553333 2.526667 2.540000 2.506667 2.128333 2.580000 2.495000 2.550000
      S 12      S 13      S 14      S 15      S 16      S 17      S 18      S 19      S 20      S 21      S 22
2.436667 2.458333 2.581667 2.620000 2.366667 2.356667 2.605000 2.446667 2.371667 2.208333 2.460000
      S 23      S 24      S 25
2.328333 2.428333 2.388333
> s.sd<-apply(samples,2,sd)
> s.sd
      S 1      S 2      S 3      S 4      S 5      S 6      S 7      S 8      S 9      S 10
0.2133932 0.3695763 0.3628774 0.2434475 0.1434805 0.2587663 0.2597435 0.2698456 0.1113553 0.1504327
      S 11      S 12      S 13      S 14      S 15      S 16      S 17      S 18      S 19      S 20
0.2589208 0.2331237 0.1888297 0.1803792 0.1801111 0.3977269 0.2499333 0.1663430 0.3915440 0.4214934
      S 21      S 22      S 23      S 24      S 25
0.3213980 0.4166533 0.2655874 0.2696974 0.1646107

```

03)

```

#Question03
#calculate the mean and standard deviation of the 25 sample means
samplemean<-mean(s.means)
samplemean
samplesd<-sd(s.means)
samplesd

#state therelationship of them with true mean and true standard deviation
popmn
samplemean

truesd=popsd/sqrt(6)
samplesd

```

```

> #Question03
> #calculate the mean and standard deviation of the 25 sample means
> samplemean<-mean(s.means)
> samplemean
[1] 2.451667
> samplesd<-sd(s.means)
> samplesd
[1] 0.1223317
> #state therelationship of them with true mean and true standard deviation
> popmn
[1] 2.468
> samplemean
[1] 2.451667
> truesd=popsd/sqrt(6)
> samplesd
[1] 0.1223317

```

R ▾

Global Environment ▾