


Probability and Statistics - IT2120

Labsheet 08

Dilsandi P.D.S



	Nicotine	var2	var3	var4	var5	var6	var7
1	1.09						
2	1.74						
3	1.58						
4	2.11						
5	1.64						
6	1.79						
7	1.37						
8	1.75						
9	1.92						
10	1.47						
11	2.03						
12	1.86						
13	0.72						
14	2.46						
15	1.93						
16	1.63						
17	2.31						
18	1.97						
19	1.7						

```

setwd("E:\\IT24101734")
getwd()

# Read the data file
weights <- read.table("Exercise - Laptopsweights.txt", header = TRUE)
fix(weights)
attach(weights)

#Q1
popmn<-mean(weight.kg.)
popmn
popsd<-sd(weight.kg.)
popsd

# Q2
samples<-c()
n<-c()
for(i in 1:25){
  s<-sample(weight.kg.,6,replace = TRUE)
  samples<-cbind(samples,s)
  n<-c(n,paste('s',i))
}
colnames(samples)=n
s.means<-apply(samples,2,mean)
s.means
s.sd<-apply(samples,2,sd)
s.sd

#Q3
#calculate the mean and standard deviation of the 25 sample means
samplemean<-mean(s.means)
samplemean
samplesd<-sd(s.means)
samplesd

#state the relationship of them with true mean and true standard deviation
popmn
samplemean

truesd=popsd/sqrt(6)
samplesd

```

Q1)

```
> setwd("E:\\IT24101734")
> getwd()
[1] "E:/IT24101734"
> # Read the data file
> weights <- read.table("Exercise - Laptopsweights.txt", header = TRUE)
> fix(weights)
> attach(weights)

> #Q1
> popmn<-mean(weight.kg.)
> popmn
[1] 2.468
> popsd<-sd(weight.kg.)
> popsd
[1] 0.2561069
>
```

Q2)

```
> # Q2
> samples<-c()
> n<-c()
> for(i in 1:25){
+   s<-sample(weight.kg.,6,replace = TRUE)
+   samples<-cbind(samples,s)
+   n<-c(n,paste('s',i))
+ }
> colnames(samples)=n
> s.means<-apply(samples,2,mean)
> s.means
      s 1      s 2      s 3      s 4      s 5
2.526667 2.278333 2.606667 2.388333 2.248333
      s 6      s 7      s 8      s 9      s 10
2.171667 2.446667 2.496667 2.488333 2.278333
      s 11      s 12      s 13      s 14      s 15
2.645000 2.365000 2.575000 2.416667 2.551667
      s 16      s 17      s 18      s 19      s 20
2.510000 2.585000 2.391667 2.573333 2.485000
      s 21      s 22      s 23      s 24      s 25
2.635000 2.376667 2.581667 2.370000 2.401667
> s.sd<-apply(samples,2,sd)
> s.sd
      s 1      s 2      s 3      s 4
0.2482472 0.2951892 0.2098253 0.2934905
      s 5      s 6      s 7      s 8
0.3485063 0.3949895 0.1923192 0.1214359
      s 9      s 10      s 11      s 12
0.2056615 0.3287806 0.1201249 0.1961377
      s 13      s 14      s 15      s 16
0.2476893 0.2915933 0.2208544 0.2551862
      s 17      s 18      s 19      s 20
0.1232477 0.2430158 0.1415156 0.2746452
      s 21      s 22      s 23      s 24
0.1261348 0.2986414 0.1259233 0.4049691
      s 25
0.3162541
```

Q3)

```
> #Q3
> #calculate the mean and standard deviation of the 25 sample means
> samplemean<-mean(s.means)
> samplemean
[1] 2.455733
> samplesd<-sd(s.means)
> samplesd
[1] 0.1280951
>
> #state the relationship of them with true mean and true standard deviation
> popmn
[1] 2.468
> samplemean
[1] 2.455733
>
> truesd=popstd/sqrt(6)
> samplesd
[1] 0.1280951
```