


Probability and Statistics - IT2120

Labsheet 08

Nandanayaka H.P.N.M



	Nicotine	var2	var3	var4	var5	var6	var7
1	1.09						
2	1.74						
3	1.58						
4	2.11						
5	1.64						
6	1.79						
7	1.37						
8	1.75						
9	1.92						
10	1.47						
11	2.03						
12	1.86						
13	0.72						
14	2.46						
15	1.93						
16	1.63						
17	2.31						
18	1.97						
19	1.7						

```
IT24101981.R x
Source on Save Run
1 setwd("c:\\Users\\malsh\\Desktop\\IT24101981")
2 getwd()
3
4 # Read the data file
5 weights <- read.table("Exercise - Laptopsweights.txt", header = TRUE)
6 fix(weights)
7 attach(weights)
8
9 #Q1
10 popmn<-mean(weight.kg.)
11 popmn
12 popsd<-sd(weight.kg.)
13 popsd
14
15 # Q2
16 samples<-c()
17 n<-c()
18 for(i in 1:25){
19   s<-sample(weight.kg.,6,replace = TRUE)
20   samples<-cbind(samples,s)
21   n<-c(n,paste('s',i))
22 }
23 colnames(samples)=n
24 s.means<-apply(samples,2,mean)
25 s.means
26 s.sd<-apply(samples,2,sd)
27 s.sd
28
29 #Q3
30 #calculate the mean and standard deviation of the 25 sample means
31 samplemean<-mean(s.means)
32 samplemean
33 samplesd<-sd(s.means)
34 samplesd
35
36 #state the relationship of them with true mean and true standard deviation
37 popmn
38 samplemean
39
40 truesd=popsd/sqrt(6)
41 samplesd
42
43
44
45
46
```

Q1)

```
> setwd("c:\\Users\\malsh\\Desktop\\IT24101981")
> getwd()
[1] "C:/Users/malsh/Desktop/IT24101981"
> weights <- read.table("Exercise - Laptopsweights.txt", header = TRUE)
> fix(weights)
> attach(weights)
> popmn<-mean(weight.kg.)
> popmn
[1] 2.468
> popsd<-sd(weight.kg.)
> popsd
[1] 0.2561069
> popmn<-mean(weight.kg.)
> popmn
```

Q2)

```
> # Q2
> samples<-c()
> n<-c()
> for(i in 1:25){
+   s<-sample(weight.kg.,6,replace = TRUE)
+   samples<-cbind(samples,s)
+   n<-c(n,paste('s',i))
+ }
> colnames(samples)=n
> s.means<-apply(samples,2,mean)
> s.means
      s 1      s 2      s 3      s 4      s 5
2.526667 2.278333 2.606667 2.388333 2.248333
      s 6      s 7      s 8      s 9      s 10
2.171667 2.446667 2.496667 2.488333 2.278333
      s 11     s 12     s 13     s 14     s 15
2.645000 2.365000 2.575000 2.416667 2.551667
      s 16     s 17     s 18     s 19     s 20
2.510000 2.585000 2.391667 2.573333 2.485000
      s 21     s 22     s 23     s 24     s 25
2.635000 2.376667 2.581667 2.370000 2.401667
> s.sd<-apply(samples,2,sd)
> s.sd
      s 1      s 2      s 3      s 4
0.2482472 0.2951892 0.2098253 0.2934905
      s 5      s 6      s 7      s 8
0.3485063 0.3949895 0.1923192 0.1214359
      s 9      s 10     s 11     s 12
0.2056615 0.3287806 0.1201249 0.1961377
      s 13     s 14     s 15     s 16
0.2476893 0.2915933 0.2208544 0.2551862
      s 17     s 18     s 19     s 20
0.1232477 0.2430158 0.1415156 0.2746452
      s 21     s 22     s 23     s 24
0.1261348 0.2986414 0.1259233 0.4049691
      s 25
0.3162541
```

Q3)

```
> #Q3
> #calculate the mean and standard deviation of the 25 sample means
> samplemean<-mean(s.means)
> samplemean
[1] 2.455733
> samplesd<-sd(s.means)
> samplesd
[1] 0.1280951
>
> #state the relationship of them with true mean and true standard deviation
> popmn
[1] 2.468
> samplemean
[1] 2.455733
>
> truesd=popsd/sqrt(6)
> samplesd
[1] 0.1280951
```