Dolamulla H.D.K.P.D

IT24103522

Probability and Statistics

1. Assume that the time taken to bake a batch of cookies is normally distributed with mean 45 minutes and standard deviation 2 minutes.

   i. Generate a random sample of size 25 for the baking time.

```
1  setwd("C:/Users/ROG/Desktop/IT24103522")
2
3  #(i)
4  set.seed(123)
5  bake_time <- rnorm(25, mean = 45, sd = 2)
6  bake_time
> setwd("C:/Users/ROG/Desktop/IT24103522")
> #(i)
> set.seed(123)
> bake_time <- rnorm(25, mean = 45, sd = 2)
> bake_time
 [1] 43.87905 44.53965 48.11742 45.14102 45.25858 48.43013 45.92183 42.46988 43.62629 44.10
[11] 47.44816 45.71963 45.80154 45.22137 43.88832 48.57383 45.99570 41.06677 46.40271 44.05
[21] 42.86435 44.56405 42.94799 43.54222 43.74992
```

   ii. Test whether the average baking time is less than 46 minutes at a 5% level of significance.

```
 9  #(ii)
10  t.test(bake_time, mu = 46, alternative = "less",conf.level = 0.95)
> #(ii)
> t.test(bake_time, mu = 46, alternative = "less",conf.level = 0.95)

        One Sample t-test

data:  bake_time
t = -2.8167, df = 24, p-value = 0.004776
alternative hypothesis: true mean is less than 46
95 percent confidence interval:
     -Inf 45.58124
sample estimates:
mean of x
 44.93334
```