



Faculty of Computing

Year 2 Semester 1 (2025)

IT2120 - Probability and Statistics

Lab Sheet 08

Exercise

1. Calculate the population mean and population standard deviation of the laptop bag weights.

	Weight.kg.	var2	var3	var4	var5	var6
1	2.46					
2	2.45					
3	2.47					
4	2.71					
5	2.46					
6	2.05					
7	2.6					
8	2.42					
9	2.43					
10	2.53					
11	2.57					
12	2.85					
13	2.7					
14	2.53					
15	2.28					
16	2.2					
17	2.57					
18	2.89					
19	2.51					

```
> setwd("C:\\Users\\acer\\OneDrive\\Desktop\\IT24104092")
>
> getwd()
[1] "C:/Users/acer/OneDrive/Desktop/IT24104092"
>
> # Read the data file
> weights <- read.table("Exercise - Laptopsweights.txt", header = TRUE)
> fix(weights)
> attach(weights)
```

```

> #Q1
> popmn<-mean(weight.kg.)
> popmn
[1] 2.468
> popsd<-sd(weight.kg.)
> popsd
[1] 0.2561069

```

2. Draw 25 random samples of size 6 (with replacement) and calculate the sample mean and sample standard deviation for each sample.

```

> # Q2
> samples<-c()
> n<-c()
> for(i in 1:25){
+   s<-sample(weight.kg.,6,replace = TRUE)
+   samples<-cbind(samples,s)
+   n<-c(n,paste('s',i))
+ }
> colnames(samples)=n
> s.means<-apply(samples,2,mean)
> s.means
      s 1      s 2      s 3      s 4      s 5      s 6
2.336667 2.460000 2.458333 2.696667 2.456667 2.513333
      s 7      s 8      s 9      s 10     s 11     s 12
2.358333 2.466667 2.425000 2.591667 2.471667 2.520000
      s 13     s 14     s 15     s 16     s 17     s 18
2.288333 2.446667 2.378333 2.463333 2.236667 2.463333
      s 19     s 20     s 21     s 22     s 23     s 24
2.456667 2.603333 2.528333 2.385000 2.561667 2.420000
      s 25
2.401667
- - - - -
> s.sd<-apply(samples,2,sd)
> s.sd
      s 1      s 2      s 3      s 4      s 5
0.2280058 0.3312401 0.1682161 0.1602082 0.2307957
      s 6      s 7      s 8      s 9      s 10
0.3731309 0.1979310 0.1467878 0.2137054 0.1333292
      s 11     s 12     s 13     s 14     s 15
0.3539727 0.2813539 0.3257248 0.2293178 0.3361200
      s 16     s 17     s 18     s 19     s 20
0.2196057 0.3023684 0.2166718 0.2967603 0.1366260
      s 21     s 22     s 23     s 24     s 25
0.2614128 0.1803053 0.1859480 0.2319483 0.2540407

```

3. Calculate the mean and standard deviation of the 25 sample means and state the relationship of them with true mean and true standard deviation.

```
> #Q3
> #calculate the mean and standard deviation of the 25 sample means
> samplemean<-mean(s.means)
> samplemean
[1] 2.455533
> samplesd<-sd(s.means)
> samplesd
[1] 0.09972151
> #state the relationship of them with true mean and true standard deviation
> popmn
[1] 2.468
> samplemean
[1] 2.455533
> truesd=popsd/sqrt(6)
> samplesd
[1] 0.09972151
```