

IT24104102

Amarakeerthi H.K.K.U

PSLabsheet 07

Q1)

```
1 setwd("/Users/kavinduumayanga/Desktop/IT24104102_Lab_07")
2 getwd()
3
4 #Question 01
5 #Uniform Distribution
6 #Here, random variable X has follows distribution with a=0 and b=40.
7 #p(10<X<25)=p(x<=25)-p(X<=10)
8 punif(25,min=0,max=40,lower.tail = TRUE)-punif(10,min=0,max=40,lower.tail = TRUE)
```

```
> setwd("/Users/kavinduumayanga/Desktop/IT24104102_Lab_07")
> getwd()
[1] "/Users/kavinduumayanga/Desktop/IT24104102_Lab_07"
> #Question 01
> #Uniform Distribution
> #Here, random variable X has follows distribution with a=0 and b=40.
> #p(10<X<25)=p(x<=25)-p(X<=10)
> punif(25,min=0,max=40,lower.tail = TRUE)-punif(10,min=0,max=40,lower.tail = TRUE)
[1] 0.375
```

Q2)

```
10 #Question 02
11 #Exponential Distribution
12 #Here, random variable X has exponential distribution with lamda=0.33
13 #p(X<=2)
14 pexp(2,rate = 1/3,lower.tail = TRUE)
```

```
> #Question 02
> #Exponential Distribution
> #Here, random variable X has exponential distribution with lamda=0.33
> #p(X<=2)
> pexp(2,rate = 1/3,lower.tail = TRUE)
[1] 0.4865829
```

Q3)

```
16 #Question 03
17 #Normal Distribution
18 #Here, random variable X has normal distribution with mean=100 and standard deviation=15.
19
20 #Part 01
21 #p(X>130)
22 pnorm(130,mean=100,sd=15,lower.tail = FALSE)
23 #or
24 1-pnorm(130,mean=100,sd=15,lower.tail = TRUE)
25
```

```
> #Question 03
> #Normal Distribution
> #Here, random variable X has normal distribution with mean=100 and standard deviation=15.
>
> #Part 01
> #p(X>130)
> pnorm(130,mean=100,sd=15,lower.tail = FALSE)
[1] 0.02275013
> #or
> 1-pnorm(130,mean=100,sd=15,lower.tail = TRUE)
[1] 0.02275013
>
```

```
26 #Part 02
27 #p(X=95/100)=0.95
28 qnorm(0.95,mean=100,sd=15,lower.tail = TRUE)
```

```
> #Part 02
> #p(X=95/100)=0.95
> qnorm(0.95,mean=100,sd=15,lower.tail = TRUE)
[1] 124.6728
```