

Ashfak

IT24100326

(1)

```
> setwd("C:\\Users\\IT24100326\\Desktop\\IT24100326")
> branch_data <- read.table("Exercise.txt", header = TRUE, sep = ",")
> str(branch_data)

> branch_data <- read.table("Exercise.txt", header = TRUE, sep = ",")
> str(branch_data)
'data.frame': 30 obs. of 4 variables:
 $ Branch      : int  1 2 3 4 5 6 7 8 9 10 ...
 $ Sales_X1    : num  3.4 4.1 2.8 5 3.7 4.5 3 4.9 3.2 2.5 ...
 $ Advertising_X2: int 120 150 90 200 110 175 95 185 105 80 ...
 $ Years_X3    : int  4 7 3 10 5 6 2 9 4 1 ...
```

(2)

branch-Categorical(nominal)


Sales_X1-Quantitative (Ratio scale, continuous)

Advertising_X2-Quantitative (Ratio scale, continuous)

Years_X3-Quantitative (Ratio scale, discrete – whole years)

(3)

```
boxplot(branch_data$Sales, main = "Boxplot of Sales",ylab = "sales")
> boxplot(branch_data$sales, main = "Boxplot of Sales",ylab = "sales")
```



(4)

```
quantile(branch_data$Advertising_X2)
summary(branch_data$Advertising_X2)
IQR(branch_data$Advertising_X2)

> quantile(branch_data$Advertising_X2)
  0%   25%   50%   75%  100%
80.00 101.25 132.50 158.75 210.00
> summary(branch_data$Advertising_X2)
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  80.0   101.2   132.5   134.8   158.8   210.0
> IQR(branch_data$Advertising_X2)
[1] 57.5
```

(5)

```
find_outliers <- function(x) {
  Q1 <- quantile(x, 0.25)
  Q3 <- quantile(x, 0.75)
  IQR_value <- Q3 - Q1
  lower_bound <- Q1 - 1.5 * IQR_value
  upper_bound <- Q3 + 1.5 * IQR_value
  return(x[x < lower_bound | x > upper_bound])
}

find_outliers(branch_data$Years)

> find_outliers <- function(x) {
+   Q1 <- quantile(x, 0.25)
+   Q3 <- quantile(x, 0.75)
+   IQR_value <- Q3 - Q1
+   lower_bound <- Q1 - 1.5 * IQR_value
+   upper_bound <- Q3 + 1.5 * IQR_value
+   return(x[x < lower_bound | x > upper_bound])
+ }
>
> find_outliers(branch_data$Years)
integer(0)
```