

Student Id – IT24200343

Name – AVEKSHIKA A.H.E

Lab sheet 10

1

```
> setwd("C:\\Users\\Lio Computers\\Desktop\\IT24100108")
> observed <- c(120, 95, 85, 100)
> prob <- c(.25, .25, .25, .25)
> chisq.test(x=observed, p=prob)
```

```
      Chi-squared test for given
      probabilities
```

```
data:  observed
X-squared = 6.5, df = 3, p-value =
0.08966
```

2

```

> file_path <- "http://www.sthda.com/sthda/RDoc/data/housetasks.txt"
> file_path <- "http://www.sthda.com/sthda/RDoc/data/housetasks.txt"
>
> housetasks <- read.delim(file_path, row.names = 1)
> housetasks
      wife Alternating Husband Jointly
Laundry   156         14        2        4
Main_meal  124         20        5        4
Dinner     77         11        7       13
Breakfast  82         36       15        7
Tidying    53         11        1       57
Dishes     32         24        4       53
Shopping   33         23        9       55
Official   12         46       23       15
Driving    10         51       75        3
Finances   13         13       21       66
Insurance   8          1       53       77
Repairs     0          3      160        2
Holidays   0          1        6     153
> chisq <- chisq.test(housetasks)
> chisq

      Pearson's Chi-squared test

data:  housetasks
X-squared = 1944.5, df = 36,
p-value < 2.2e-16

> |

```

3

```

> ##Q3
> #Consider 25% level of significant for the class
> #Rejection Region: if the p value for the test is less than 0.25,
> #Reject the null hypothesis at 25% level of significant
> #P value for the test got as 2.2e-16
> |

```