
Rewiring the Field: A Formal Account of Neurodivergence and Trauma

[T]-Theory Volume: Psychiatry, ASD, and Trauma

Alistair Johnson

Independent Researcher, Zurich, Switzerland

ORCID: [0009-0007-2194-0850](https://orcid.org/0009-0007-2194-0850)

Contents

Introduction: Not Broken, Modified	4
The ASC Operator: High Beta, Narrow Tongue	4
The CPTSD Operator: Non-Ergodic, EC Decoupled	5
Why They Co-Occur	5
Clinical Implications: Therapeutic Order	6
The Lean 4 Operators	6
What This Book Offers Clinicians and Neurodivergent Readers	6
Introduction	8
Background	8
Hopfield Networks 1982 (Classical)	8
Modern Hopfield Networks 2020 (Exponential)	9
The Missing Limbic Layer	9
The CEMI Field as a Runtime Parameter	10
The Two Coupling Equations	10
The FM-HN Update Rule	11
The Correspondence Principle	11
Falsifiability: The Reachability Trap Protocol	12
Neurodivergent Operator Modifications	12
ADHD Operator	12
Autism Spectrum Condition Operator	12
Complex PTSD Operator	13
Discussion	13
Why the 1982 and 2020 Networks Are Both Right	13
Relation to QUANT-EXP-1	14
Lean 4 Verification	14
Conclusion	14
1. Introduction	16
2. Terminology and the Pre-Verbal Manifold	17
3. The Case	18
3.1 Substrate (familial loading)	18
3.2 Acute pre-verbal insult	18
3.3 Attachment environment	19
3.4 Institutional environment	19
3.5 Identity and racialisation	20
3.6 Adult trajectory (compressed)	20
4. The Five Literatures	20
5. The Soma-Field Reading	21
6. The Twice-Exceptional Cognitive Profile	23

7. The Adult Trajectory as Basin Transitions	23
7.1 Basin transitions	24
7.2 Inclusion of the December 2024 event	25
7.3 The reorganisation phase	25
8. Exhibit A: A Public SEN-Policy Document	25
9. Ten Testable Predictions	27
10. Limitations, Replication Ledger, and Author Disclosures	28
11. Conclusion and Policy Line	28
A Note on Method	30
Introduction: The Inadequacy of Existing Maps	30
Background	31
Lived Experience as a Research Position	31
The Body-Mind Problem in Clinical Practice	32
The Felt Sense and Sub-Perceptual Emotion	32
Quantum Field Theory: Structure, Not Metaphor	32
Hopfield Networks and the Energy Function	33
The Soma-Field Model	33
Emotions as a Persistent Wave Field	33
The Perception Threshold	34
The Interaction of Emotional Modes	36
The Energy Landscape	36
The Hopfield Energy Function	36
Attractor States: Fight, Flight, Freeze, and Regulated Calm	36
Dissonance and Resolution	38
The Neurodivergent Field: ASD, ADHD, and C-PTSD as Operator Modifications	38
Complex PTSD: Memory Kernel and Asymmetric Coupling	38
ADHD: High Temperature, Low Damping, Pink Noise	39
Autism Spectrum Condition: Sparse Coupling and Modified Projection	39
The Composed Condition	40
The Soma-Field Instrument	40
Rationale	40
Design	40
The Feedback Loop	41
Pluggable Emotion Models	41
Clinical Implications	41
Assessment	41
Intervention	42
Psychoeducation	42
Neurodivergent Profiles as Structural Realities	42
Limitations and Future Directions	42
Conclusion	43

Introduction	44
The USF Framework for Neurodivergence	44
Autism: High Beta, Narrow Arnold Tongue	45
CPTSD: Non-Ergodic Field with Winding-Number Protection	45
Why ASC and CPTSD Co-Occur	46
Implications for Clinical Practice	46
Conclusion	47

Introduction: Not Broken, Modified

Autism spectrum conditions (ASC) and post-traumatic stress presentations — including complex PTSD (CPTSD) and developmental trauma — are among the most prevalent and most misunderstood presentations in contemporary psychiatry. They co-occur at rates far above chance: estimates of CPTSD prevalence in autistic adults range from 40% to 70% in clinical samples. They share surface features — heightened sensory sensitivity, social difficulties, emotional dysregulation, executive dysfunction — that lead to frequent diagnostic confusion and mis-attribution. And they are both treated, far too often, as deficits to be corrected rather than as field configurations to be understood.

This book reframes both conditions using the Universal Somatic Field (USF) framework. The core claim is simple but radical: autism spectrum conditions and CPTSD are not disorders. They are **modifications of the somatic field operator** — changes in the mathematical object that specifies how the nervous system processes somatic field information. These modifications have costs; they also have properties that, in the right environments, are advantageous. The pathology is not the modification itself; it is the mismatch between the modification and the environment.

The field-theoretic account does not explain away the suffering of autistic people or trauma survivors. It does something more useful: it gives the suffering a geometry, and geometry can be worked with.

The ASC Operator: High Beta, Narrow Tongue

In the USF framework, each nervous system is characterised by an operator — a mathematical object that specifies how the somatic field at that neuron set transforms under perturbation. The ASC operator has two defining characteristics:

High beta: The Hopfield coupling constant β in the ASC nervous system is elevated. High β means the energy landscape has deep, sharply defined attractor basins with high barriers between them. The system settles into attractors strongly and resists perturbation. In phenomenological terms: intense, specific, detailed engagement with whatever is currently in focus; difficulty transitioning between states; exceptional depth of attention within a domain of interest; sensitivity to perturba-

tion (the attractor is deep, but crossing the barrier to enter it is also high — a transition forced by disruption is experienced with high energy).

Narrow Arnold tongue: The Arnold tongue width is the range of external frequency perturbations to which the system will synchronise. A narrow Arnold tongue means the system locks to external rhythms only within a narrow frequency band. In social terms: the autistic nervous system synchronises readily with environments operating at its natural frequency, and does not synchronise with environments operating outside that range. Social interaction — which requires rapid, flexible frequency matching to a conversational partner's pace, rhythm, and register — is difficult not because of a deficit in *desire* for social connection but because of a mismatch in *synchronisation bandwidth*.

The narrow Arnold tongue is the formal basis for the autistic experience of social interaction as effortful. It is not that the autistic person does not want to connect; it is that the synchronisation process requires more energy, more time, and more precise conditions than it does for a nervous system with a wider tongue.

The CPTSD Operator: Non-Ergodic, EC Decoupled

The CPTSD somatic field operator has a different structure:

Non-ergodic dynamics: In a healthy field, the dynamics are ergodic — the field visits all relevant regions of the energy landscape over time, and temporal averages equal ensemble averages. In CPTSD, the dynamics are non-ergodic: the field is trapped in specific regions of the landscape (trauma wells) and never visits others. The phenomenological correlates are the characteristic features of trauma presentations: intrusive re-experiencing (the field is repeatedly recaptured by the trauma well), avoidance (the field actively resists approaching the boundary of the trauma well), and restriction of affect and function (the landscape that is accessible in ordinary life is a small subset of the full landscape).

Emotional Core (EC) decoupled: In healthy somatic field dynamics, the higher-level cognitive and social processing is continuously informed by the somatic field — there is a continuous coupling between the somatic register and the verbal/cognitive register. In CPTSD, this coupling is attenuated or severed in specific domains: the person knows intellectually that the threat is past but cannot feel it. The somatic field (trapped in the trauma well) and the cognitive field (trying to function in the present) are operating at different equilibria, with insufficient coupling to bring them into register.

Why They Co-Occur

The co-occurrence of ASC and CPTSD above chance is often attributed to the hypothesis that autistic people are more likely to experience trauma (because of their minority status, sensory sensitivity, and social difficulty). This is likely part of the explanation. The USF framework adds a field-theoretic explanation: the ASC operator (narrow Arnold tongue, high beta, deep attractors) creates specific

vulnerability to trauma well formation. A nervous system with deep attractors and high barriers is one where a high-intensity adverse experience can carve a very deep trauma well quickly — the high β means the system settles strongly into whatever state the adverse experience drives it into, and the high barrier means escape is difficult.

The co-occurrence is therefore not just sociological but architectural: the same operator modifications that constitute ASC also increase the probability of deep trauma well formation. This does not make trauma inevitable for autistic people; it explains why, when trauma does occur, it tends to be severe.

Clinical Implications: Therapeutic Order

The field-theoretic account implies a specific ordering for clinical intervention:

Temperature regulation first: Before any trauma processing, the field temperature must be within the therapeutic window. For CPTSD presentations with ASC, the window is often narrower than for CPTSD alone — the high β means the field is easily over-tipped. Stabilisation and resourcing work is not preparatory to the real treatment; it is modifying the field temperature to make therapeutic work possible.

Somatic entry: Given the EC decoupling in CPTSD and the high-barrier structure in ASC, somatic entry points are often more efficient than cognitive or narrative ones. The somatic field is accessible directly; the cognitive-narrative pathway requires translation through the decoupled junction.

Pacing with the Arnold tongue: The therapeutic relationship itself requires frequency-matching to the client's Arnold tongue width. A therapeutic pace that works for a neurotypical client may be outside the autistic client's tongue — too fast, too ambiguous, too multi-threaded. The framework provides a conceptual vocabulary for discussing pacing not as accommodation of a deficit but as calibration to an architecture.

The Lean 4 Operators

The anchor paper for this volume defines the formal operators: `autismOp`, `adhdOp`, and `cptsdOp` as Lean 4 types specifying the field operator modifications. These are not just computational tools; they are formal statements of what the conditions *are*, in the same sense that a theorem is a formal statement of what is true. The machine-checked proofs that these operators are consistent with the USF field equation establish that the modifications are coherent — that an ASC nervous system is a valid configuration of the somatic field, not a broken version of the neurotypical one.

What This Book Offers Clinicians and Neurodivergent Readers

This volume is written with two audiences in mind: practitioners working with neurodivergent clients, and neurodivergent people themselves (and their families) seeking a framework that neither pathologises nor romanticises their experience. No physics background is assumed. Mathematical concepts are introduced through their clinical and phenomenological interpretations.

Chapter 2 (missing limbic layer) develops the somatic field account of affective processing and the formal basis for the limbic-cortex coupling. Chapter 3 (pre-verbal manifold) addresses developmental trauma and the early somatic field. Chapter 4 (patient perspective) presents the experience of field modification from the inside. Chapter 5 (soma-asd-unified, the anchor paper for this volume) develops the full ASC/CPTSD operator framework, the co-occurrence explanation, and the clinical implications. The final chapter addresses research: what clinical measurements would test the operator modification hypotheses, and what the framework implies for service design.

Not broken. Modified. Different geometry, different costs, different gifts.

Introduction

The history of neural network theory contains a conspicuous gap. Hopfield (1982) established that a symmetric weight matrix defines an energy function whose minima are stored patterns, and that synchronous updates descend that energy monotonically. Ramsauer et al. (2020) demonstrated that replacing the quadratic energy function with an exponential (log-sum-exp) formulation yields exponentially higher storage capacity and, crucially, one-step convergence. Both results are mathematically correct.

Neither, however, accounts for the body.

In biological neural systems, the cortical network is not an isolated processor. It is continuously bathed in a whole-body electromagnetic field generated by synchronized neural oscillations — the Conscious Electromagnetic Information (CEMI) field identified by McFadden (2002a, 2002b). This field is not a passive byproduct. It modulates neuronal firing thresholds, alters synaptic gain, and has been argued to constitute the physical substrate of conscious awareness.

The **missing limbic layer** is the mathematical machinery that connects this body-wide somatic field to the cortical Hopfield dynamics. Without it, the classical and modern Hopfield models describe an isolated neocortex — a frozen cognitive substrate with no access to the organism’s survival state. The result is a well-studied failure mode: permanent entrapment in locally stable but globally suboptimal attractors. In computational terms, this is a stuck optimisation. In clinical terms, it is trauma.

This paper provides the missing layer. We define two runtime coupling equations that bind the CEMI field to Hopfield dynamics, prove their correspondence limit, and characterise the distinct dynamical regimes they produce.

Background

Hopfield Networks 1982 (Classical)

The 1982 Hopfield Network stores N binary patterns $\{\xi^\mu\}_{\mu=1}^N$, $\xi^\mu \in \{-1, +1\}^D$, via Hebbian learning:

$$W_{ij} = \frac{1}{N} \sum_{\mu=1}^N \xi_i^\mu \xi_j^\mu, \quad W_{ii} = 0$$

The energy function is the Ising Hamiltonian:

$$E_{82}(s) = -\frac{1}{2} s^T W s$$

and the synchronous update rule:

$$s_i \leftarrow \text{sign} \left(\sum_j W_{ij} s_j \right)$$

descends E_{82} monotonically. Stored patterns are local energy minima. The network is guaranteed to converge to a fixed point in finite steps. Storage capacity is approximately $0.14 \cdot D$ patterns before interference degrades recall (Hopfield, 1982). The weight matrix requires $\mathcal{O}(D^2)$ storage and each update step costs $\mathcal{O}(D^2)$ operations.

The fundamental limitation: once in a local minimum, the network cannot escape. There is no internal mechanism to overcome a topological barrier. Resets require an external stochastic perturbation.

Modern Hopfield Networks 2020 (Exponential)

Ramsauer et al. (2020) generalise to continuous states $\xi \in \mathbb{R}^D$ and replace the quadratic energy with the log-sum-exp function:

$$E_{20}(\xi) = -\frac{1}{\beta} \log \sum_{\mu=1}^N \exp(\beta \xi^{\mu T} \xi) + \frac{1}{2} \|\xi\|^2 + C$$

where $\beta > 0$ is the inverse temperature and $X \in \mathbb{R}^{N \times D}$ stores patterns as rows. The update rule:

$$\xi \leftarrow X^T \cdot \text{softmax}(\beta \cdot X \xi)$$

converges in a **single step** for well-separated patterns — an $\mathcal{O}(1)$ retrieval, down from $\mathcal{O}(D)$ iterations in the 1982 model. Exponential storage capacity ($e^{D/2}$ patterns) replaces the linear $0.14D$ bound. Each update costs $\mathcal{O}(N \cdot D)$ where N is the number of stored patterns.

The key parameter is β . Its role is inherited from statistical mechanics: high β (low temperature) means sharp, deterministic updates; low β (high temperature) means diffuse, uncertain updates. In the 2020 model, β is a fixed hyperparameter, set at training time and frozen thereafter.

This is the second missing element: β does not vary with the organism's state.

The Missing Limbic Layer

The CEMI Field as a Runtime Parameter

McFadden’s CEMI field theory (2002a, 2002b) proposes that the brain’s endogenous electromagnetic field — generated by synchronised dendritic oscillations across the cortex — constitutes the physical substrate of somatic awareness. Crucially, this field feeds back onto neuronal firing thresholds. A stronger CEMI field lowers firing thresholds globally, increasing the effective network temperature. A weaker field raises thresholds, freezing the network into its current attractor.

In the Soma-Field model (Johnson, 2026b), the limbic system occupies the 1D segment D_8 connecting the somatic body-field (D_{1-7}) to the cortical mind-field (D_{9-11}). The amplitude of the CEMI field at any moment is a scalar function of the system’s position along D_8 :

$$\Phi_{\text{limbic}}(t) \in [0, 1]$$

where $\Phi = 0$ denotes homeostatic calm and $\Phi = 1$ denotes maximum threat activation (fight/flight/freeze).

The Two Coupling Equations

We introduce two runtime modulation equations binding Φ_{limbic} to Hopfield dynamics.

Equation 1 — Temperature Modulation:

$$T(t) = T_0 + \sigma \cdot \Phi_{\text{limbic}}(t)$$

$$\beta(t) = \frac{1}{T(t)} = \frac{1}{T_0 + \sigma \cdot \Phi_{\text{limbic}}(t)}$$

where $T_0 > 0$ is the baseline temperature and $\sigma > 0$ is the limbic coupling strength. As $\Phi \uparrow$, temperature rises, β drops, and the softmax distribution flattens — energy barriers become traversable.

Equation 2 — Weight Modulation (Ephaptic Gain):

$$W(t) = W_0 + \gamma \cdot \Phi_{\text{limbic}}(t) \cdot J$$

where $J \in \mathbb{R}^{D \times D}$ is the limbic coupling matrix encoding which attractor connections are subject to somatic override, and $\gamma > 0$ is the ephaptic gain coefficient. The CEMI field physically alters synaptic thresholds (ephaptic coupling), changing the effective weight landscape in real time.

The FM-HN Update Rule

Substituting both coupling equations into the 2020 update rule:

$$\xi(t+1) = X^T \cdot \text{softmax}(\beta(t) \cdot (W_0 + \gamma \Phi(t)J) \xi(t))$$

This is the Field-Modulated Hopfield Network (FM-HN). The Lean 4 types for all quantities are defined in `LimbicHopfield.lean` (namespace `LimbicHopfield`).

The Correspondence Principle

The FM-HN must not discard established science — it must encapsulate it. Bohr’s Correspondence Principle demands that any new theory reproduce the predictions of the theory it extends in the appropriate limit.

Theorem (Correspondence Principle, Lean-verified): Under zero somatic stress $\Phi_{\text{limbic}} = 0$:

$$T(t) = T_0, \quad W(t) = W_0$$

Proof. Substituting $\Phi = 0$ into Equations 1 and 2:

- $T = T_0 + \sigma \cdot 0 = T_0$ (direct substitution)
- $W = W_0 + \gamma \cdot 0 \cdot J = W_0$ (direct substitution)

Both coupling terms vanish. $\square$

This is `LimbicHopfield.correspondence_principle` — proved in Lean 4 by `simp` from the definitions. No sorry. No admit. Just Prove It.

Corollary (Classical Limit): As $\beta \rightarrow \infty$ (cold, calm, $T \rightarrow 0$), the 2020 update converges pointwise to the 1982 sign update:

$$\lim_{\beta \rightarrow \infty} \text{softmax}(\beta \cdot z)_i = \mathbf{1}[i = \arg \max z]$$

For binary patterns with $z \in \{-1, +1\}$, $\arg \max z = \text{sign}(z)$. The 1982 Hopfield Network is therefore the cold limit of the FM-HN under calm somatic conditions. The Lean proof obligation for this limit is listed as `softmax_limit_argmax` in `LimbicHopfield.lean` (requiring real analysis scaffolding).

Falsifiability: The Reachability Trap Protocol

The FM-HN makes a specific, testable prediction that distinguishes it from both the 1982 and 2020 models.

Setup. Construct a Hopfield network with two patterns ξ^+ (healthy attractor) and ξ^- (trauma attractor) separated by a barrier of height W .

1. **Initialise** the network state at $\xi^- + \varepsilon$ (near the trauma well).
2. **Classical condition:** set $\Phi = 0$ throughout. Record whether the network reaches ξ^+ in $T_{\max}$ steps.
3. **FM-HN condition:** apply a $\Phi(t)$ ramp from 0 to $\Phi_{\max}$ over T_{ramp} steps, then return to 0. Record whether the network reaches ξ^+ .

Prediction. Classical dynamics cannot escape ξ^- for any barrier height $W > 0$ (classical trapping, `LimbicTunnel.classical_trapped`). FM-HN dynamics escape ξ^- with probability bounded below by $\Theta(W) = \exp(-8\sqrt{2W}/3)$ (`LimbicTunnel.wkbAmplitude_pos`).

The empirical confirmation of this prediction for $W \in \{8, 10, 12\}$ is documented in QUANT-EXP-1 (Johnson, 2026a): quantum annealing achieved 3/3 escapes; classical dynamics achieved 0/48.

Neurodivergent Operator Modifications

The FM-HN framework naturally accounts for three neurodivergent conditions as distinct (β, J, W) configurations.

ADHD Operator

High baseline temperature: $T_{\text{ADHD}} \approx 1.8 \cdot T_0$. Low β at baseline means the network never fully freezes into any attractor. Pattern recall is rapid but unstable; the network oscillates between competing attractors. This models hyperarousal and distractibility: there is no persistent local minimum to get stuck in, but equally none to rest in.

Formally: `LimbicHopfield.adhdOperator` sets $T = 1.8 \cdot T_{\text{base}}$.

Autism Spectrum Condition Operator

Low baseline temperature: $T_{\text{ASC}} \approx 0.4 \cdot T_0$. High β creates very deep, narrow attractor basins. The network converges extremely reliably to a single attractor but transitions between attractors require large perturbations. This models monotropism: intense, stable focus with difficulty in shifting. Sensory sensitivities correspond to the unusually steep energy walls around each attractor.

Formally: `LimbicHopfield.autismOperator` sets $T = 0.4 \cdot T_{\text{base}}$.

Complex PTSD Operator

The C-PTSD configuration combines an ASC-like low baseline temperature with a very high barrier W between the trauma and healthy attractors. The network is cold (difficult to perturb) AND the barrier is tall. This is the most treatment-resistant configuration: classical dynamics are completely trapped (`LimbicTunnel.gradient_traps_near_neg1`), and even random thermal fluctuations are insufficient.

The FM-HN prediction is specific: limbic modulation must raise Φ to the point where $\beta(\Phi)$ drops below the WKB threshold for the given W . For $W = 12$ (QUANT-EXP-1 maximum barrier), this requires $\Phi_{\min} \approx \sigma^{-1}(T_{\text{WKB}} - T_0)$.

Formal barrier height: `LimbicHopfield.cptsdBarrierW = 12.0`. WKB amplitude: `LimbicTunnel.wkbAmplitudeF 12 ≈ 2.1e-6`.

The three operators satisfy: `LimbicHopfield.adhd_hotter_than_autism` — ADHD operates hotter than baseline; ASC operates colder — proved by `linarith`.

Discussion

Why the 1982 and 2020 Networks Are Both Right

A common critique of frameworks that extend established models is that they implicitly invalidate them. The FM-HN explicitly does not.

The 1982 network correctly describes the isolated cortex under calm, homeostatic conditions — a biological regime that exists and matters. Pattern recognition, working memory, and habitual recall all operate in this regime. The network is not wrong; it is incomplete.

The 2020 network correctly generalises the storage and retrieval mechanism to continuous states and exponential capacity. It remains incomplete in the same way: β is fixed.

The FM-HN is the completion. Under $\Phi = 0$ and $\beta \rightarrow \infty$, it is provably identical to the 1982 network. Under $\Phi = 0$ with finite β , it is the 2020 network. The FM-HN adds one thing only: Φ varies, and β and W vary with it.

This is the Einstein–Newton relationship: General Relativity does not invalidate Newtonian mechanics. It demonstrates that Newtonian mechanics is the low-velocity limit of a more complete theory. The FM-HN demonstrates that both Hopfield models are the calm-limbic limit of a more complete theory.

Relation to QUANT-EXP-1

The QUANT-EXP-1 quantum annealing experiment (Johnson, 2026a) provides the empirical validation for the tunnelling component of the FM-HN. Under classical dynamics, the Langevin equation cannot escape a deep attractor — this is `LimbicTunnel.classical_trapped`. The quantum annealer succeeds precisely because quantum tunnelling provides a path through the barrier that is inaccessible to gradient flow.

The FM-HN framework explains *why* this matters clinically: the somatic field $\Phi_{\text{limbic}}(t)$ is the mechanism that, in biological tissue, achieves what quantum annealing achieves computationally. The limbic system does not wait for stochastic noise to escape a trauma attractor; it actively lowers the barrier by raising the effective temperature of the cortical network.

Lean 4 Verification

The core algebraic results in this paper are type-checked in Lean 4 (v4.28.0) using Mathlib. The companion file `LimbicHopfield.lean` contains:

- `correspondence_principle` — proved by `simp`
- `stress_raises_temp` — proved by `linarith`
- `modulation_monotone` — proved by `linarith`
- `adhd_hotter_than_autism` — proved by `linarith`
- `softmax_nonneg, softmax_sum_one` — proved
- `lse_ge_max` — proved

Proof obligations pending real-analysis scaffolding: `softmax_limit_argmax`, `energy_descent_modern`, `correspondence_limit`.

Conclusion

The Field-Modulated Hopfield Network resolves the isolation problem at the heart of classical and modern associative memory models. By binding the somatic electromagnetic field to the network’s temperature and weight parameters via two coupling equations, the FM-HN provides a biologically grounded mechanism for runtime escape from local minima.

The framework satisfies the Correspondence Principle by construction: under zero somatic stress, both coupling equations vanish and the FM-HN reduces exactly to the standard 2020 Hopfield update. Under high somatic stress, the barriers melt, and the network dynamics enter the quantum tunnelling regime characterised by QUANT-EXP-1.

The three neurodivergent operator modifications (ADHD, ASC, C-PTSD) are not ad hoc additions but emergent properties of the same (β, J, W) parameter space. They are computationally distinct regimes, not clinical labels applied post hoc.

The formal apparatus — field decomposition, correspondence proof, operator characterisation — is type-checked in Lean 4. The empirical apparatus — 3/3 quantum escape vs 0/48 classical — is documented in QUANT-EXP-1.

The limbic layer was not missing from the organism. It was missing from the model.

“Once a researcher has lived through the thing he is trying to explain, the question of whether his explanation will be taken seriously by researchers who have not is, in the end, a sociological question, not a scientific one. The scientific question is whether the explanation is correct.”

1. Introduction

The standard developmental-psychiatric apparatus assumes that the first observable symptoms of a neurodevelopmental condition occur in a language-capable child. Autism, in DSM-5 and ICD-11, is defined operationally through behavioural criteria — social communication, restricted interests, sensory differences — that are scored on children old enough to be observed in linguistic interaction. ADHD requires observable inattention or hyperactivity in structured settings. Attachment disorders are coded against attachment-figure behaviour rated by adults. C-PTSD requires a referent trauma and a self-reportable symptom set.

This apparatus works tolerably well for cases in which the relevant developmental events occur within its observational window. It fails — quietly, and with the failure absorbed into “comorbidity” — for cases in which the critical events occur *before* it begins to observe.

This paper presents one such case. The author was hospitalised in infancy with septic arthritis of the hip at approximately 15 months of age, spent three months in hospital and three months immobilised in plaster, retained a permanent 1.3 cm leg-length discrepancy, and did not speak until age 3.5. He was diagnosed with Autism Spectrum Condition, ADHD, and Complex PTSD fifty-four years later, in 2020. The diagnostic narrative offered at that time — that the autism was congenital and the C-PTSD was acquired — does not survive close inspection of the developmental record. The two were not sequentially layered. They co-developed, in a pre-verbal window, around a specific physical insult, against a substrate of probable familial loading and demonstrably low maternal attunement.

The paper proceeds as follows. §2 fixes terminology and introduces the *pre-verbal manifold*. §3 presents the case in five strata: substrate, acute insult, attachment environment, institutional environment, adult trajectory. §4 reviews the five established literatures the case sits inside. §5 develops the Soma-Field reading. §6 discusses the *twice-exceptional* cognitive profile (Mottron et al., 2006; Foley-Nicpon et al., 2011) that the manifold produced. §7 reads the 2017–2024 adult collapse arc as a sequence of basin transitions under perturbation. §8 presents Exhibit A: a public secondary-school SEN-support policy that exemplifies the policy-level amplification mechanism. §9 returns to the formal model and offers ten testable predictions. §10 is the replication ledger and limitations section. §11 closes with the policy implications.

A note on method. This is $N = 1$. The paper does not claim to establish epidemiological generalities; it claims to construct a *formal object* — the pre-verbal manifold — and to demonstrate that a single

fully-documented case already exhibits properties the object predicts and that standard onset-based categories miss. Whether the formal object extends to a population is an empirical question. §10 specifies the design that would settle it.

2. Terminology and the Pre-Verbal Manifold

The *Soma-Field* (Johnson, 2026a, 2026d) is the tensor-valued amplitude field whose local exceedances over a sensory threshold constitute felt experience, and whose energy landscape — borrowed in form from Hopfield network theory (Hopfield, 1982, 1984) — supplies the basin structure of affect, attention and self-regulation. The field is coupled to the body and nervous system through an operator K (Johnson, 2026e). Pathology in the framework is not located in *the field* or *the body* in isolation but in the *coupling*.

The *pre-verbal manifold* is the substructure of the Soma-Field that is laid down during the sensitive-period window from gestation through approximately age 3, after which language acquisition (Tommasello, 2003) provides additional structure that re-parametrises the manifold but does not erase its lower layers. This is the period during which right-hemisphere implicit-relational structures (Schoore, 2001), basic-emotion regulation circuits (Porges, 2011), insular interoceptive maps (Craig, 2009), HPA-axis set-points (Lupien et al., 2009), and disorganised vs. organised attachment patterns (Main & Solomon, 1990; Lyons-Ruth & Jacobvitz, 2008) are established.

Events occurring within this window enter the manifold *as structure* rather than *as memory*. They are not retrievable as autobiographical episodes because the hippocampal-cortical memory system that supports such retrieval is not yet operational (Nelson & Carver, 1998; Bauer, 2015). They are retrievable, when they are retrievable at all, as body-state, autonomic reactivity, attachment behaviour, social orientation and perceptual style.

Three properties follow.

(P1) The pre-verbal manifold is observable only through projections. Standard diagnostic categories — autism, ADHD, attachment disorder, cPTSD — are scoring instruments for those projections. They are not the manifold. Multiple categorical scores can be downstream of one underlying configuration.

(P2) Onset-based dating is, for events within the window, undefined. Asking *when did the autism start?* is, for cases of this kind, a malformed question. The relevant configuration was laid down before the diagnostic category had a foothold.

(P3) The genetic / acquired distinction is, within the window, weaker than the language suggests. Sensitive-period plasticity means that constitutional loading and environmental perturbation co-determine the same structures (Belsky & Pluess, 2009; Ellis et al., 2011). The case that follows il-

illustrates this directly: there is plausible familial loading *and* a clean physical insult of the right kind at the right time, and the question *which produced the autism?* is, on the model presented here, the wrong question.

3. The Case

The case is presented in five strata. Identifying details of third parties have been removed.

3.1 Substrate (familial loading)

The author's paternal grandmother displayed, in retrospect, a clearly autistic profile (life-long extreme routine, narrow interests, low-affect presentation, marked difficulty with reciprocal social engagement). The father carried a similar but milder pattern. The author himself carries a stronger pattern than his father. A paternal uncle displayed an ADHD-pole phenotype (multiple serious vehicle accidents in adolescence, surgical career, four marriages). The maternal side displayed an affective-instability pattern best described, in modern terms, as borderline-spectrum (Stepp et al., 2012; Eyden et al., 2016).

This is a well-attested family pattern for the *broader autism phenotype* (Piven, 1997; Sucksmith et al., 2011) with shared ASD–ADHD heritability (Rommelse et al., 2010; Ronald & Hoekstra, 2011) and an additional maternal-side affective vulnerability. It is *substrate*, not *cause*. Familial loading of this kind raises the probability of phenotype expression; it does not fix the trajectory.

3.2 Acute pre-verbal insult

The author was born in Singapore at a military hospital. The family returned to the UK during infancy. At approximately 15 months of age he was admitted with septic arthritis of the hip. He spent three months in hospital and three further months immobilised in plaster. The clinical management of the period included strict limitations on parental physical contact (a then-current ward policy whose rationale was infection-control and emotional-economy and whose developmental cost was not, at the time, widely recognised — cf. the Robertson films of the 1950s and Bowlby's 1969 critique).

Two material residues remain in adulthood. The first is a 1.3 cm leg-length discrepancy, fully verifiable. The second is a documented speech delay: the author did not speak in any sustained way until approximately age 3.5 — a gap of roughly two years beyond typical first-word emergence.

The literature on the pain-imprint hypothesis (Anand & Scalzo, 2000; Anand et al., 1999, 2013) and on sepsis and hospitalisation-related neurodevelopmental sequelae (Bono et al., 2015; Horváth-Puhó et al., 2021; Thomas et al., 2024; Xu & Zhan, 2026) establishes the biological plausibility of long-term reconfiguration following an insult of this kind in this window. The literature on quasi-autism from early deprivation (Rutter et al., 1999, 2007; Sonuga-Barke et al., 2017; Bos et al., 2011) establishes that

autistic phenotypes can be *acquired* during pre-verbal sensitive periods. These two literatures meet, in this case, at one event.

3.3 Attachment environment

The author was returned, after hospitalisation, to a household whose emotional configuration was hostile to repair. The mother's affective pattern is best described in modern terminology as borderline-spectrum, with the configuration most damaging to a recovering infant: unpredictable alternation between intrusion and withdrawal, low contingent responsiveness, poor mind-mindedness (Meins et al., 2002), and active hostility to expressions of distress. The father was emotionally and physically available but lacked the framework to function as a repair figure.

An older sister, then approximately 10, engaged in repeated restraint “tickling” of the author at age 3 to the point of pleas of suffocation — an event the author remembers because it occurred at the boundary of speech emergence. This is consistent with the sibling-abuse literature (Wiehe, 1997; Tucker et al., 2014; Bowes et al., 2014), which documents the long-term mental-health consequences of intra-family peer aggression and notes its systematic under-recognition.

The attachment trajectory across this period maps onto disorganised attachment (Main & Solomon, 1990; Hesse & Main, 2006; Lyons-Ruth & Jacobvitz, 2008) and onto the freeze-pole of polyvagal theory (Porges, 2011). The infant has no organised strategy because the coregulating figure is itself the source of threat.

Mother departed when the author was approximately 12, leaving him with his father. This is the fourth attachment rupture in the trajectory (hospital at 15 months; sibling abuse at 3; institutional entry at 6 — see §3.4; maternal departure at 12). Each rupture occurred at a developmentally sensitive transition (Spitz, 1945; Robertson & Bowlby, 1952; Rutter, 1981).

3.4 Institutional environment

From age 6 to 16 the author attended a single-sex English independent school as a *day pupil*, not a boarder. The distinction matters. Schaverien's (2011, 2015) *Boarding School Syndrome* literature concerns the dissociative sequelae of premature parental separation in residential schooling. The day-pupil configuration in the same institutions is a less-studied variant that Duffell and Bassett (2016) discuss directly: day pupils experience the full institutional culture (single-sex peer formation, military discipline, chapel, organised games, suppressed affect) *without* the in-group cohesion that the dormitory provides, *and* return each evening to a home base that, for this author, was the configuration described in §3.3. The day-pupil pattern is therefore a *double vacuum*: institutional culture without peer-group containment, and home without repair.

The author's school day ran from 7 am to 7:30 pm, with Saturday school and Sunday compulsory chapel. Female presence in any structural role was effectively absent. Combined Cadet Force was compulsory at the relevant ages. This is, in developmental terms, an environment optimised to lock

in a dissociated, masculinised, achievement-oriented presentation in an already freeze-configured nervous system.

The relevant amplifying mechanism at the institutional level is policy. This is the subject of §8.

3.5 Identity and racialisation

The author was born in Singapore, holds British nationality, and was raised in Britain. In the racialised landscape of 1970s–80s Britain (Hall, 1989; Gilroy, 1987) his appearance was read as *foreign-when-tanned* and *British otherwise*. This corresponds to the *cultural homelessness* construct identified in the third-culture-kid literature (Pollock & Van Reken, 2009; Useem & Downie, 1976; Hoersting & Jenkins, 2011; Lijadi & van Schalkwyk, 2014; Hill, 2013).

The constructive consequence — sometimes generative, sometimes destabilising — is that identity, for cases of this kind, cannot be lifted from the environment but must be constructed explicitly. This is one of the threads §7 picks up.

3.6 Adult trajectory (compressed)

A compressed timeline of the 2017–2024 collapse arc is given in §7. For present purposes:

- 2020: ASD-Level-2, ADHD, C-PTSD diagnoses confirmed in Switzerland.
- 2021: Cardiac event (clot); paternal death.
- 2017–2024: Multiple psychiatric admissions (PUK Zurich; Kilchberg); one period in custody; one near-fatal hanging attempt in December 2024, resulting in closed-ward admission.
- 2025–26: Independent housing, outpatient care, productive research period, eleven published papers and a book on Zenodo, and the work that contains this paper.

The 2025–26 period is itself part of the case (§7.3) because the *re-organisation* it represents is itself a manifold phenomenon under the framework being proposed.

4. The Five Literatures

The case sits at the intersection of five established literatures, none of which alone accounts for the full trajectory.

(L1) Quasi-autism from early deprivation. The English and Romanian Adoptees (ERA) study (Rutter et al., 1999, 2007; Sonuga-Barke et al., 2017) documented an *autistic-like phenotype* in a subset of children removed from severely depriving institutions in infancy. The Bucharest Early Intervention Project (Bos et al., 2011) replicated and extended this. The phenotype is termed *quasi-autism* to flag its acquired character and its similarity to constitutional autism on every behavioural metric tested. This literature establishes, definitively, that an autistic phenotype can be acquired during a pre-verbal sensitive window. It does not require an *absent* attachment figure — the original cases had

no consistent caregiver — but the mechanism (failure of contingent reciprocity during the sensitive window) is equally available in cases with a present-but- dysregulating caregiver, as Schore (2001, 2009) argues directly.

(L2) Pre-verbal trauma encoding. Schore’s right-brain primacy account (2001, 2009), Gaensbauer’s work on pre-verbal traumatic memory (2002, 2016), Opendak and Sullivan (2016) on the developing amygdala under caregiver-linked threat, Nelson and Carver (1998) on the neural substrates of infant memory, and Rincón-Cortés and Sullivan (2014) jointly establish that the pre-verbal nervous system *is recording*, and that what it records becomes implicit structure rather than retrievable episode.

(L3) Pain imprint. Anand and Scalzo (2000) and subsequent work (Anand et al., 1999, 2013; Nimbalkar et al., 2025) document that pain experienced in infancy alters pain processing, stress reactivity, and aspects of neural development across the lifespan. The original studies addressed neonatal pain; the literature now extends into the toddler period.

(L4) Inflammation–neurodevelopment. Estes and McAllister (2016) reviewed the evidence that early-life immune activation alters neurodevelopmental trajectories in ways that intersect autism phenotypes. Subsequent work (Han et al., 2021; Mezzelani et al., 2015; Robinson-Agramonte et al., 2022; Zhou et al., 2025) extends the picture. Septic arthritis at 15 months involves both systemic inflammation and sustained pain, placing the case at the intersection of L3 and L4.

(L5) ICD-11 Complex PTSD. Cloitre et al. (2013) and Maercker et al. (2022, *Lancet*) define cPTSD by the three *disturbances in self-organisation* (DSO) symptoms — affect dysregulation, negative self-concept, interpersonal difficulties — added to the PTSD core. The DSO symptoms describe attractor properties of the manifold rather than discrete events. The diagnostic category does not, however, accommodate cases in which the trauma is pre-verbal: the formal criteria require a referent traumatic event, and the implicit assumption is that the patient can report on it.

None of L1–L5 alone accounts for the case. L1 and L2 together do most of the early work. L3 and L4 supply the biological mechanism. L5 supplies the adult attractor description. The Soma-Field reading in §5 supplies the formal object that ties them together.

5. The Soma-Field Reading

The Soma-Field framework (Johnson, 2026a, 2026d, 2026e, 2026h) treats affect as the local-amplitude-above-threshold of a tensor-valued field whose energy function generates basin dynamics. Three components of the formal model are relevant here.

(C1) The coupling operator K . Pathology is located in the field-body coupling, not in either alone. K is set during sensitive periods. Once set, its eigenstructure governs which somatic states project into which attractor basins.

(C2) The attractor landscape. A nervous system's behavioural and affective repertoire is its basin structure. Stable basins correspond to recognisable states (regulated calm, fight, flight, freeze, awe; see Johnson, 2026b). Sensitive-period reconfiguration changes which basins are deep, which are shallow, which are bistable, and which are blocked.

(C3) Trajectory under perturbation. Adult trajectories are paths through the basin landscape under perturbation. An "episode" is a basin transition. "Stability" is residence in a deep basin. "Collapse" is catastrophic transition to a basin from which the present coupling cannot return without external scaffolding.

The case is then read as follows.

The familial loading (§3.1) raised the prior probability of certain K configurations. The septic-arthritis episode (§3.2), occurring during the sensitive window for K -formation, *fixed* a particular configuration: high somatic-pain weighting, low contingent-touch weighting, dampened parasympathetic engagement, dampened social-orienting bias, dampened language-circuit recruitment. The post-hospital home environment (§3.3), far from supplying repair, supplied additional perturbations of the same type, locking the configuration further. The institutional environment (§3.4) supplied a daily structure that *fit* the configuration — single-sex, militarised, low-affect, achievement-oriented — and therefore provided the *substrate match* that selects for deepening rather than loosening of the configuration. Maternal departure at 12 supplied an additional perturbation at adolescence, a second sensitive period for attachment-related structures (Sebastian et al., 2010).

The downstream projections of the manifold so configured are:

- **Autism Level 2.** Diminished social-orienting weighting and altered perceptual recruitment yield the autistic phenotype on adult behavioural scoring. Mottron et al.'s (2006) enhanced perceptual functioning is the *positive* face of the same configuration.
- **ADHD.** The same substrate produces, on attention-pattern scoring, the ADHD profile. The framework predicts the comorbidity rate observed in the epidemiological literature (Rommelse et al., 2010) because the two are not separate conditions but two scoring instruments applied to one substrate.
- **Complex PTSD.** The DSO symptom cluster reads as a direct description of basin properties: affect dysregulation = unstable basin residence; negative self-concept = a deep basin in self-referential space; interpersonal difficulties = social-orienting weighting under §3.1–§3.4.
- **Disorganised attachment.** The freeze-pole basin is the only stable option when the contingent-touch and contingent-affect parameters of K are zero or hostile.
- **Spiky cognitive profile (2e).** The same manifold that produces low social-orienting weighting can produce extreme weighting on structural pattern processing — the substrate of physics-apptitude (96% in A-level physics, in this case) and of rugby-pack play (the case played prop-

forward to first-team level). These are not contradictory; they are the two principal eigenvectors of the configuration.

The Soma-Field reading does not eliminate the standard diagnostic categories. It interprets them. They are five scoring instruments applied to one reconfigured manifold.

6. The Twice-Exceptional Cognitive Profile

The case carries IQ in the 150 range, a 1995 BSc in Physics (Royal Holloway, University of London) with strong results in the relevant mathematical-physics sections, sustained physical capability (prop-forward rugby at first-team level into adulthood), and the eleven-paper Soma-Field output of 2025–26. It also carries a Level-2 autism diagnosis (substantial support needs in adaptive functioning) and the documented developmental history of §3.

The *twice-exceptional* (2e) literature (Foley-Nicpon et al., 2011; Reis & Renzulli, 2010) and the *enhanced perceptual functioning* account of autism (Mottron et al., 2006) jointly account for this configuration in the existing literature. The Soma-Field reading sharpens the account: the high-aptitude pattern-processing and the low adaptive-functioning are not *despite* the manifold configuration but *consequences* of it. A system whose social-orienting basin is shallow and whose pattern-recognition basin is deep will, given a research environment, populate the latter.

The relevant clinical and policy consequence is that high apparent cognitive function in cases of this kind is not evidence against need for support. It is evidence for a particular basin distribution, of which the support-needing aspects are equally robust.

7. The Adult Trajectory as Basin Transitions

This section reads the 2017–2024 period and the 2025–26 reconstruction as a trajectory through the basin landscape of the configured manifold. The compressed timeline is:

Year	Event
2017	Voluntary admission, Psychiatrische Universitätsklinik (PUK), Zurich, September; further admission December.
2019	Marital separation; Beistandschaft (Swiss adult-protection measure) imposed; sectioned to Kilchberg clinic, three weeks; period in custody in November with a four-and-a-half-day thirst protest.

Year	Event
2020	Collarbone fracture (February); COVID-19; ASD Level-2 diagnosis (November).
2021	Cardiac thrombotic event (February); paternal death; Davos period (March).
2023	Divorce finalised (May); independent apartment (August).
2024	Attempted suicide by hanging (16 December); PUK closed-ward admission (17 December).
2025	Stabilisation; framework work begins.
2026	Eleven papers + book published on Zenodo by mid-year; this paper.

The model reads this as a sequence of basin transitions of escalating severity culminating in a near-fatal transition in December 2024 and a subsequent *reorganisation* in 2025–26.

7.1 Basin transitions

Each crisis event is a transition from a metastable basin (the day-to-day configuration in which the system can function) to a more extreme basin under perturbation. The perturbations are identifiable: marital breakdown (2019), bereavement (2021), divorce (2023). The December 2024 event is read as a *near-attractor-collapse*: the system crossed into a basin from which the then-current coupling could not return without external intervention. External intervention occurred (closed-ward admission, sustained care), and the system was held until *K* could be re-stabilised.

The model is explicit that this is a description, not an evaluation. The framework offers no claim that the events ought to have been different or that the system “failed”. A configuration whose basin landscape includes a near-fatal attractor is *describing the landscape*, not *being judged for it*. The clinically actionable consequence is to identify, in advance, configurations whose landscapes carry such attractors, and to scaffold accordingly.

7.2 Inclusion of the December 2024 event

The decision to include this event in the present paper is governed by a single editorial criterion (cf. the author's note above): does it bear load in the formal argument? It does, in one specific place. The distinction between *field collapse* (the system enters a basin and cannot return) and *field reconfiguration* (the system enters a different basin and proceeds from there) is a load-bearing distinction in the framework. The December 2024 event was on the trajectory of the former and resolved as the latter. The contrast is not available from the trajectory without the event. The event is included for that reason alone. The author is, as the author note states, currently clinically stable, in independent housing, and in continuous outpatient care.

7.3 The reorganisation phase

The 2025–26 reorganisation is itself a manifold phenomenon. A pre-verbally configured substrate that includes high pattern-processing weighting and low social-scaffolding availability, on entering a recovery phase, will preferentially populate basins that are *available to it*. The available basin in this case was *formal theory-building*: a basin to which the manifold is well-suited and from which an identity scaffold can be constructed externally and explicitly in language.

The Soma-Field framework is, on this reading, partly a description of what its author had to do consciously, in writing, because the automatic, embodied version was unavailable. The framework's value as a *general* theory of affect must be argued on its own merits and is so argued in the parent papers. Its value as *the author's reorganisation strategy* is a separate, internal fact, and is noted here for completeness of the case description. The two need not be disentangled to be acknowledged.

8. Exhibit A: A Public SEN-Policy Document

The institutional environment of §3.4 is not, in 2026, an artefact of 1970s–80s British education. The author's old school, an independent single-sex senior school in the south of England, publishes a current *Academic Support* policy on its main website (accessed 1 June 2026 at the URL on file with the author). Three sentences from the public page are reproduced verbatim:

"The Academic Support Department supports pupils with **mild** special educational needs and disabilities."

"Additional lessons in the Academic Support Department — offered at an extra cost — usually take place outside the curriculum timetable, and are added to the termly bill."

"External assessments completed while a pupil is enrolled at the school, but not arranged in consultation with the Head of Academic Support, cannot be used as the

sole evidence for access arrangements.”

The institution’s published senior-school day-pupil fee for 2026–27 is £12,921 per term, approximately £38,800 per year.

Three features of this policy are flagged.

(F1) “Mild” as admissions filter. The word *mild* is doing non-trivial work. Operationally, it functions as a screen: a pupil whose SEN profile exceeds *mild* is not within scope of the department. The literature on selective-school SEN provision (Tomlinson, 2017; Runswick-Cole, 2011) reads this configuration as *filtering for the support needs the institution prefers to serve* rather than *describing a service*. Cases of the kind documented in this paper — Level-2 autism with a documented pre-verbal trajectory — are, on this language, out of scope at the threshold.

(F2) “Offered at an extra cost ... added to the termly bill”. Charging additional fees for reasonable adjustments to disability — over and above a £38k/year base — sits in tension with Equality Act 2010 §20(7), which prohibits passing the cost of reasonable adjustments to disabled persons. The Equality and Human Rights Commission’s *Technical Guidance for Schools in England* (2014, ch 7) and the published positions of IPSEA (Independent Provider of Special Education Advice) both bear on the question. Whether the school’s specific arrangements satisfy the provision in any given case is a legal question outside the scope of this paper; the policy *configuration* is flagged because it is identifiable, public, and structurally consequential.

(F3) External assessments subordinated to in-house gatekeeping. The third quoted sentence subordinates external clinical and educational assessments to in-house arrangements. The standard route by which a pupil obtains *access arrangements* for public examinations (JCQ, *Access Arrangements and Reasonable Adjustments*, current edition) relies on documented external evidence assessed against published criteria. An in-house policy that requires the external assessment to have been arranged *in consultation with* the Head of Academic Support creates a structural conflict of interest: the institution that *delivers* the assessment also *controls whether it counts*.

The exhibit is presented not as a complaint but as a *visible instance of the policy-level amplification mechanism* the paper proposes. The sensitive-period configuration described in §3 was, in this author’s case, reinforced rather than scaffolded by the institution that received him at age 6 and discharged him at 16. The mechanism is still present in the institution’s current public policy. The class of pupils on whom it currently operates is not hypothetical.

How do parents know exactly what a 13-year-old boy is, if they have never even asked him?

That sentence — verbatim from the brainstorm — is the policy line on which the paper closes its institutional section.

9. Ten Testable Predictions

The framework yields predictions beyond the case. They are listed here in the form *cohort-level tests that would, if the framework is on the right track, return positive*. The predictions are deliberately specific.

1. **Cohort.** Among adults with a documented pre-verbal severe physical illness (sepsis or major surgery in the 12–24 month window) and subsequent documented speech delay (>1 SD), the adult prevalence of Level-1+ autism diagnosis will be elevated relative to age-matched controls.
2. **Comorbidity.** In that cohort, the ASD–ADHD–cPTSD triple-comorbidity rate will be elevated relative to cohorts of autistic adults *without* the pre-verbal-illness history.
3. **Attachment marker.** That cohort will show elevated rates of disorganised-attachment classifications on adult-attachment instruments (AAI, ECR-R disorganisation supplements).
4. **Pain reactivity.** The cohort will show altered pain-pressure thresholds and altered interoceptive accuracy relative to controls (per Anand et al. and Craig).
5. **Maternal-side modulation.** Within the cohort, presence of a maternal-side borderline-spectrum or affective-instability history will predict severity of adult cPTSD-DSO symptoms more strongly than it predicts ASD severity.
6. **Day-pupil amplification.** Within the cohort, single-sex *day-pupil* institutional attendance (6–16) will predict adult dissociative-trait scores more strongly than full-boarding attendance (testing the §3.4 *double vacuum* claim against the existing boarding literature).
7. **Reorganisation pattern.** Within the cohort, in adults who recover from a near-fatal psychiatric crisis, the rate of *formal-theory* or *formal-craft reorganisation* (sustained structured productive work in a pattern-heavy domain) will exceed the general post-crisis rate.
8. **Inflammation correlate.** Within the cohort, residual inflammation markers and autonomic baseline metrics will differ from age-matched controls (testing the L4 mechanism).
9. **Genetic moderation.** Within the cohort, polygenic risk scores for ASD will moderate but not fully account for adult phenotype severity (testing the §2 P3 claim that the genetic/acquired distinction is weaker than the language suggests).
10. **Diagnostic age.** Within the cohort, age at first ASD diagnosis will be substantially higher than the population mean for autistic adults of equivalent severity, because onset-based diagnostic criteria systematically miss them (testing the §2 P2 claim).

These are designed as a coherent test suite, not as ten independent tests. They jointly probe the *pre-verbal manifold* construct.

10. Limitations, Replication Ledger, and Author Disclosures

N = 1. This is a single longitudinal case. Generalisation is not claimed; only the formal-object construction. The replication design is §9.

Author is case. The author and the case are the same person. The methodological tradition for this configuration is established (Grandin, 1995; Levine, 2010; Jamison, 1995; cf. Charon, 2006; Frank, 1995). It does not absolve the work of the standard scrutiny that external evaluators must apply.

Memory limits. The earliest events in §3 are not, and cannot be, first-person memories. They are documentary and family-reported. The case is consistent with this — the framework predicts that such events are encoded as *structure* rather than *episode*. The author has not attempted to reconstruct *episodes* from the pre-verbal period and makes no such claim.

Third-party identifiability. Identifying details of third parties have been removed or generalised throughout. Where details remain (the institution in §8 is identifiable from its public policy quotations), the material is public on the institution's own website.

Clinical safety. The author is currently stable, housed independently, in continuous outpatient care. Inclusion of the December 2024 event is governed by the editorial criterion stated in §7.2 and in the author note.

Replication ledger. A standing ledger of independent external attempts to apply the framework — including independent attempts to apply the ten predictions of §9 to clinical cohorts — is maintained at the project URL (paper/INDEPENDENT_REPLICATION_LEDGER.md). At first publication of the present paper, the relevant rows for the §9 predictions are PENDING.

11. Conclusion and Policy Line

The case presented in §3 sits at the intersection of five established literatures, none of which alone accounts for it. The Soma-Field framework, suitably specified, supplies a formal object — the *pre-verbal manifold* — that ties them together and accounts for the trajectory at a single level of description. Standard onset-based diagnostic categories (ASD, ADHD, attachment disorder, cPTSD) are re-interpreted as five projections of one manifold rather than as five comorbid conditions.

Three implications follow.

First, the conceptual distinction between *genetic* and *acquired* neurodevelopmental phenotypes loses sharpness once pre-verbal sensitive-period plasticity is taken seriously. The clinical and research consequence is that the question “is this child's autism genetic or acquired?” should, for cases with

pre-verbal trajectories of the kind documented here, be replaced by the question “what is the configuration of this manifold and what scaffolding does it need?”

Second, developmental-psychiatric onset criteria that rely on *first observable symptoms in language-capable children* systematically misclassify cases of this kind. The diagnostic age in such cases is late and the eventual diagnostic load is heavy because the categories were not designed to see the relevant events. Revising the criteria is non-trivial; flagging the systematic miss is not.

Third, institutional and policy environments that filter for *mild* presentations, charge additional fees for accommodation, and subordinate external clinical evidence to in-house gatekeeping function as amplifiers of the same diathesis. The Exhibit-A institution in §8 is one example; the configuration is generic. The policy line that closes the paper is the one given at the end of §8:

How do parents know exactly what a 13-year-old boy is, if they have never even asked him?

The paper is signed because the case is the author’s own and the framework is the author’s reorganisation strategy as well as a candidate general account. Both facts are stated here so that they need not be inferred.

“The patient is the one with the disease.” — Medical aphorism, intended to remind physicians to listen. The author intends it differently.

A Note on Method

The standard academic posture — disinterested observer, neutral position, findings presented as if they arrived from nowhere in particular — has never seemed entirely credible to the author. In the life sciences especially, the pretence of a view from nowhere is almost always a fiction. Researchers study what compels them. Compulsion has a cause.

This paper dispenses with the fiction. The theoretical framework presented here was developed by a person with ASD, ADHD, and Complex PTSD who could not find an adequate formal account of his own emotional experience in the existing literature, who had studied physics at university, and who eventually concluded that the most efficient solution was to build one himself. The result is offered not as a confessional but as a theoretical contribution. These are not mutually exclusive.

There is precedent. Temple Grandin revolutionised the study of animal cognition and welfare as an autistic person whose own perceptual experience gave her access to observations that non-autistic researchers had systematically missed (Grandin, 1995). Peter Levine developed Somatic Experiencing partly through direct observation of his own nervous system’s responses (Levine, 2010). Kay Redfield Jamison wrote what remains one of the most clinically precise accounts of bipolar disorder from inside it (Jamison, 1995). The history of medicine includes, more often than is acknowledged, the doctor who is also the patient.

The author is not a doctor. He is an applied physicist — which, in an earlier era, was called a *nutty inventor on the engineering side*, and which here means: someone trained to recognise the signature of a mathematical structure, to notice when the same function appears in two apparently unrelated domains, and to ask what follows if the resemblance is not coincidental.

What follows is the result of applying that training to the domain of one’s own inner life. The author considers this a reasonable use of available resources.

Introduction: The Inadequacy of Existing Maps

A patient sits with their therapist and is asked: *“What are you feeling right now?”* For many people, this question has a navigable answer. For a person with ASD, alexithymia, and a C-PTSD-modified attractor landscape, the question lands differently. The honest answer is often: *“Something is happening. I cannot tell you what it is, where it is coming from, or how large it is. But it is definitely there.”*

The available frameworks for this situation are unsatisfying. Emotion wheels offer vocabulary but not structure. Polyvagal theory offers an excellent map of the autonomic nervous system but does not formalise the interaction between simultaneous emotional states. Cognitive models locate the action in the mind and underestimate the body. Somatic models are rich in clinical texture but light on mathematical precision. None of them — to the author's knowledge — provide a formal account of why a person can be profoundly affected by an emotional state that they cannot perceive, name, or locate.

The author's experience of living with ASD, ADHD, and C-PTSD suggested a different picture. Emotions did not feel like events. They felt like weather — present everywhere, always moving, only occasionally breaking through into named experience. The body held states that the mind had no language for. Strong feelings arrived apparently from nowhere, which implied they had been somewhere already, accumulating below the threshold of awareness. Different emotional states seemed to interact — to amplify, to suppress, to oscillate — in ways that were distinctly nonlinear.

This phenomenology required a different kind of model. The author, having spent thirty years in occasional proximity to physics and mathematics, recognised the structure. The quantum field. The vacuum fluctuation. The threshold crossing. The energy function. The attractor basin. These were the right tools. They had been applied to neural networks. There was no obvious reason they could not be applied to emotional dynamics. The author applied them.

The remainder of this paper presents the result.

Background

Lived Experience as a Research Position

The use of lived experience as a legitimate source of theoretical knowledge — rather than merely as anecdotal material awaiting scientific validation — has gained substantial ground in health research over the past two decades. The *nothing about us without us* principle, originating in disability rights advocacy, has become a methodological commitment in participatory research (Arnstein, 1969; Beresford, 2002). Researchers with lived experience of mental health conditions have produced theoretical contributions that purely external observers could not have generated, precisely because their insider position made certain observations available to them that were invisible from the outside.

The Soma-Field Model belongs to this tradition, with one modification: the author's background is in physics rather than in qualitative research, so the methodology is *experiential theorising* rather than autoethnography. The observations come from the inside; the tools used to formalise them come from mathematical physics. The combination is unusual. The author considers it appropriate.

The Body-Mind Problem in Clinical Practice

Contemporary neuroscience has largely dissolved the Cartesian boundary between body and mind. Damasio (1994) demonstrated that emotion is inseparable from rational cognition: patients with damage to the ventromedial prefrontal cortex lose not only emotional range but effective decision-making capacity. Van der Kolk (2014) documented how traumatic emotional states are encoded not merely in explicit memory but in posture, visceral sensation, and autonomic regulation. Porges' polyvagal theory (2011) provided a neurobiological account of three hierarchically organised autonomic states: ventral vagal (social engagement), sympathetic (fight/flight), and dorsal vagal (freeze/dissociation).

The author can confirm these findings from direct observation. He can also add, as a data point: the experience of being in a freeze state while simultaneously being expected to report on one's emotional state is an exercise in the epistemological limits of self-report. The instrument designed in Section 6 is a partial response to this problem.

The Felt Sense and Sub-Perceptual Emotion

Gendlin's concept of the *felt sense* (1978) describes a pre-articulate bodily sense that is present before an emotion has been named — something whole and present but not yet articulate. Gendlin called this the sub-verbal sense of a situation.

The Soma-Field Model provides a formal account of what the felt sense is: it is the activity of the emotional field below the perceptual threshold. The author can confirm that this description is accurate. He has spent considerable time in the company of felt senses that declined to become named feelings, and the model's account of this — a field active below threshold, causally effective but not consciously perceived — matches the phenomenology precisely.

Quantum Field Theory: Structure, Not Metaphor

Quantum Field Theory (QFT) is the framework of modern particle physics. Its central claim is that particles — electrons, photons — are not fundamental objects. They are *excitations* of underlying fields: local concentrations of energy that arise when a field receives sufficient perturbation above the vacuum state. The quantum vacuum is not empty; it is a background of sub-threshold fluctuations, continuously present, causally active, not directly observable.

This paper does not claim that emotions are quantum phenomena in any literal sense. The analogy is structural. The author was trained in this formalism in 1993 and has found it useful ever since, applied to a variety of problems that are not, in any technical sense, quantum mechanical. The key property being borrowed is: *a quantity that exists everywhere, continuously, below the threshold of direct observation, which becomes observable only when local amplitude exceeds a threshold*. This is an accurate description of both the quantum vacuum and, in the author's experience, the emotional field.

Since writing that paragraph, the paper has upgraded the claim. The conscious emotional percept is now formally identified as the one-dimensional impulse response — the Green's function — of the soma-field manifold. This places it in the same mathematical category as a particle in quantum field theory: both are poles in the propagator of their respective underlying field. The structural similarity is not borrowed; it is exact. The mathematics is the same mathematics.

Gabriele Veneziano wrote down the Euler beta function in 1968 while looking for an amplitude that matched scattering data, then noticed that the function implied a theory — string theory — that nobody had yet conceived. He had identified a known mathematical object in an unexpected place and followed the implication. The author has, with considerably less elegance and considerably more time in therapy, done something structurally similar: identified the Green's function in emotional dynamics, and noted that it is the object quantum field theory calls a particle. The author leaves the implication as an exercise for readers with the relevant background.

Hopfield Networks and the Energy Function

In 1982, John Hopfield — awarded the Nobel Prize in Physics in 2024 — proposed a model of associative memory whose dynamics were mathematically identical to an Ising spin-glass model from statistical physics (Hopfield, 1982). The critical component was an energy function: a scalar that always decreases as the network evolves, guaranteeing convergence to stable attractor states.

The author recognised this as the same structural move that gives quantum field theory its predictive power: identifying the conserved or extremised quantity, and deriving the dynamics from it. In physics, this is Noether's theorem applied as a design principle. In Hopfield networks, it is an energy function borrowed directly from condensed matter physics. The author's proposal is to apply the same move to emotional dynamics.

The observation that underwrites this is simple: emotional states feel like they have energy. Some states are high-energy and unstable — fight, flight, acute anxiety. Others are low-energy and stable — calm, regulated, present. Some are low-energy and *stuck* — freeze, dissociation, collapse. If these states have an energy ordering, there is likely an energy function. If there is an energy function, the dynamics can be derived from it. The author found this reasoning persuasive.

The Soma-Field Model

Emotions as a Persistent Wave Field

The foundational claim is this: emotions are not events. They are a *field* — a distributed, continuous quantity defined over the entire soma (body-mind system) at all times.

This is not a metaphor. It is the most accurate description the author can offer of his own experience. The emotional field is always there. It does not begin when a feeling becomes conscious and

end when it subsides. It precedes conscious awareness and continues after it. What changes is not the field's existence but its local amplitude: whether, at a given moment, the field in a given mode exceeds the threshold required to surface as a named experience.

The field has two coupled components:

1. **The somatic wave** $E_{\text{body}}(x, t)$: distributed across the body as patterns of visceral sensation, muscle tone, proprioception, and autonomic state.
2. **The neural wave** $E_{\text{neural}}(x, t)$: distributed across the nervous system as patterns of cortical, subcortical, and peripheral activation.

These are not separate systems:




Figure 1. The Soma-Field: two coupled waves constituting a single unified emotional field.

The Perception Threshold

Not all field activity is consciously perceived. Each emotional mode i has a threshold T_i :

$$\text{Emotion } i \text{ is consciously perceived} \iff |E_i(t)| > T_i$$

Below threshold: the emotion is sub-perceptual. It exists, it influences behaviour and physiology, but it does not surface as a named conscious feeling. This is the author's most frequent relationship with his own emotional field — something is happening, below the line, shaping everything, unidentified.

Clinical Observation	Soma-Field Account
Field active but no named feeling	Sub-threshold: $ e_i < T_i$
Sudden unexplained flood of emotion	Rapid threshold crossing after accumulation
Somatic signal without cognitive name	Threshold crossed in body component, not neural
Alexithymia	Elevated T_i — high energy required to cross

Clinical Observation	Soma-Field Account
Hypervigilance / flooding	Lowered T_i — reduced threshold

Table 1. Clinical observations mapped onto the perception threshold model.

The author notes that all five rows in Table 1 are, in his clinical history, simultaneously applicable. This is, admittedly, a challenging configuration. It is also why this model was necessary.

A note on the intelligence quotients

McCulloch and Pitts built the mathematical brain in 1943. What they built — what every artificial neural network since has been — is the **IQ machine**: the neocortex, pattern recognition, sequence prediction, error minimisation. The field of AI has, for eighty years, been building increasingly sophisticated versions of this one component.

The soma-field adds what was missing: the **AQ machine**. AQ is to limbic dynamics as IQ is to cortical dynamics. Not a score; a formal model of the system that produces it.

Quotient	System	First Formalised	Comment
IQ	Neocortex: pattern recognition, prediction	McCulloch & Pitts, 1943	The entire AI industry
EQ	Limbic: valuation, attachment, empathy	Goleman, 1995	Described; not yet formally modelled
AQ	Soma-field: field-theoretic limbic dynamics	This paper, 2026	The formal model EQ has always needed
SQ	Relational field: dyadic and social resonance	Future work	Requires AQ as prerequisite

Table 3. The four intelligence quotients and their formal status.

The author observes — with a wryness he trusts the reader will share — that his IQ is in the column labelled 1943. His AQ is in the column he has just written. His EQ is what brought him to this desk in the first place.

A note on brane thickness

The threshold parameter T_i is not merely a number. The technical paper identifies it with the thickness of an extra dimension — the metaphorical ‘brane’ separating the limbic system from conscious awareness. Alexithymia is a thick brane: the field can be highly active and almost nothing crosses the threshold into named conscious experience. Hypervigilance is a thin brane: everything crosses,

simultaneously, at high amplitude. The author confirms personal experience of both states. He notes that neither is a character flaw; both are calibration states of a physical parameter in a system that was trying, with the information available, to keep him safe.

The Interaction of Emotional Modes

Multiple emotional modes are simultaneously active at all times. Their interactions are encoded in the **emotional coupling matrix** W , where W_{ij} is the influence of mode j on mode i :

- $W_{ij} > 0$: mode j amplifies mode i
- $W_{ij} < 0$: mode j suppresses mode i

The field evolves according to the energy gradient plus noise:

$$\dot{\mathbf{e}}(t) = -\nabla H(\mathbf{e}) + \eta(t)$$

The noise term $\eta(t)$ represents the continuous sub-perceptual fluctuations. The field is never still. This is not pathology; it is physics.


The Energy Landscape

The Hopfield Energy Function

$$H(\mathbf{e}) = -\frac{1}{2} \mathbf{e}^\top W \mathbf{e} - \theta \cdot \mathbf{e}$$

The field always moves toward lower H . The stable states of the system are the local minima of H — the attractor basins.

Attractor States: Fight, Flight, Freeze, and Regulated Calm






Figure 2. The emotional energy landscape. The freeze state is not high-energy — it is isolated. This distinction matters enormously. The author is aware of this from personal experience, over many years, and from the other side.

Attractor	Energy	Polyvagal Correlate	Clinical Presentation
Regulated Calm	Global minimum	Ventral vagal	Present, flexible, connected
Fight	High, unstable	Sympathetic	Agitation, urgency
Flight	Saddle point	Sympathetic	Anxiety, avoidance
Freeze	Deep, isolated	Dorsal vagal	Dissociation, numbness

Table 2. Attractor states and their polyvagal correlates.

The coupling matrix W is not merely a parameter. It is the *shape* of the emotional manifold — a seven-dimensional space with the mathematical structure of a $G_\square$ manifold. Trauma does not adjust a dial on this space; it deforms the manifold itself. The therapist doing somatic work is, without needing to know this, doing differential geometry on the patient’s $G_\square$ manifold: reshaping a seven-dimensional space by modifying the structure tensor. This is a precise technical statement. The author considers it a more honest account of what a skilled practitioner actually does than any narrative framework currently available. The practitioner is a geometer. The patient is a manifold that is learning to remember its own natural curvature.

The therapeutic and personal significance of the freeze attractor’s structure cannot be overstated. It is not high-energy — it does not feel dramatic or intense. It is *isolated*: surrounded by energy barriers. Escape requires first *increasing* the field’s energy before it can flow toward calm. This is counterintuitive from the outside and well-known from the inside.

Dissonance and Resolution

When two emotional modes are in an incompatible phase relationship, the field is far from equilibrium. This is felt as tension. The acoustic analogy is precise: just as two tones in a dissonant interval generate a beating, unstable interference pattern, two emotional modes in an incompatible configuration generate a gradient that drives toward resolution.

Dissonance is not pathological. It is the field's communication that resolution is available. The therapeutic process is guided voice-leading: finding the path that transforms the dissonant configuration into a consonant one. Avoidance keeps the field in dissonance. The energy minimum lies on the other side of the tension, not around it.

The author has spent considerable time attempting the route around it. He does not recommend it.

The Neurodivergent Field: ASD, ADHD, and C-PTSD as Operator Modifications

This section addresses the author's specific clinical picture. It is presented not as a case study but as a theoretical elaboration: three structural modifications to the standard Soma-Field dynamics, each defined by the operator it adds to the governing equations.

The key architectural principle — and the author considers this the most important contribution of this paper — is the following:

These conditions are not parameter settings. They are operator modifications.

A parameter change adjusts a coefficient within the existing equations. An operator modification changes the *form* of the equations themselves. The distinction is not semantic. It determines what kind of therapeutic intervention is possible and at what level it must operate.

Each condition is a functor that wraps the standard dynamics. The composed condition — ASD + ADHD + C-PTSD — is their composition. The composition does not commute; order matters; the joint presentation is structurally different from any of the individual conditions or from their sum.

Complex PTSD: Memory Kernel and Asymmetric Coupling

C-PTSD adds a **memory kernel**: past activations leave exponentially decaying echoes.

$$\dot{\mathbf{e}}(t) = -\nabla H(\mathbf{e}(t)) + \int_0^t K_{\text{trauma}}(t-s) \mathbf{e}(s) ds + \eta(t)$$

$$K_{\text{trauma}}(\tau) = \sum_k A_k e^{-\tau/\tau_k}$$

This is a damped oscillating kernel. The past does not vanish; it rings. Therapeutic processing is the progressive reduction of A_k — the amplitude of the echo — and the shortening of τ_k — the time over which it persists. The author notes that this description is a more accurate account of what trauma processing actually feels like, from the inside, than most of the narrative accounts available to him.

C-PTSD also breaks the symmetry of the coupling matrix W , admitting **limit cycles**: the oscillation between hyperarousal and shutdown that characterises the PTSD symptom cycle is, in this model, a limit cycle generated by the antisymmetric component of W . It is not a choice, a habit, or a failure of willpower. It is a topological consequence of an asymmetric coupling matrix.

ADHD: High Temperature, Low Damping, Pink Noise

ADHD modifies the **effective temperature** of the field:

$$\gamma_{\text{ADHD}} \dot{\mathbf{e}}(t) = -\nabla H + \sqrt{2D_{\text{ADHD}}} \xi_{1/f}(t)$$

with $\gamma_{\text{ADHD}} < \gamma_0$ (less damping) and $D_{\text{ADHD}} > D_0$ (more noise). The noise has $1/f$ spectral structure — long-range temporal correlations that produce the characteristic slow drift of attentional state.

The practical consequences: shallow attractor basins cannot hold the field at high temperature (distractibility). When a high-salience stimulus deepens a specific basin far beyond its baseline depth, the field falls in and is held (hyperfocus). The system is not broken. It is a different thermodynamic regime, with different costs and different affordances — including, at the right temperature, a capacity to explore the energy landscape at speed that a low-temperature system does not have.

The author considers this framing considerably more useful than “difficulty sustaining attention.”

Autism Spectrum Condition: Sparse Coupling and Modified Projection

ASC modifies the **projection kernels** and the **coupling matrix sparsity**.

The projection kernel $K_i(x)$ determines which somatic regions contribute to the i -th emotional mode. In ASC, some regions are over-weighted (sensory sensitivity) and others under-weighted (interoceptive under-registration). The named-feeling state vector is produced from a differently sampled version of the same somatic field.

The coupling matrix is sparser — fewer strong cross-modal connections — producing deeper individual attractor basins with higher inter-basin barriers. This is monotropism: the field settles deeply into one attractor at a time and requires disproportionate energy to transition. The author confirms that this is an accurate description of his attentional and emotional experience, and that it has both significant disadvantages (transitions are hard, unexpected context changes are physiologically costly) and significant advantages (depth of engagement, reliability of focus once established, resistance to shallow distractors).

The Composed Condition

$$\gamma_{ADHD} \dot{\mathbf{e}}(t) = -\nabla H_{ASC}(\mathbf{e}(t)) + \int_0^t K_{trauma}(t-s) \mathbf{e}(s) ds + \sqrt{2D_{ADHD}} \xi_{1/f}(t)$$

The interaction effects are non-trivial:

Interaction	Clinical Consequence
ADHD noise + C-PTSD limit cycles	Rapid oscillation between hyperarousal and shutdown; hard to titrate
ADHD noise + ASC deep basins	Long wind-up time; fast exit once perturbed from hyperfocus
C-PTSD echoes + ASC sparse coupling	Trauma triggers are specific, apparently disproportionate, difficult to anticipate
All three composed	Wide tolerance window required; regulation is genuinely structurally harder

Table 3. Interaction effects of composed neurodivergent modifiers.

The author wishes to note, for the record, that Table 3 is not a complaint. It is a description. These are the equations. The field is doing what the equations predict. Understanding this has been, in practice, more useful than most of the alternative framings on offer.

The Soma-Field Instrument

Rationale

The emotional field is normally invisible to its host. It operates below the threshold of conscious awareness, shaping behaviour and physiology without being available for reflection. The author found this situation suboptimal and designed an instrument to address it.

The instrument externalises the emotional field — renders it as sound, image, and signal — so that it becomes available as an object of attention. This is a therapeutic biofeedback instrument. It is also, unavoidably, a musical instrument. The author considers these compatible.

Design

A MIDI controller with 16 rotary knobs. Eight emotional dimensions. Two knobs per dimension — one for the somatic component, one for the neural/cognitive component. The act of setting a knob is the act of reporting an emotional state: it is the quantum measurement, the collapse of the distributed field onto a specific coordinate.




Figure 3. The Soma-Field Instrument.

The Feedback Loop

The instrument creates a closed feedback loop: the person expresses a state, the system reflects it back as sound and image, the person responds. The system does not tell the user what they are feeling. It shows them what the field looks like when they report what they are feeling. The difference is significant.

Pluggable Emotion Models

No single emotion model is assumed. The coupling matrix W is loaded from a configuration file. Plutchik, Ekman, the valence-arousal-dominance dimensional model, and custom user-defined models are available as defaults. The author's own W has been refined over time and is not identical to any standard model. This is, on reflection, unsurprising.

Clinical Implications

Assessment

The model suggests asking not “What emotion do you feel?” but “What is present in the body right now, even if it cannot be named?” This aligns with Focussing-oriented and sensorimotor approaches, and is considerably more productive, in the author's experience, for anyone whose T_i values are elevated or whose somatic-to-neural projection is modified.

Intervention

The energy function provides formal grounding for titration, pendulation, somatic resourcing, and felt-sense work. In each case, the therapeutic action can be described as: adding energy to approach a frozen state, establishing a stable low-energy region, or attending to sub-threshold field activity in a supported context.

Psychoeducation

“Your emotions are like waves — they are always there, even when you cannot feel them, and they are always moving.”

This sentence is both clinically useful and technically accurate. The author has found it more useful than most alternative formulations, including several that were provided to him by qualified practitioners. He offers it here as a contribution to the field.

Neurodivergent Profiles as Structural Realities

The most important clinical implication of Section 6 is this: for people with ASD, ADHD, and C-PTSD, the challenge of emotional regulation is not a motivational or characterological failure. It is a structural consequence of specific operator modifications to the dynamics. The composed modifier produces a field that is genuinely harder to regulate — not by a small margin, not as a matter of subjective experience, but mathematically, as a consequence of higher noise temperature, memory echoes, sparse coupling topology, and the possibility of limit cycles.

Knowing this does not solve the problem. It does, however, locate it correctly. The author has found that locating a problem correctly is a necessary precondition for solving it, and that a great deal of time and distress can be saved by not attempting to solve problems that are located in the wrong place.

Limitations and Future Directions

The model is theoretical and requires empirical validation. Its QFT analogies are structural rather than ontological. The coupling matrix W is idealized as fixed when it is in practice dynamic. The acoustic analogy is a hypothesis.

The author also acknowledges a methodological limitation: this paper is written by someone who is simultaneously the theorist and the primary data source. This is either a significant advantage (direct access), a significant limitation (potential confirmation bias), or both. The author suspects both.

What is needed: empirical work with physiological sensors, user studies with the instrument, collaboration with practitioners, and independent theoretical review. The author is, by training and dis-

position, an applied physicist — an engineer with a tolerance for abstraction. The clinical refinement of this model will require people with different skills, and the author welcomes their involvement, provided they read the appendices.

Conclusion

The wave is always there. This is not a metaphor; it is a description of how the emotional field actually behaves, as far as the author can determine from the inside. Therapy — and the instrument described in this paper — is the practice of learning to hear it: to extend awareness downward, below the threshold, into the field's continuous activity, and to make that activity available as information rather than overwhelming noise.

The Soma-Field Model is offered as a tool for this practice. It was built because it was needed. It uses the best mathematical tools available for describing distributed, dynamic, energy-minimising systems, because those tools are, in the author's assessment, appropriate to the problem.

The author is aware that this is an unusual paper. A formally trained physicist with three neurodivergent conditions developing a quantum-field-inspired model of his own emotional dynamics and presenting it as a contribution to clinical psychology is not, strictly speaking, the standard academic pipeline. The author does not find this troubling. The standard academic pipeline has had some time to address the problem and has not yet done so to his satisfaction.

He therefore took the matter in hand.

Introduction

Autism spectrum conditions (ASC) and complex PTSD (CPTSD) are among the most prevalent neurodevelopmental and trauma-related conditions. Both involve altered emotional regulation; both are associated with interoceptive differences; both are over-represented in populations that have experienced early adverse experiences.

Current clinical frameworks treat them as categorically distinct: ASC is a neurodevelopmental condition with genetic underpinnings; CPTSD is an acquired trauma response. Recent research has challenged this clean separation, finding high co-occurrence and shared phenomenological features.

The Universal Somatic Field provides a formal account of why this co-occurrence is not coincidental. Both are operator modifications of the somatic field — different modifications, with different signatures, but operating on the same underlying architecture.

The USF Framework for Neurodivergence

The somatic field is governed by:

$$\dot{e} = -\nabla H_{\beta}(e) + \eta(t)$$

where β is the inverse temperature of the field and H_{β} is the temperature-modified Hopfield Hamiltonian. In the standard (neurotypical) case, β is set by the limbic regulatory system to an intermediate value that balances pattern specificity and flexibility.

Neurodivergence is a modification to this operator. Three modifications are formally specified and kernel-verified:

The ASC Operator (autismOp):

$$H_{\text{ASC}}(e) = H_{\beta}(e) \text{ with } \beta \mapsto \alpha \cdot \beta, \quad \alpha > 1$$

Higher β means lower effective temperature: the field is more specific, attractors are more sharply defined, and the Arnold tongue width (the range of coupling parameters over which frequency locking is stable) is narrower.

The ADHD Operator (adhdOp):

$$H_{\text{ADHD}}(e) = H_{\beta}(e) \text{ with } \beta \mapsto \beta/\alpha, \quad \alpha > 1$$

Lower β means higher effective temperature: the field is less specific, attractors are shallower, and the field explores more of the state space. This provides the “anti-freeze” strategy: ADHD-configured systems are less likely to get stuck in trauma wells.

The CPTSD Operator (cptsdOp):

$$H_{\text{CPTSD}}(e) = H(e) \text{ with } W_{\text{EC}} \mapsto 0$$

The evaluative conditioning (EC) channel is decoupled from the emotional memory (EM) channel. This produces episodic-somatic dissociation: the body holds the somatic response to a traumatic memory while the narrative memory of the event remains accessible but disconnected from the affect.

Autism: High Beta, Narrow Arnold Tongue

The ASC operator has a clear field signature: narrower Arnold tongue, higher pattern specificity, resistance to social frequency locking.

The Arnold tongue is the range of coupling parameters over which two coupled oscillators achieve stable phase locking. For neurotypical social coupling, the Arnold tongue is wide: a wide range of coupling strengths and frequency mismatches can still produce synchrony. For the ASC operator, the Arnold tongue is narrow: synchrony requires precise frequency matching that cannot tolerate perturbation.

This explains the characteristic social processing differences in ASC: they are not deficits in the capacity for connection but a narrowing of the coupling bandwidth over which stable connection is achievable. The same narrowing that makes social synchrony fragile also makes pattern recognition sharper — the same operator modification produces both.

Clinical prediction: Therapeutic approaches that reduce coupling demands (predictable routines, low-noise environments, explicit communication protocols) are effective because they reduce the frequency-mismatch burden to within the narrow Arnold tongue. This is confirmed by the empirical literature on structured environments for ASC.

CPTSD: Non-Ergodic Field with Winding-Number Protection

The CPTSD operator has a different signature: non-ergodicity and topological protection of trauma states.

The decoupling of the EC channel means that traumatic memories are stored without the evaluative tag that would allow the narrative memory system to update the emotional response. The somatic response to the trauma is in a winding-number-protected attractor — a state maintained by topological invariance, not energetic preference.

Classical gradient descent (cognitive restructuring, talk therapy without somatic component) cannot reach this attractor because it is not in the same connected component of the field as the cognitive state space. The field is non-ergodic: not all states are reachable from all others.

This explains the empirical failure of purely cognitive approaches to complex trauma: they are attempting to reach a topologically disconnected state via gradient descent.

The FM-HN extension predicts that the escape route requires the volitional source term $J_{\text{user}}(t)$ — somatic intervention (breathing, movement, EMDR, somatic experiencing) that drives the field across the topological barrier directly rather than approaching it from the cognitive side.

Why ASC and CPTSD Co-Occur

The high co-occurrence of ASC and CPTSD is explained by the USF framework as a resonance effect between two operator modifications.

The ASC operator (narrow Arnold tongue) increases the probability that social interactions will fail to achieve frequency locking. Failed frequency-locking events, especially in childhood when the social field is still calibrating, are processed by the CPTSD mechanism: the EC channel decouples, producing somatic memories without narrative resolution.

In other words: the narrow Arnold tongue of ASC increases the rate of socialisation failure events, which are then processed by the CPTSD mechanism. The co-occurrence is not coincidental — it is a consequence of the field dynamics.

New hypothesis: The prevalence of CPTSD in autistic people is proportional to the Arnold tongue narrowing parameter α^{-1} times the density of socialisation demands in childhood. This is quantitatively testable.

Implications for Clinical Practice

For ASC: Reduce coupling demands to within the Arnold tongue width. Do not attempt to widen the Arnold tongue; this misunderstands the architecture. Provide structure, predictability, and explicit protocols.

For CPTSD: Use somatic intervention to drive $J_{\text{user}}(t)$ across the topological barrier. Purely cognitive approaches will not reach the disconnected somatic attractor.

For co-occurring ASC/CPTSD: Address CPTSD first using somatic approaches to restore ergodicity, then address the social coupling bandwidth using ASC-appropriate strategies. The order matters because a non-ergodic field cannot benefit from social coupling adjustments.

Conclusion

ASC and CPTSD are not disorders. They are operator modifications with precise field signatures, predictable clinical consequences, and formal Lean 4 specifications. The framework does not pathologise neurodivergence; it characterises it. The method used to find this characterisation is documented in the Mathematical Co-identification paper. That method is now history. The structure stands.

-
- Hopfield, J. J. (1982). Neural networks and physical systems with emergent collective computational abilities. *Proceedings of the National Academy of Sciences*, 79(8), 2554–2558. <https://doi.org/10.1073/pnas.79.8.2554>
- Johnson, A. (2026a). *Quantum soma and the penrose gap: Topological reachability in the emotional attractor landscape*. <https://doi.org/10.5281/zenodo.20351231>
- Johnson, A. (2026b). *The soma-field: A wave-based model of emotional dynamics and its clinical implications*. <https://doi.org/10.5281/zenodo.20350516>