Gaussian Fermions

E. Miles Stoudenmire[1]

[1] *Center for Computational Quantum Physics, Flatiron Institute, 162 5th Avenue, New York, NY 10010, USA*

I. FERMIONIC VECTOR SPACES

Fermions are conventionally defined in terms of creation and annihilation operators. Here we will start from state vectors, or fermionic vector spaces. In the next section, we will see that the properties of the operators derive from the properties of the states.

The simplest fermionic vector space V has the basis

$$\mathcal{B}(V) = \{|0\rangle, |1\rangle\} \tag{1}$$

which is the basis for a single "spinless fermion," with $|0\rangle$ being the vacuum or empty state and $|1\rangle$ being the filled state or state of one fermion. So far the fermionic nature is not very obvious except that we can only have at most one fermion.

Many-body spaces can be formed by taking Cartesian products of spaces (tensor products of basis vectors) so we could consider a space V_1 with basis $\{|0\rangle_1, |1\rangle_1\}$ and space V_2 with basis $\{|0\rangle_2, |1\rangle_2\}$, then the space $V_1 \times V_2$ has a basis

$$\mathcal{B}(V_1 \times V_2) = \{|0\rangle_1 \otimes |0\rangle_2,\ |1\rangle_1 \otimes |0\rangle_2,\ |0\rangle_1 \otimes |1\rangle_2,\ |1\rangle_1 \otimes |1\rangle_2\}\ . \tag{2}$$

A generic vector (or "state") in this space has the form $|\psi\rangle = a\,|0\rangle_1 \otimes |0\rangle_2 + b\,|0\rangle_1 \otimes |1\rangle_2 + c\,|1\rangle_1 \otimes |0\rangle_2 + d\,|1\rangle_1 \otimes |1\rangle_2$.

The defining property of fermions is how one identifies vectors in the space $V_1 \times V_2$ with vectors in the space $V_2 \times V_1$. We can define this identification, or vector space isomorphism, by defining it on the basis vectors. It is given by:

$$\begin{aligned}
|0\rangle_1 \otimes |0\rangle_2 &\leftrightarrow +\,|0\rangle_2 \otimes |0\rangle_1 \\
|1\rangle_1 \otimes |0\rangle_2 &\leftrightarrow +\,|0\rangle_2 \otimes |1\rangle_1 \\
|0\rangle_1 \otimes |1\rangle_2 &\leftrightarrow +\,|1\rangle_2 \otimes |0\rangle_1 \\
|1\rangle_1 \otimes |1\rangle_2 &\leftrightarrow -\,|1\rangle_2 \otimes |1\rangle_1 \quad .
\end{aligned} \tag{3}$$

Note the minus sign in the last line. Plus signs are shown for emphasis.

We can summarize the vector space isomorphism as

$$|n_1\rangle_1 \otimes |n_2\rangle_2 \ \leftrightarrow\ (-1)^{n_1 \cdot n_2}\,|n_2\rangle_2 \otimes |n_1\rangle_1 \quad . \tag{4}$$

This means the ordering of fermionic basis vectors matters: if two odd basis vectors are permuted, the basis vector changes by a minus sign. [1]

Note that tensor products of basis states such as $|1\rangle_i \otimes |1\rangle_j$ are commonly written in the more compact notation $|1\rangle_i\,|1\rangle_j$ or even $|1_i 1_j\rangle$. Tensor product symbols "$\otimes$" are thus generally dropped when understood from context, and will be dropped frequently in the rest of these notes.

II. FERMIONIC OPERATORS

For the simplest case of a "spinless fermion" space V with basis $\mathcal{B}(V) = \{|0\rangle, |1\rangle\}$, we can define the following operators:

$$\hat{n} = |1\rangle\langle 1| \tag{5}$$

$$\hat{c} = |0\rangle\langle 1| \tag{6}$$

$$\hat{c}^\dagger = |1\rangle\langle 0| \quad . \tag{7}$$

Together with the identity operator, this set actually defines a complete operator basis.

The $\hat{n}$ operator is known as the "number operator" since it "detects" the filling of a state:

$$\begin{aligned}
\hat{n}\,|0\rangle &= 0 \cdot |0\rangle \\
\hat{n}\,|1\rangle &= 1 \cdot |1\rangle \quad .
\end{aligned} \tag{8}$$

The $\hat{c}$ and $\hat{c}^\dagger$ are known as the "annihilation" and "creation" operators, respectively. This is simply because they map a filled state to an empty state or vice versa:

$$\hat{c}\,|1\rangle = |0\rangle \tag{9}$$
$$\hat{c}^\dagger\,|0\rangle = |0\rangle \quad . \tag{10}$$

One can of course define tensor products of operators in the usual way, such as $\hat{c}_1\hat{c}_2$ or $\hat{c}_2^\dagger\hat{c}_1$. [2] These operators act on the spaces $V_1 \times V_2$ or $V_2 \times V_1$, respectively.

Now comes a crucial point: the creation and annihilation operators have non-trivial commutation relations, and these are entirely a consequence of the state properties. For operators on state spaces V_i and V_j (with $i \neq j$), the commutation relations are

$$\hat{c}_i\hat{c}_j = -\hat{c}_j\hat{c}_i \tag{11}$$
$$\hat{c}_i^\dagger\hat{c}_j^\dagger = -\hat{c}_j^\dagger\hat{c}_i^\dagger \tag{12}$$
$$\hat{c}_i\hat{c}_j^\dagger = -\hat{c}_j^\dagger\hat{c}_i \tag{13}$$

The first two of these are commonly summarized as:

$$\begin{aligned}\{\hat{c}_i, \hat{c}_j\} &= 0\\ \{\hat{c}_i^\dagger, \hat{c}_j^\dagger\} &= 0\end{aligned} \tag{14}$$

where $\{a, b\} = a\,b + b\,a$ is the anti-commutator.

The third relation Eq. (13) has the special case that when $i = j$, $(\hat{c}_i\hat{c}_i^\dagger + \hat{c}_i^\dagger\hat{c}_i) = 1$. Therefore it is commonly summarized as

$$\{\hat{c}_i, \hat{c}_j^\dagger\} = \delta_{ij} \quad . \tag{15}$$

In many presentations, the anticommutation relations of the operators are stated as an axiom, but they actually follow from the properties of the state spaces. For example,

$$\begin{aligned}\hat{c}_1\hat{c}_2 &= |0\rangle_1 \otimes \langle 1|_1 \otimes |0\rangle_2 \otimes \langle 1|_2\\ &= |0\rangle_1 \otimes (\langle 1|_1 \otimes \langle 1|_2) \otimes |0\rangle_2\\ &= |0\rangle_1 \otimes (-\langle 1|_2 \otimes \langle 1|_1) \otimes |0\rangle_2\\ &= -|0\rangle_2 \otimes \langle 1|_2 \otimes \langle 0|_1 \otimes |1\rangle_1\\ &= -\hat{c}_2 \otimes \hat{c}_1 \quad .\end{aligned} \tag{16}$$

III. MANY-BODY STATES AND GAUSSIAN STATES

Many-body or many-particle basis states can be defined or constructed as

$$|1_i 1_j 1_k\rangle = \hat{c}_i^\dagger\hat{c}_j^\dagger\hat{c}_k^\dagger\,|0\rangle \quad . \tag{17}$$

The most general many-particle state is a sum of these:

$$|\Psi\rangle = \sum_{n_1, n_2, \ldots, n_N} \Psi^{n_1 n_2 \cdots n_N}\,|n_1 n_2 \cdots n_N\rangle \tag{18}$$

where $\Psi^{n_1 n_2 \cdots n_N}$ is an arbitrary tensor of complex amplitudes. The anti-commuting nature of the fermions is automatically satisfied by the properties of the basis, and no additional symmetry properties need to be assumed for the amplitudes $\Psi^{n_1 n_2 \cdots n_N}$.

A typical extra assumption, though, is that for a closed system the total number of particles N_p is fixed. Mathematically this means $n_1 + n_2 + \ldots + n_N = N_p$ otherwise $\Psi^{n_1 n_2 \cdots n_N} = 0$. (Superconducting states are a common exception to this rule, though these are probably more properly understood as mixed states.)

A special subset of many-particle fermionic states are the *Gaussian states*, more commonly known as *Slater determinants*. The general form of a Gaussian state with four fermions is:

$$\begin{aligned}|\Psi\rangle &= \hat{A}^\dagger \hat{B}^\dagger \hat{C}^\dagger \hat{D}^\dagger |0\rangle \\ &= \left(\sum_{i_1=1}^{N} A^{i_1} \hat{c}_{i_1}^\dagger\right)\left(\sum_{i_2=1}^{N} B^{i_2} \hat{c}_{i_2}^\dagger\right)\left(\sum_{i_3=1}^{N} C^{i_3} \hat{c}_{i_3}^\dagger\right)\left(\sum_{i_4=1}^{N} D^{i_4} \hat{c}_{i_4}^\dagger\right)|0\rangle \quad .\end{aligned} \tag{19}$$

Above we have introduced operators such as $\hat{A} = \sum_{i_1=1}^{N} A^{i_1} \hat{c}_{i_1}^\dagger$. One describes the state above by saying that it contains four fermions which are in the "orbitals" A, B, C and D. These orbitals are vectors $\vec{A} = [A^1\ A^2\ \ldots A^N]$ which are unit-normalized. The vectors are not required to be orthogonal, though the properties of the state can be simpler to deduce if they are.

The Gaussian states are quite special and far from being a general fermionic state, even though they are elements of the general vector space of fermionic many-body states. To see that the Gaussian states are not general, let us write a two-particle Gaussian state $|AB\rangle = \hat{A}^\dagger \hat{B}^\dagger |0\rangle$ in the general form Eq. (18) for a fermionic state. We see that for the state $|AB\rangle$, the amplitudes $\Psi^{n_1 n_2 \cdots n_N}$ take the form:

$$\Psi^{0_1 0_2 \cdots 1_i \cdots 1_j \cdots 0_N} = A_i B_j \quad \text{for } \sum_i n_i = 2 \tag{20}$$

$$\Psi^{n_1 n_2 \cdots n_N} = 0 \qquad \text{otherwise for } \sum_i n_i \neq 2 \quad . \tag{21}$$

So the amplitude tensor Ψ of $|AB\rangle$ is constructed from two vectors (it is rank-2 in a certain sense [3]) and is not fully general.

To understand the origin of the name Slater determinant, consider the basis defined as $|i\rangle = \hat{c}_i^\dagger |0\rangle$, the so-called "real-space basis". Expressing states in this basis is known as *first quantization*. We can express the amplitudes of $|AB\rangle$ in this basis, as $\psi_{AB}(i,j) = \langle i,j|AB\rangle = \langle 0| c_i^\dagger c_j^\dagger |AB\rangle$. One finds that

$$\psi_{AB}(i,j) = A^i B^j - B^i A^j \tag{22}$$

such that the amplitudes in real space take the form of an antisymmetric function. It was pointed out by Slater that expressions of the above type can be written as determinants of matrices whose columns are the orbital vectors $\vec{A}, \vec{B}, \ldots$ and whose rows are the spatial positions $i = 1, 2, \ldots, N$, hence the name Slater determinant.

The first-quantized basis can be convenient for proving facts such as if $\vec{A} = \vec{B}$, then the state $|AB\rangle = 0$, as is evident from Eq. (22) above. In the more general setting, the requirement that orbitals required be independent is known as the *Pauli exclusion principle*. It is not a separate axiom about fermions, but follows from their vector space properties.

IV. STATIC PROPERTIES OF GAUSSIAN STATES

Define g operators.

The most generic form of a Gaussian state or Slater determinant is

$$|m_1 m_2 \ldots m_n\rangle_g = \hat{g}_{m_1}^\dagger \hat{g}_{m_2}^\dagger \cdots \hat{g}_{m_n}^\dagger |0\rangle \tag{23}$$

$$= \prod_{m=1} (\hat{g}_m^\dagger)^{\nu_m} |0\rangle \tag{24}$$

where ν_m is the *occupancy* of state m in the g basis. The occupancy is defined here as $\nu_m = 1$ if $m \in \{m_1, m_2, \ldots, m_n\}$ and $\nu_m = 0$ otherwise.

V. DYNAMICS OF GAUSSIAN STATES

VI. MATHEMATICAL RESULTS

Correlation functions and expectation values.

- *Correlation matrix:* Given a Gaussian state $|g\rangle = \hat{g}^\dagger_{m_1}\hat{g}^\dagger_{m_2}\cdots\hat{g}^\dagger_{m_n}|0\rangle$ where $\hat{g}^\dagger_m = g^i_m\hat{f}^\dagger_i$, the correlation matrix C^j_i in the f basis is given by $C^j_i = \langle g|\,\hat{f}^\dagger_i\hat{f}^j\,|g\rangle = \sum_n \bar{g}^n_i\nu_n g^j_n$, where $\nu_m = \langle g|\,\hat{n}_m\,|g\rangle$ is the occupancy of state m in the g basis ($\nu_m = 1$ if $m \in \{m_1, m_2, m_n\}$, $\nu_m = 0$ otherwise).

- *Same-basis correlation matrix:* Given a Gaussian state $|f\rangle = \hat{f}^\dagger_{j_1}\hat{f}^\dagger_{j_2}\cdots\hat{f}^\dagger_{j_n}|0\rangle$, the correlation matrix C^j_i in the f basis is given by $C^j_i = \langle f|\,\hat{f}^\dagger_i\hat{f}^j\,|f\rangle = \delta^i_j\nu_j$ where $\nu_j = \langle f|\,\hat{n}_j\,|f\rangle$ is the occupancy of orbital j in the f basis.

- *Expectation value:* Given an operator $\hat{M} = m^i_j\hat{c}^\dagger_i c^j$ and a Gaussian state $|f\rangle$ with correlation matrix $C^j_i = \langle f|\,\hat{c}^\dagger_i\hat{c}^j\,|f\rangle$, the expected value of $\hat{M}$ is $\langle\hat{M}\rangle = \mathrm{Tr}[mC]$.

Action of operators.

- *Action of creation operator:* Given state $|g\rangle = \hat{g}^\dagger_{m_1}\hat{g}^\dagger_{m_2}\cdots\hat{g}^\dagger_{m_n}|0\rangle$ where $\hat{g}^\dagger_m = g^i_m\hat{f}^\dagger_i$ and an operator $\hat{v}^\dagger = v^j\hat{f}^\dagger_i$, the correlation matrix of the state $|g'\rangle = \hat{v}^\dagger|g\rangle$ is given by $C' = ||v_0||^2C + v_0v_0^\dagger$, where $v_0 = (1-C)v$ is the projection of v onto the unoccupied subspace of C.

- *Action of annihilation operator:* Given state $|g\rangle = \hat{g}^\dagger_{m_1}\hat{g}^\dagger_{m_2}\cdots\hat{g}^\dagger_{m_n}|0\rangle$ where $\hat{g}^\dagger_m = g^i_m\hat{f}^\dagger_i$ and an operator $\hat{w} = \bar{w}_i\hat{f}^i$, the correlation matrix of the state $|g'\rangle = \hat{w}|g\rangle$ is given by $C' = ||w_1||^2C - w_1w_1^\dagger$, where $w_1 = Cw$ is the projection of w onto the occupied subspace of C.

Dynamics. All of these results assume the Hamiltonian $\hat{H} = h^i_j\hat{c}^\dagger_i\hat{c}^j$ has a hopping matrix which can be diagonalized as $h^i_j = \phi^i_n\epsilon_n\phi^n_j$. The ground state of $\hat{H}$ is $|\Phi_0\rangle$.

- *Heisenberg time evolution of operators:* Define the operators $\hat{\phi}^\dagger_n = \sum_n\phi^j_n\hat{c}^\dagger_j$ and $\hat{\phi}^n = \sum_n\bar{\phi}^n_j\hat{c}^j$. These evolve in the Heisenberg picture as $\hat{\phi}^n(t) = e^{i\hat{H}t}\hat{\phi}^n e^{-i\hat{H}t} = e^{-i\epsilon_n t}\hat{\phi}^n$ and $\hat{\phi}^\dagger_n(t) = e^{i\epsilon_n t}\hat{\phi}^\dagger_n$.

- *Lesser Greens function:* Defined as $G^{<\,i}_j(t) = i\langle\Phi_0|\,\hat{c}^\dagger_j\hat{c}^i(t)\,|\Phi_0\rangle$, it is given by $G^{<\,i}_j(t) = i\sum_n\nu_n\phi^i_n e^{-i\epsilon_n t}\bar{\phi}^n_j$ where ν_n is the occupancy of the ground state in the ϕ basis.

- *Greater Greens function:* Defined as $G^{>\,i}_j(t) = -i\langle\Phi_0|\,\hat{c}^i(t)\hat{c}^\dagger_j\,|\Phi_0\rangle$, it is given by $G^{>\,i}_j(t) = -i\sum_n(1-\nu_n)\phi^i_n e^{-i\epsilon_n t}\bar{\phi}^n_j$ where ν_n is the occupancy of the ground state in the ϕ basis.

- *Retarded Greens function:* Defined as $G^{R\,i}_j(t) = \theta(t)(G^{>\,i}_j(t) - G^{<\,i}_j(t))$, it is given by $G^{R\,i}_j(t) = -i\theta(t)\sum_n\phi^i_n e^{-i\epsilon_n t}\bar{\phi}^n_j = -i\theta(t)e^{-iht}$.

- *Propagator:* Define the propagator as $g^i_j(t) = i(G^{>\,i}_j(t) - G^{<\,i}_j(t))$. It can be written simply as $g^i_j(t) = e^{-iht} = \sum_n\phi^i_n e^{-i\epsilon_n t}\bar{\phi}^n_j$.

- *Evolution of correlation matrix:* For a pure, Gaussian state $|f\rangle$ with correlation matrix $C^j_i = \langle f|\,\hat{c}^\dagger_i\hat{c}^j\,|f\rangle = \bar{f}^n_i\nu_n f^j_n$, the orbitals evolve, in the Schrödinger picture, as $f^j_n(t) = g^i_j(t)f^j_n$. So $C^j_i(t) = \bar{g}^{i'}_i(t)C^{j'}_{i'}g^j_{j'}(t)$.

---

[1] In some mathematical literature, an isomorphism of this type is described as "having a $\mathbb{Z}_2$ grading" (meaning that one assigns non-trivial properties based on the parity or grading of the basis) or having a "braided" or "graded" vector space isomorphism.

[2] When an operator like $\hat{c}_1$ is written in isolation, it actually stands for $\hat{c}_1\otimes\hat{I}_2$. Products of operators such as $\hat{c}^\dagger_1\hat{c}_2$ are actually shorthand for $(\hat{c}^\dagger_1\otimes\hat{I}_2)(\hat{I}_1\otimes\hat{c}_2) = \hat{c}^\dagger_1\otimes\hat{c}_2$. If the operators act on the same space, they "stack", meaning for example that $(\hat{c}^\dagger_1\otimes\hat{I}_2)(\hat{c}_1\otimes\hat{I}_2) = (\hat{c}^\dagger_1\hat{c}_1)\otimes\hat{I}_2$. All of this is easier to see with tensor diagrams!

[3] The amplitude tensor for $|AB\rangle$ or a general two-particle Gaussian state is actually a bond-dimension 2 or rank 2 matrix product state.