

Hodgkin-Huxley Neuron Model Equations (with Extensions)

1 Membrane Potential Dynamics

The voltage dynamics for a single-compartment neuron is described by:

$$C_m \frac{dV}{dt} = -(I_L + I_{Na} + I_K + I_{CaL} + I_{CaT} + I_{AHP} + I_H + I_{Ks}) + I_{ext} + I_{noise}$$

where C_m is the membrane capacitance, I_{ext} is external current, and I_{noise} is noise current.

2 Ionic Currents

Belows are equations for various ionic currents in the neuron model:

$$I_L = g_L(V - V_L) \quad (1)$$

$$I_{Na} = g_{Na} m_{Na}^3 h_{Na} (V - V_{Na}) \quad (2)$$

$$I_K = g_K n_K^4 (V - V_K) \quad (3)$$

$$I_{CaL} = g_{CaL} m_{CaL}^2 (V - V_{CaL}) \quad (4)$$

$$I_{CaT} = g_{CaT} (m_{CaT}^\infty)^2 h_{CaT} (V - V_{CaT}) \quad (5)$$

$$I_{AHP} = g_{AHP} x_{AHP} (V - V_{AHP}) \quad (6)$$

$$I_{CAN} = g_{CAN} x_{CAN} (V - V_{CAN}) \quad (7)$$

$$I_{Ks} = g_{Ks} m_{Ks} h_{Ks} (V - V_{Ks}) \quad (8)$$

$$I_H = g_H m_H (V - V_H) \quad (9)$$

Gating and Activation Variables

Sodium Channel (Na)

$$m_{Na}(V) = \frac{1}{1 + \exp\left(-\frac{V+30}{9.5}\right)} \quad (10)$$

$$h_{Na,\infty}(V) = \frac{1}{1 + \exp\left(\frac{V+53}{7}\right)} \quad (11)$$

$$\tau_h(V) = 0.37 + 2.78 \cdot \frac{1}{1 + \exp\left(\frac{V+40.5}{6}\right)} \quad (12)$$

Potassium Channel (K)

$$n_{K,\infty}(V) = \frac{1}{1 + \exp\left(-\frac{V+30}{10}\right)} \quad (13)$$

$$\tau_n(V) = 0.37 + 1.85 \cdot \frac{1}{1 + \exp\left(\frac{V+27}{15}\right)} \quad (14)$$

L-type Calcium Channel (CaL)

$$m_{CaL,\infty}(V) = \frac{1}{1 + \exp\left(-\frac{V+12}{7}\right)} \quad (15)$$

$$\tau_{CaL}(V) = 10^{(0.6-0.02 \cdot V)} \quad (16)$$

T-type Calcium Channel (CaT)

$$m_{CaT,\infty}(V) = \frac{1}{1 + \exp\left(-\frac{V+57}{6.2}\right)} \quad (17)$$

$$h_{CaT,\infty}(V) = \frac{1}{1 + \exp\left(\frac{V+81}{4}\right)} \quad (18)$$

$$\tau_{mT}(V) = 0.612 + \frac{1}{\exp\left(\frac{V+132}{-16.7}\right) + \exp\left(\frac{V+16.8}{18.2}\right)} \quad (19)$$

$$\tau_{hT}(V) = \begin{cases} \exp\left(\frac{V+467}{66.6}\right) & \text{if } V < -80 \\ \exp\left(-\frac{V+22}{10.5}\right) + 28 & \text{otherwise} \end{cases} \quad (20)$$

Hyperpolarization-Activated Channel (H)

$$m_{H,\infty}(V) = \frac{1}{1 + \exp\left(\frac{V - V_{\tau,peak}}{k_{\tau}}\right)} \quad (21)$$

$$\tau_{mH}(V) = \tau_{min} + \frac{\tau_{diff}}{\exp\left(\frac{V - V_{\tau,peak}}{k_{\tau}}\right) + \exp\left(-\frac{V - V_{\tau,peak}}{k_{\tau}}\right)} \quad (22)$$

CAN Channel

$$x_{CAN,\infty}([Ca^{2+}]) = \frac{a_{CAN} \cdot [Ca^{2+}]}{a_{CAN} \cdot [Ca^{2+}] + b_{CAN}} \quad (23)$$

$$\tau_{xCAN}([Ca^{2+}]) = \frac{1}{a_{CAN} \cdot [Ca^{2+}] + b_{CAN}} \quad (24)$$

AHP Channel

$$x_{AHP,\infty}([Ca^{2+}]) = \frac{a_{AHP} \cdot [Ca^{2+}]}{a_{AHP} \cdot [Ca^{2+}] + b_{AHP}} \quad (25)$$

$$\tau_{xAHP}([Ca^{2+}]) = \frac{1}{a_{AHP} \cdot [Ca^{2+}] + b_{AHP}} \quad (26)$$

Slow Potassium Channel (Ks)

$$m_{Ks,\infty}(V) = \frac{1}{1 + \exp\left(-\frac{V+44}{5}\right)} \quad (27)$$

$$h_{Ks,\infty}(V) = \frac{1}{1 + \exp\left(\frac{V+74}{9.3}\right)} \quad (28)$$

$$\tau_{hKs}(V) = 200 + \frac{4800}{1 + \exp\left(-\frac{V+50}{9.3}\right)} \quad (29)$$

General Form

All gating variables follow first-order kinetics:

$$\tau_x \frac{dx}{dt} = (x_{\infty} - x) \quad (30)$$

where x represents the gating variable, x_{∞} is its steady-state value, and τ_x is its voltage-dependent or calcium-dependent time constant.