

The Role of Cell Size on bacterial community assembly

Yingcai Hu

May 17, 2023

1 Introduction

2 Methods

2.1 Defining bacterial taxa and resource types

In this study, each bacterial taxon i are defined by their resource preference and average cell mass m_i . The preferred resources for each taxa are chosen uniformly

2.2 Microbial Consumer-Resource Model (MiCRM)

Here we adopt the framework from @iiiAvailableEnergyFluxes2019: the energy per unit time or, The metabolic rate, v_{ij}^{in} of species i using resource j depends on the concentration of resource j , R_j and species preference on a particular resource type p_{ij} :

$$v_{ij}^{in} = \sigma(p_{ij}R_j) = \frac{v_{max}p_{ij}R_j}{R_h + R_j} \quad (1)$$

where the function $\sigma(x) = x_{\max} \frac{x}{k+x}$ is the Monod function that maps the resource available to the resources taken by bacteria. The total mass, v_i^{grow} , used for growth of biomass will be defined by setting a fraction (l_j) of mass returned to the environment:

$$v_i^{grow} = \sum_{j=1}^M (1 - l_j) v_{ij}^{in} \quad (2)$$

Therefore, assuming there are N species and M types of resources, the dynamics of biomass abundance C_i of species i^{th} and resource concentration R_j of type j^{th} is

$$\frac{dC_i}{dt} = \mu C_i (v_i^{grow} - \phi_i) \quad (3)$$

$$\frac{dR_j}{dt} = \rho_j - k_m \sum_{i=1}^N C_i (v_{ij}^{out} - v_{ij}^{in}) \quad (4)$$

Here, U_{ij} is uptake rate of resource j by species i , l_{ij} is the leakage of resource j to the environment in the form of resource k . m_i is the maintenance required for species i . The terms are explained in Table 1.

Table 1: Definition and Units of Parametres

Symbols	Definition	Units
C	Biomass content	<i>mass</i>
R	Resources content	<i>mass</i>
U	Uptake rate	<i>time</i> ⁻¹
l	Fraction of leakage	<i>None</i>
m	Maintenance coefficient	<i>time</i> ⁻¹
ρ	External resource supply	<i>mass/time</i>
μ	Constant that scales the resource uptake to growth	<i>mass</i> ⁻¹
k_m	Constant that scales the maintenance	<i>mass</i> ⁻¹
k_{ab}	Constant that scales the body mass and resource intake	<i>mass</i> ⁻¹

Here v_{ij} , v^{grow} and ϕ_i are scaled by biomass for investigating the effect of biomass on population dynamics within the bacterial community.

2.3 Body mass Parametrisation on MiCRM

2.4 Linear Stability Analysis

Let $x = [C_1, C_2, \dots, C_N, R_1, R_2, \dots, R_M] \in S$, then $x^* \in S$ is the steady state (i.e. $f'_n(x^*) = 0$ and $f'_{N+m}(x^*) = 0$ for $n = 1, 2, 3, \dots, N, m = 1, 2, 3, \dots, N$)

. The jacobian matrix of the MiRCM will be defined as followed:

$$J = \begin{bmatrix} \frac{\partial f_1}{\partial C_1} & \frac{\partial f_1}{\partial C_2} & \cdots & \frac{\partial f_1}{\partial C_N} & \frac{\partial f_1}{\partial R_1} & \frac{\partial f_1}{\partial R_2} & \cdots & \frac{\partial f_1}{\partial R_M} \\ \frac{\partial f_2}{\partial C_1} & \frac{\partial f_2}{\partial C_2} & \cdots & \frac{\partial f_2}{\partial C_N} & \frac{\partial f_2}{\partial R_1} & \frac{\partial f_2}{\partial R_2} & \cdots & \frac{\partial f_2}{\partial R_M} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_N}{\partial C_1} & \frac{\partial f_N}{\partial C_2} & \cdots & \frac{\partial f_N}{\partial C_N} & \frac{\partial f_N}{\partial R_1} & \frac{\partial f_N}{\partial R_2} & \cdots & \frac{\partial f_N}{\partial R_M} \\ \frac{\partial f_{N+1}}{\partial C_1} & \frac{\partial f_{N+1}}{\partial C_2} & \cdots & \frac{\partial f_{N+1}}{\partial C_N} & \frac{\partial f_{N+1}}{\partial R_1} & \frac{\partial f_{N+1}}{\partial R_2} & \cdots & \frac{\partial f_{N+1}}{\partial R_M} \\ \frac{\partial f_{N+2}}{\partial C_1} & \frac{\partial f_{N+2}}{\partial C_2} & \cdots & \frac{\partial f_{N+2}}{\partial C_N} & \frac{\partial f_{N+2}}{\partial R_1} & \frac{\partial f_{N+2}}{\partial R_2} & \cdots & \frac{\partial f_{N+2}}{\partial R_M} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_{N+M}}{\partial C_1} & \frac{\partial f_{N+M}}{\partial C_2} & \cdots & \frac{\partial f_{N+M}}{\partial C_N} & \frac{\partial f_{N+M}}{\partial R_1} & \frac{\partial f_{N+M}}{\partial R_2} & \cdots & \frac{\partial f_{N+M}}{\partial R_M} \end{bmatrix}$$

The f_n and f_{N+m} are vector-valued function that returns Equation 3 and Equation 4, therefore the elements in J are defined as:

$$\frac{\partial f_n}{\partial C_n} = \mu(v_n^{\text{grow}} - \phi_n) \quad (5)$$

$$\frac{\partial f_n}{\partial R_m} = \mu C_n (1 - l_m) \frac{v_{nm}^{\text{max}} p_{nm} R_m}{(R_h + p_{nm} R_m)^2} \quad (6)$$

$$\frac{\partial f_{N+m}}{\partial C_n} = k_m C_n (v_{nm}^{\text{out}} - v_{nm}^{\text{in}}) \quad (7)$$

$$\frac{\partial f_{N+m}}{\partial R_m} = k_m \sum_{i=1}^N C_i (D_{mm} l_m - 1) \frac{v_{nm}^{\text{max}} p_{nm} R_m}{(R_h + p_{nm} R_m)^2} \quad (8)$$

To investigate the stability of the system on steady state x^* , eigenvalue decomposition will be performed on the jacobian evaluated at x^* , i.e., $J|_{x=x^*} = P^{-1} \Lambda P$, using numpy library.