

Essentials of Analytical Geometry and Linear Algebra. Lecture 12.

Vladimir Ivanov

Innopolis University

November 27, 2020

Main question for today's lecture

Quadratic surfaces in 3D or **Quadrics**

Lecture 12. Outline

- Part 1. Types of Quadrics and their equations
- Part 2. Cylindrical / Spherical coordinates.
- Part 3. Applications

Part 1. Types of Quadrics and their equations

Part 1. Types of Quadrics and their equations



Part 1. Types of Quadrics and their equations

Well, it is still a quadratic equation, but with 3 variables:

x, y, z

Part 1. Types of Quadrics and their equations

Well, it is still a quadratic equation, but with 3 variables:

x, y, z

In general form,

$$ax^2 + by^2 + cz^2 + 2dxy + 2exz + 2fyz + 2gx + 2hy + 2mz + k = 0$$

Part 1. Types of Quadrics and their equations

Well, it is still a quadratic equation, but with 3 variables:

x, y, z

In general form,

$$ax^2 + by^2 + cz^2 + 2dxy + 2exz + 2fyz + 2gx + 2hy + 2mz + k = 0$$

In matrix/vector form the **same** equation can be presented as:

$$\begin{bmatrix} x & y & z \end{bmatrix} Q \begin{bmatrix} x \\ y \\ z \end{bmatrix} + P \begin{bmatrix} x \\ y \\ z \end{bmatrix} + R = 0$$

Q is a 3×3 matrix, P is a vector, R is a scalar.

Part 1. Types of Quadrics and their equations

Well, it is still a quadratic equation, but with 3 variables:

x, y, z

In general form,

$$ax^2 + by^2 + cz^2 + 2dxy + 2exz + 2fyz + 2gx + 2hy + 2mz + k = 0$$

In matrix/vector form the **same** equation can be presented as:

$$\begin{bmatrix} x & y & z \end{bmatrix} Q \begin{bmatrix} x \\ y \\ z \end{bmatrix} + P \begin{bmatrix} x \\ y \\ z \end{bmatrix} + R = 0$$

Q is a 3×3 matrix, P is a vector, R is a scalar.

Matrix of the **quadratic form** is Q

Demo in Geogebra

In the demo, have you recognised any types of 3D surfaces?

In the demo, have you recognised any types of 3D surfaces?
Which (make a list)?

Let us check the types:

Should be 9 types

 Ellipsoid

Let us check the types:

Should be 9 types

- Ellipsoid
- Hyperboloid of 1 sheet
- Cone
- Hyperboloid of 2 sheets

Let us check the types:

Should be 9 types

- Ellipsoid
- Hyperboloid of 1 sheet
- Cone
- Hyperboloid of 2 sheets
- Elliptic paraboloid
- Hyperbolic paraboloid

Let us check the types:

Should be 9 types

- Ellipsoid
- Hyperboloid of 1 sheet
- Cone
- Hyperboloid of 2 sheets
- Elliptic paraboloid
- Hyperbolic paraboloid
- Elliptic cylinder
- Hyperbolic cylinder
- Parabolic cylinder

All the difference is in the equation (in the canonical form).


Cylinders

Properties

In the equation, some variable is missing...

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

(for any z this will be an ellipse)



Example



Example




Cylinders...

Properties

In the equation, some variable is missing...

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

(for any z this will be a hyperbola)




Cylinders...

Properties


In the equation, some variable is missing...

$$\frac{x^2}{a^2} + 2hy = 0$$

(for any z this will be a parabola)



Other cylindrical surfaces




Ellipsoid

Properties

We have all "pluses":

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$



Example




Sphere is a special case of an ellipsoid

Properties

We have all "pluses" and $a = b = c$:

$$\frac{x^2}{a^2} + \frac{y^2}{a^2} + \frac{z^2}{a^2} = 1$$



Example





Hyperboloids

Properties

We have either one or two minuses:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = -1$$



Example




Cone as a limiting case of hyperboloids

Properties

We have zero on the right-hand side of the equation:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0$$



Example




Paraboloids

(Elliptic) Paraboloid


Properties

We have a linear term (without a quadratic one), two other are positive:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - z = 0$$



Example




Hyperbolic Paraboloid

Properties

We have a linear term (without a quadratic one), but one of the quadratic terms has a minus:


$$\frac{x^2}{a^2} - \frac{y^2}{b^2} - z = 0$$



Example



Break, 5-10 min.




<http://graphics.berkeley.edu/papers/Andrews-TCD-2013-06/Andrews-TCD-2013-06.pdf>

Part 2. Cylindrical / Spherical coordinates

Part 2. Cylindrical / Spherical coordinates




Cylindrical coordinates



$$P(r, \theta, z)$$

Cylindrical coordinates. Change to Cartesian



- $x =$
- $y =$
- $z =$

Example

Given the cylindrical coordinates, identify each of the three surfaces

$$r = 5$$

$$r^2 + z^2 = 100$$

$$z = r$$

Example


Transform point P from cylindrical coordinates to Cartesian

$$P(2, 2\pi/3, 9)$$

Transform point P from Cartesian coordinates to cylindrical


$$P(3, -3, 8)$$

Spherical coordinates



$$P(\rho, \theta, \phi)$$

Spherical coordinates. Change to Cartesian



- $X =$
- $y =$
- $Z =$

Example

Transform point P from Spherical coordinates to Cartesian

$$P(2, \pi/4, \pi/3)$$

Transform point P from Cartesian coordinates to Spherical


$$P(0, 2\sqrt{3}, -2)$$

MEME time: Cylindrical / Spherical coordinates



Cylindrical / Spherical coordinates

Cylindrical / Spherical coordinates



Part 3. Applications

Simplify of integrals

Measurements of distances (e.g. on sphere)

...

Extra. Tangent planes to quadrics

Given a function $f(x, y)$ that defines a surface, you need to define a tangent plane at a point with coordinates (x_0, y_0)

The tangent plane is given by the first-order Taylor polynomial approximation:

$$z = f(x_0, y_0) + f'_x(x - x_0) + f'_y(y - y_0)$$

You can vary the point of tangency.

Find an equation of tangent plane to the surface $z = -x^2 - y^2$ at point $(x_0, y_0) = (-1, -1)$

Useful links

- <https://www.geogebra.org>
- https://youtu.be/fNk_zzaMoSs
- <http://immersivemath.com/ila>
- <https://en.wikipedia.org/wiki/Quadric>