

Contents

1	Combinatorics	5
1.1	Binomial Coefficients	5
1.1.1	Odd numbers in the i -th line	5
1.1.2	Properties	5
1.2	Partition	7
1.3	4 fundamental problems of distribution	8
1.3.1	N equal balls in K equal boxes	8
1.3.2	N equal balls in K distinct boxes	8
1.3.3	N distinct balls in K equal boxes (Stirling's Number of Second Kind)	9
1.3.4	N distinct balls in K distinct boxes	10
2	Boolean Algebra (Logic)	10
2.1	Symbols	10
2.1.1	Implication (\rightarrow)	10
3	Geometry	10
3.1	Trigonometry	10
3.2	Line	11
3.2.1	General equation	11
3.2.2	General equation from two points	11
3.2.3	Line inclination from two points	11
3.2.4	Check if a point belongs to the line	12
3.2.5	Distance from a point to a line	12
3.3	Convex and Concave Polygons	12
3.3.1	Regular polygon circumradius	13
3.3.2	Regular polygon inscribed circle radius	13
3.3.3	Area of regular polygons	14
3.3.4	Sum of internal angle of a regular polygon	14
3.4	Triangle	14
3.4.1	Semiperimeter	14
3.4.2	Area	14
3.4.3	Circumradius	14
3.4.4	Inradius	15
3.4.5	Length of bisector	15
3.4.6	Law of sines	15
3.4.7	Law of cosines	15
3.4.8	Law of tangents	15

3.5	Quadrilaterals	15
3.6	Trapezium	16
3.6.1	Area, knowing base and height	16
3.6.2	Area knowing only sides	16
3.7	Sphere	16
3.7.1	Equation	16
3.7.2	Equation in spherical coordinate system	17
3.7.3	Area	17
3.7.4	Volume	17
3.8	Cube	17
3.8.1	Facial & Body diagonal	17
3.8.2	Area	18
3.8.3	Volume	18
3.8.4	Circumscribed sphere	18
3.8.5	Inscribed Sphere	18
3.8.6	Tangent Sphere	18
3.9	Parallelepiped	19
3.9.1	Area delimited by two vectors	19
3.9.2	Volume	19
3.10	Cylinder	19
3.10.1	Area	19
3.10.2	Volume	20
3.11	Cone	20
3.11.1	Area	20
3.11.2	Volume	20
3.12	Truncated Cone	20
3.12.1	Area	20
3.12.2	Volume	21
3.13	Dot product	21
3.14	Magnitude	21
4	Algebra	21
4.1	Absolute Value	21
4.1.1	Definition	21
4.1.2	Properties and Theorems	22
4.2	Sums	22

5	Graphs	23
5.0.1	Bipartite Graph	23
5.1	Topological Sorting	23
5.2	Strongly Connected Components	23
5.3	Minimum spanning tree	24
5.3.1	Propeties	24
5.4	Eulerian path	24
5.5	Network	25
5.6	Flow network	25
5.6.1	Propeties	25
5.7	Prufer Code	25
5.8	Prüfer's Sequence	26
6	Trees	27
6.1	Centroid	27
6.2	Centroid decomposition	27
7	Number Theory	28
7.1	Fermat's Theorems and Lemmas	28
7.2	Goldbach's Conjecture	28
7.3	Linear Diophantine Equations	28
7.3.1	Solution(s)	29
7.4	Wilson's theorem	29
7.5	Fundamental theorem of arithmetic	30
7.5.1	LCM and GCD	30
7.6	Taking modulo at the exponent	30
8	Identities	30
9	Linear Algebra	31
9.1	Matrix Multiplication	31
10	Probability Theory	31
10.1	Discrete distributions	31
10.1.1	Binomial distribution	31
10.1.2	First success distribution	32
10.1.3	Poisson distribution	32
10.2	Continuous distributions	32
10.2.1	Uniform distribution	32
10.2.2	Exponential distribution	33
10.2.3	Normal distribution	33

10.3 Markov chains	33
10.3.1 Stationary distribution	33
10.3.2 Ergodicity	34
10.3.3 Absorption	34
11 Polynomial	34
11.1 Bhaskara	34
11.2 Pascal's Triangle	34
11.3 N-th first terms of P-th column in Pascal Triangle	34
11.4 Number of odd numbers in the N-th line of pascal triangle	35
12 Trees	35
12.1 Heavy-Light Decomposition	35
13 Bitwise	35
13.1 Binary to gray code	35
13.2 Gray code to binary	35
14 Game Theory	36
14.1 Impartial Games	36
14.2 Sprague-Grundy Theorem	37
14.3 Nim variation Subtract game	37
15 Group Theory	37
15.1 Permutations	37
15.1.1 Odd permutations	38
15.1.2 Even permutations	38
16 Others	38
16.1 Unimodal Functions	38
16.2 Critérios de divisibilidade	39
16.2.17	39
16.2.211	40
16.2.313	40
16.2.417	40
16.2.519	41
16.2.623	41

1 Combinatorics

1.1 Binomial Coefficients

Binomial coefficients $\binom{n}{k}$ are the number of ways to select a set of k elements from n different elements without taking into account the order of arrangement of these elements (i.e., the number of unordered sets).

Binomial coefficients are also the coefficients in the expansion of $(a + b)^n$ (so-called binomial theorem):

$$(a + b)^n = \binom{n}{0}a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{k}a^{n-k}b^k + \dots + \binom{n}{n}b^n$$

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Recurrence formula** (which is associated with the famous "Pascal's Triangle"):

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

1.1.1 Odd numbers in the i-th line

O número de elementos ímpares na n -ésima linha do triângulo de pascal é 2^c , onde c é o número de bits na representação binária de n .

1.1.2 Properties

Binomial coefficients have many different properties. Here are the simplest of them:

- Symmetry rule:

$$\binom{n}{k} = \binom{n}{n-k}$$

- Factoring in:

$$\binom{n}{k} = \frac{n}{k} \binom{n-1}{k-1}$$

- Sum over k :

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

- Sum over n :

$$\sum_{m=0}^n \binom{m}{k} = \binom{n+1}{k+1}$$

- Sum over n and k :

$$\sum_{k=0}^m \binom{n+k}{k} = \binom{n+m+1}{m}$$

- Sum of the squares:

$$\binom{n}{0}^2 + \binom{n}{1}^2 + \dots + \binom{n}{n}^2 = \binom{2n}{n}$$

- Weighted sum:

$$1 \binom{n}{1} + 2 \binom{n}{2} + \dots + n \binom{n}{n} = n 2^{n-1}$$

- Connection with the [Fibonacci numbers](../algebra/fibonacci-numbers.md):

$$\binom{n}{0} + \binom{n-1}{1} + \dots + \binom{n-k}{k} + \dots + \binom{0}{n} = F_{n+1}$$

1.2 Partition

In number theory and combinatorics, a partition of a non-negative integer n , also called an integer partition, is a way of writing n as a sum of positive integers. Two sums that differ only in the order of their summands are considered the same partition. (If order matters, the sum becomes a composition.) For example, 4 can be partitioned in five distinct ways:

$$\begin{aligned} &4 \\ &3 + 1 \\ &2 + 2 \\ &2 + 1 + 1 \\ &1 + 1 + 1 + 1 \end{aligned}$$

The only partition of zero is the empty sum, having no parts. An individual summand in a partition is called a *part*. The number of partitions of n is given by the partition function $p(n)$. So $p(4) = 5$. The notation $\lambda \vdash n$ means that λ is a partition of n .

1	1	2	3
5	7	11	15
22	30	42	56
77	101	135	176
231	297	385	490
627	792	1002	1255
1575	1958	2436	3010
3718	4565	5604	6842
8349	10143	12310	14883
17977	21637	26015	31185
37338	44583	53174	63261
75175	89134	105558	124754
147273	173525		

Table 1: Values of $a(n)$ for n from 0 to 49

For instance, whenever the decimal representation of n ends in the digit 4 or 9, the number of partitions of n will be divisible by 5.

$P(n, k)$ denotes the number of ways of writing n as a sum of exactly k terms or, equivalently, the number of partitions into parts of which the largest is exactly k . $P(n, k)$ can be computed from the recurrence relation

$$P(n, k) = P(n - 1, k - 1) + P(n - k, k) \quad (1)$$

With $P(n, k) = 0$ for $k > n$, $P(n, n) = 1$, and $P(n, 0) = 0$. The triangle of $P(k, n)$ is given by :

N \ K	0	1	2	3	4	5	6	7	8	9	10
0	1	0	0	0	0	0	0	0	0	0	0
1	0	1	0	0	0	0	0	0	0	0	0
2	0	1	1	0	0	0	0	0	0	0	0
3	0	1	1	1	0	0	0	0	0	0	0
4	0	1	2	1	1	0	0	0	0	0	0
5	0	1	2	2	1	1	0	0	0	0	0
6	0	1	3	3	2	1	1	0	0	0	0
7	0	1	3	4	3	2	1	1	0	0	0
8	0	1	4	5	5	3	2	1	1	0	0
9	0	1	4	7	6	5	3	2	1	1	0
10	0	1	5	8	9	7	5	3	2	1	1

1.3 4 fundamental problems of distribution

1.3.1 N equal balls in K equal boxes

Considering that no box can be empty, it's given by the partition function when the number of terms is limited by k .

$$P(n, k) = P(n - 1, k - 1) + P(n - k, k) \quad (2)$$

1.3.2 N equal balls in K distinct boxes

Equivalent to count the number of solutions for the equation:

$$\begin{aligned} x_1 + \dots + x_K &= N \\ \text{where } x_i &> 0, i \in [1, K], N > 0 \end{aligned} \quad (3)$$

It's given by the formula:

$$\binom{N+1}{K-1} \quad (4)$$

If some boxes may be **empty** ($x_i \geq 0$), then it's given by:

$$\binom{N+K-1}{N}$$

1.3.3 N distinct balls in K equal boxes (Stirling's Number of Second Kind)

Also known as **Stirling's Number of Second Kind** let's define $S(N, K)$ as how many distinct ways to distribute N distinguishable balls in K indistinguishable boxes.

Some special cases:

$$\begin{aligned} S(0, 0) &= 1 \\ S(n, 0) &= 0 \\ S(n, k) &= 0 \text{ if } n < k \\ S(n, 1) &= 1 \\ S(n, k) &= 1 \text{ if } n = k \\ S(n, 2) &= 2^{n-1} - 1 \\ S(n, 3) &= \frac{1}{2}(3^{n-1} + 1) - 2^{n-1} \\ S(n, n-1) &= \binom{n}{2} \\ S(n, n-2) &= \binom{n}{3} + 3\binom{n}{4} \end{aligned} \quad (5)$$

Can be found by the following recurrence relationship.

$$S(n, k) = S(n-1, k-1) + kS(n-1, k) \quad (6)$$

There is also this formula :

$$S(n, k) = \sum_{r=0}^k (-1)^r \frac{(k-r)^n}{r!(k-r)!} \quad (7)$$

$\begin{array}{c c} & K \\ \hline N & \end{array}$	0	1	2	3	4	5	6	7	8	9
0	1	1	0	0	0	0	0	0	0	0
1	0	1	0	0	0	0	0	0	0	0
2	0	1	1	0	0	0	0	0	0	0
3	0	1	3	1	0	0	0	0	0	0
4	0	1	7	6	1	0	0	0	0	0
5	0	1	15	25	10	1	0	0	0	0
6	0	1	31	90	65	15	1	0	0	0
7	0	1	63	301	350	140	21	1	0	0
8	0	1	127	966	1701	1050	266	28	1	0
9	0	1	255	3025	7770	6951	2646	462	36	1

1.3.4 N distinct balls in K distinct boxes

$$K^N \quad (8)$$

2 Boolean Algebra (Logic)

2.1 Symbols

2.1.1 Implication (\rightarrow)

$$a \rightarrow b \Leftrightarrow \neg a \vee b \quad (9)$$

3 Geometry

3.1 Trigonometry

$$\sin(v + w) = \sin v \cos w + \cos v \sin w \quad (10)$$

$$\cos(v + w) = \cos v \cos w - \sin v \sin w \quad (11)$$

$$\tan(v + w) = \frac{\tan v + \tan w}{1 - \tan v \tan w} \quad (12)$$

$$\sin v + \sin w = 2 \sin \frac{v+w}{2} \cos \frac{v-w}{2} \quad (13)$$

$$\cos v + \cos w = 2 \cos \frac{v+w}{2} \cos \frac{v-w}{2} \quad (14)$$

$$(V+W) \tan(v-w)/2 = (V-W) \tan(v+w)/2$$

where V, W are lengths of sides opposite angles v, w . (15)

$$\begin{aligned} a \cos x + b \sin x &= r \cos(x - \phi) \\ a \sin x + b \cos x &= r \sin(x + \phi) \end{aligned} \quad (16)$$

where $r = \sqrt{a^2 + b^2}, \phi = \text{atan2}(b, a)$.

3.2 Line

3.2.1 General equation

$$ax + by + c = 0 \quad (17)$$

Note that a same line can have multiple representations (just multiply the equation by any real number), so to make each line have a single equation divide everything by a , or b if a is zero.

3.2.2 General equation from two points

Let P and Q be the points that define the line.

$$\begin{aligned} a &= P_y - Q_y \\ b &= Q_x - P_x \\ c &= P_x Q_y - P_y Q_x \end{aligned}$$

3.2.3 Line inclination from two points

Let P and Q be two points that belongs to the line, such that $P_x < Q_x$ the inclination m or angular coefficient is given by:

$$m = \frac{Q_y - P_y}{Q_x - P_x} \quad (18)$$

3.2.4 Check if a point belongs to the line

Let r be a line such that $ax + by + c = 0$ and P a point. $P \in r$ if and only if :

$$aP_x + bP_y + c = 0$$

3.2.5 Distance from a point to a line

The distance from a point P and a line r is defined as the shortest distance possible between every point that belongs to r and P . Such distance will be the distance from P and the intersection between r and the orthogonal projection from P to r , and can be found by:

$$\frac{|aP_x + bP_y + c|}{\sqrt{a^2 + b^2}}$$

The coordinates of the point Q are given by:

$$Q_x = \frac{b(bP_x - aP_y) - ac}{a^2 + b^2}$$

$$Q_y = \frac{a(-bP_x + aP_y) - bc}{a^2 + b^2}$$

3.3 Convex and Concave Polygons






Figure 1: Convex Polygon Figure 2: Concave Polygon

Figure 3: Two Types of Polygons

3.3.1 Regular polygon circumradius

$$R = \frac{s}{2} \csc \frac{\pi}{n}$$

3.3.2 Regular polygon inscribed circle radius

$$r = R \cos \frac{\pi}{n}$$

3.3.3 Area of regular polygons

- Let n be the number of sides of the regular polygon, the area can be found using one of the values below:
 - the length of one of the sides (s)
 - apothem, the radius of the inscribed circle (r)
 - the radius of the circumscribed circle (R)

$$A = \frac{1}{2}nrs = \frac{1}{4}ns^2 \cot \frac{\pi}{n} = nr^2 \tan \frac{\pi}{n} = \frac{1}{2}nR^2 \sin \frac{2\pi}{n}$$

3.3.4 Sum of internal angle of a regular polygon

A regular polygon with n sides have $(n - 2)180$ degrees as sum of it's internal angle.

3.4 Triangle

Let the length of the sides of the triangle be a, b, c .

3.4.1 Semiperimeter

Let p be the semiperimeter defined as:

$$p = \frac{a + b + c}{2} \quad (19)$$

3.4.2 Area

Let A be the area defined as:

$$\sqrt{p(p - a)(p - b)(p - c)} \quad (20)$$

3.4.3 Circumradius

$$R = \frac{abc}{4A} \quad (21)$$

3.4.4 Inradius

$$r = \frac{A}{p} \quad (22)$$

3.4.5 Length of bisector

$$s_a = \sqrt{bc \left[1 - \left(\frac{a}{b+c} \right)^2 \right]} \quad (23)$$

3.4.6 Law of sines

$$\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c} = \frac{1}{2R} \quad (24)$$

3.4.7 Law of cosines

$$a^2 = b^2 + c^2 - 2bc \cos \alpha \quad (25)$$

3.4.8 Law of tangents


$$\frac{a+b}{a-b} = \frac{\tan \frac{\alpha+\beta}{2}}{\tan \frac{\alpha-\beta}{2}} \quad (26)$$

3.5 Quadrilaterals

Let its sides length be a, b, c, d


3.6 Trapezium

3.6.1 Area, knowing base and height



$$A = \frac{(B + b)h}{2}$$

3.6.2 Area knowing only sides



$$e = \frac{d^2 - b^2 + a^2 - 2ac + c^2}{2a - 2c}$$

$$h = \sqrt{d^2 - e^2}$$

$$A = \frac{h(a + c)}{2}$$

3.7 Sphere

3.7.1 Equation

$$(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2$$

3.7.2 Equation in spherical coordinate system

r is the radius, θ is an angle that goes from 0 to 2π , and φ is an angle that goes from 0 to π .

$$\begin{aligned}x &= x_0 + r \cos \theta \sin \varphi \\y &= y_0 + r \sin \theta \sin \varphi \\z &= z_0 + r \cos \varphi\end{aligned}$$

3.7.3 Area

$A = 4\pi r^2$, where r is the radius, comes from :

$$A = \int_0^\pi \int_0^{2\pi} r^2 \sin(\varphi) d\theta d\varphi$$

3.7.4 Volume

$$V = \frac{4}{3}\pi r^3$$


comes from :

$$V = \int_0^R \int_0^\pi \int_0^{2\pi} r^2 \sin(\varphi) d\theta d\varphi dR$$

3.8 Cube**3.8.1 Facial & Body diagonal**

Facial diagonal join two vertices at the same face.

Body diagonal join two vertices from opposite faces.



$$\text{Facial diagonal} = L\sqrt{2}$$

$$\text{Body diagonal} = L\sqrt{3}$$

3.8.2 Area

$$A = 6L^2$$

3.8.3 Volume

$$V = L^3$$

3.8.4 Circumscribed sphere

Pass through the 8 vertices, radius equal to $L(\frac{\sqrt{3}}{2})$.

3.8.5 Inscribed Sphere

Tangent to the 6 faces, radius equal to $\frac{L}{2}$.

3.8.6 Tangent Sphere

Tangent to the edges, radius equal to $\frac{L}{\sqrt{2}}$.

3.9 Parallelepiped

Let it be defined by three linear independent vectors

$$\vec{u}, \vec{v}, \vec{w}$$

3.9.1 Area delimited by two vectors

$$A = |\vec{u} \times \vec{v}|$$

3.9.2 Volume


$$V = |(\vec{u} \times \vec{v}) \cdot \vec{w}|,$$

$$(\vec{u} \times \vec{v}) \cdot \vec{w} = \det \begin{bmatrix} u_x & u_y & u_z \\ v_x & v_y & v_z \\ w_x & w_y & w_z \end{bmatrix}$$

or if you know the sides a, b, c , and the angles α, β, γ :

$$V = abc \sqrt{1 + 2 \cos \alpha \cos \beta \cos \gamma - \cos^2 \alpha - \cos^2 \beta - \cos^2 \gamma}$$

3.10 Cilinder




3.10.1 Area

$$A = 2\pi r h + 2\pi r^2 = 2\pi r(h + r),$$

3.10.2 Volume


$$A = \pi r^2 h$$

3.11 Cone**3.11.1 Area**

$$A = \pi r^2 + \pi r \sqrt{r^2 + H^2}$$

3.11.2 Volume

$$V = \frac{1}{3} \pi r^2 H$$

3.12 Truncated Cone**3.12.1 Area**

$$l = \sqrt{h^2 + R^2}$$

$$A = \pi(R + r)l + \pi(R^2 + r^2)$$

3.12.2 Volume

$$V = \frac{1}{3}\pi h(R^2 + Rr + r^2),$$

3.13 Dot product

The dot product of vectors \mathbf{u} and \mathbf{v} in n dimensions is given by:

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 \cdot v_1 + u_2 \cdot v_2 + \dots + u_n \cdot v_n$$

3.14 Magnitude

The magnitude of a vector \mathbf{v} in n dimensions is given by:

$$|\mathbf{v}| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$$

The dot product of two Euclidean vectors \mathbf{u} and \mathbf{v} is defined by

$$\langle \mathbf{u}, \mathbf{v} \rangle = |\mathbf{u}| \cdot |\mathbf{v}| \cdot \cos(\theta)$$

4 Algebra

4.1 Absolute Value

4.1.1 Definition

The absolute value of x , denoted by $|x|$ is defined by:

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases} \quad (27)$$

4.1.2 Properties and Theorems

$$|x| < a \iff -a < x < a, \text{ where } a > 0 \quad (28)$$

$$|x| \leq a \iff -a \leq x \leq a, \text{ where } a > 0 \quad (29)$$

$$|x| > a \iff (x > a) \vee (x < -a), \text{ where } a > 0 \quad (30)$$

$$|x| \geq a \iff (x \geq a) \vee (x \leq -a), \text{ where } a > 0 \quad (31)$$

$$|ab| = |a| \cdot |b|, \text{ where } a, b \in \mathbb{R} \quad (32)$$

$$\left| \frac{a}{b} \right| = \frac{|a|}{|b|}, \text{ where } a, b \in \mathbb{R}, \text{ and } b \neq 0 \quad (33)$$

$$|a + b| \leq |a| + |b|, \text{ where } a, b \in \mathbb{R} \quad (34)$$

$$|a - b| \leq |a| + |b|, \text{ where } a, b \in \mathbb{R} \quad (35)$$

$$|a| - |b| \leq |a - b|, \text{ where } a, b \in \mathbb{R} \quad (36)$$

4.2 Sums

$$\begin{aligned} c^a + c^{a+1} + \dots + c^b &= \frac{c^{b+1} - c^a}{c - 1}, c \neq 1 \\ 1 + 2 + 3 + \dots + n &= \frac{n(n+1)}{2} \\ 1^2 + 2^2 + 3^2 + \dots + n^2 &= \frac{n(2n+1)(n+1)}{6} \\ 1^3 + 2^3 + 3^3 + \dots + n^3 &= \frac{n^2(n+1)^2}{4} \\ 1^4 + 2^4 + 3^4 + \dots + n^4 &= \frac{n(n+1)(2n+1)(3n^2+3n-1)}{30} \end{aligned} \quad (37)$$

5 Graphs

5.0.1 Bipartite Graph

Bipartite Graph is a special graph with the following characteristics: the set of vertices V can be partitioned into two disjoint sets V_1 and V_2 and all undirected edges $(u, v) \in E$ have the property that $u \in V_1$ and $v \in V_2$. This makes a Bipartite Graph free from odd-length cycle. Not that by this definition is possible to have isolated vertices.

5.1 Topological Sorting

Definition: You are given a **directed** graph with n vertices and m edges. You have to find an order of the vertices, so that every edge leads from the vertex with a smaller index to a vertex with a larger one.

Topological order can be non-unique !

A Topological order **may not exist** at all. It only exists, if the directed graph **contains no cycles**. Otherwise because there is a contradiction: if there is a cycle containing the vertices a and b , then a needs to have a smaller index than b (since you can reach b from a) and also a bigger one (as you can reach a from b). Every acyclic directed graph contains at least one topological order.

5.2 Strongly Connected Components

You are given a directed graph G with vertices V and edges E . It is possible that there are loops and multiple edges. Let's denote n as number of vertices and m as number of edges in G .

Strongly connected component is a maximal subset of vertices C such that any two vertices of this subset are reachable from each other, i.e. for any $u, v \in C$:

$$u \mapsto v, v \mapsto u$$

where \mapsto means reachability, i.e. existence of the path from first vertex to the second.

The most important property of the condensation graph is that it is a **DAG**. Indeed, suppose that there is an edge between C and C' , let's prove that there is no edge from C' to C . Suppose that $C' \mapsto C$. Then there are two vertices $u' \in C$ and $v' \in C'$ such that $v' \mapsto u'$. But since u and u' are in the same strongly connected component then there is a path between them; the same for v and v' . As a result, if we join these paths we have that $v \mapsto u$ and at the same time $u \mapsto v$. Therefore u and v should be at the same strongly connected component, so this is contradiction. This completes the proof.

5.3 Minimum spanning tree

5.3.1 Properties

- A minimum spanning tree of a graph is unique, if the weight of all the edges are distinct. Otherwise, there may be multiple minimum spanning trees. (Specific algorithms typically output one of the possible minimum spanning trees).
- Minimum spanning tree is also the tree with minimum product of weights of edges. (It can be easily proved by replacing the weights of all edges with their logarithms)

5.4 Eulerian path

A Eulerian path is a path in a graph that passes through all of its edges exactly once. A Eulerian cycle is a Eulerian path that is a cycle.

An Eulerian cycle exists if and only if the degrees of all vertices are even

And an Eulerian path exists if and only if the number of vertices with odd degrees is two (or zero, in the case of the existence of a Eulerian cycle).

5.5 Network

A **network** is a directed graph G with vertex V and edges E combined with a function c , which assigns each edge $e \in E$ a non-negative integer value, the *capacity* of e .

5.6 Flow network

Is a **network** with two vertices labeled as **source** and **sink**.

5.6.1 Properties

- The flow of an edge cannot exceed the capacity

$$f(e) \leq c(e)$$

- And the sum of the incoming flow of a vertex u has to be equal to the sum of the outgoing flow of u except in the source and sink vertices.
- The source vertex s only has an outgoing flow, and the sink vertex t has only incoming flow.

5.7 Prüfer Code

The Prüfer code is a way of encoding a labeled tree with n vertices using a sequence of $n - 2$ integers in the interval $[0; n - 1]$. This encoding also acts as a bijection between all spanning trees of a complete graph and the numerical sequences.

The Prüfer code is constructed as follows. We will repeat the following procedure $n - 2$ times: we select the leaf of the tree with the smallest number, remove it from the tree, and write down the number of the vertex that was connected to it. After $n - 2$ iterations there will only remain 2 vertices, and the algorithm ends.

Thus the Prüfer code for a given tree is a sequence of $n - 2$ numbers, where each number is the number of the connected vertex, i.e. this number is in the interval $[0, n - 1]$.

The algorithm for computing the Prüfer code can be implemented easily with $O(n \log n)$ time complexity, simply by using a data structure to extract the minimum (for instance `set` or `priority_queue` in C++), which contains a list of all the current leaves.

After constructing the Prüfer code two vertices will remain. One of them is the highest vertex $n - 1$, but nothing else can be said about the other one.

Each vertex appears in the Prüfer code exactly a fixed number of times - its degree minus one. This can be easily checked, since the degree will get smaller every time we record its label in the code, and we remove it once the degree is 1. For the two remaining vertices this fact is also true.

5.8 Prüfer's Sequence

The Prüfer sequence is a bijection between labeled trees with n vertices and sequences with $n - 2$ numbers from 1 to n . To get the sequence from the tree:

- While there are more than 2 vertices, remove the leaf with smallest label and append it's neighbour to the end of the sequence.

To get the tree from the sequence:

- The degree of each vertex is 1 more than the number of occurrences of that vertex in the sequence. Compute the degree d , then do the following: for every value x in the sequence (in order), find the vertex with smallest label y such that $d(y) = 1$ and add an edge between x and y , and also decrease their degrees by 1. At the end of this procedure, there will be two vertices left with degree 1; add an edge between them.

6 Trees

6.1 Centroid

A centroid of a tree is defined as a node such that when the tree is rooted at it, no other nodes have a subtree of size greater than $\frac{N}{2}$.

- Every tree have a centroid.
- There is at most two centroids (mostly only one)
- All centroids are adjacent
- Let $s(v)$ be the sum of the distance from v to every other node, $\sum_{u \neq v} \text{dist}(u, v)$, then if, if c is the centroid, it implies that $s(c)$ is the smallest one among every other node

6.2 Centroid decomposition

The centroid decomposition of a tree is another tree defined recursively as:

- Its root is the centroid of the original tree.
- Its children are the centroid of each tree resulting from the removal of the centroid from the original tree.

Properties:

- The tree formed contains all the N nodes.
- The height of the centroid decomposition is $O(\log N)$.
- Let A and B be two arbitrary nodes, let C be the $LCA(A, B)$ in the centroid decomposition tree.
 - C lies on the path from A to B in the original tree.
 - Every path in the tree can be decomposed in A to C , and C to B .
 - As the height of the tree is $O(\log N)$ the LCA can be found in $O(\log N)$ (just like with binary lifting)

7 Number Theory

7.1 Fermat's Theorems and Lemmas

Let p be a prime number and $a, b \in \mathbb{Z}$:

$$a^p \equiv a \pmod{p} \quad (38)$$

$$a^{p-1} \equiv 1 \pmod{p} \quad (39)$$

$$(a + b)^p \equiv a^p + b^p \pmod{p} \quad (40)$$

$$a^{-1} \equiv a^{p-2} \pmod{p} \quad (41)$$

7.2 Goldbach's Conjecture

"Every pair number greater than 2 can be written as the sum of two primes"

Valid for every integer in range from 4 to 10^{18} , but without proof

For an odd x number it can be written as the sum of two primes if $x - 2$ is also prime, or three primes, 3 and the two primes that results in $x - 3$.

7.3 Linear Diophantine Equations

A Linear Diophantine Equation (in two variables) is an equation of the general form:

$$ax + by = c$$

Where a, b, c are given integers, and x, y are unknown integers.

If $a = b = 0$, we have infinite solutions if $c = 0$, and 0 otherwise.

7.3.1 Solution(s)

Let $g = \gcd(a, b)$ such that $ax_g + by_g = g$, then we only have a solution if and only if $g \mid c$, and if it have a solution it have infinite.

The solutions will be of the form :

$$\begin{aligned} x_0 &= x_g \cdot \frac{c}{g}, y_0 = y_g \cdot \frac{c}{g}. \\ a \cdot x_0 + b \cdot y_0 &= c \end{aligned} \quad (42)$$

With the initial solution, we can can find every solution, with :

$$x = x_0 + k \cdot \frac{b}{g}, y = y_0 - k \cdot \frac{a}{g} \quad (43)$$

To find the solution that minimize $x + y$ we use the fact that:

$$\begin{aligned} x' &= x + k \cdot \frac{b}{g}, \\ y' &= y - k \cdot \frac{a}{g}. \end{aligned}$$

Note that

$x + y$ change as follows:

$$x' + y' = x + y + k \cdot \left(\frac{b}{g} - \frac{a}{g} \right) = x + y + k \cdot \frac{b - a}{g}$$

If $a < b$, we need to select smallest possible value of k . If $a > b$, we need to select the largest possible value of k . If $a = b$, all solution will have the same sum $x + y$.

7.4 Wilson's theorem

Wilson's theorem states that a natural number $n > 1$ is a **prime number** if and only if the product of all the positive integers less than n is one less than a multiple of n .

That is :

$$(n - 1)! \equiv -1 \pmod{n} \quad (44)$$

In other words, any integer $n > 1$ is a prime number if, and only if, $(n - 1)! + 1$ is divisible by n

7.5 Fundamental theorem of arithmetic

Every integer greater than 1 can be represented uniquely as a product of prime numbers, up to the order of the factors.

$$n = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_k^{\alpha_k} \quad (45)$$

7.5.1 LCM and GCD

$$\begin{aligned} a &= p_1^{\alpha_1} p_2^{\alpha_2} \dots p_k^{\alpha_k} \\ b &= p_1^{\beta_1} p_2^{\beta_2} \dots p_k^{\beta_k} \\ (a, b) &= p_1^{\min \alpha_1, \beta_1} p_2^{\min \alpha_2, \beta_2} \dots p_k^{\min \alpha_k, \beta_k} \\ [a, b] &= p_1^{\max \alpha_1, \beta_1} p_2^{\max \alpha_2, \beta_2} \dots p_k^{\max \alpha_k, \beta_k} \end{aligned} \quad (46)$$

7.6 Taking modulo at the exponent

If $\gcd(a, m) = 1$ then:

$$a^m \equiv a^{n \bmod \varphi(m)} \pmod{m} \quad (47)$$

8 Identities

$$\begin{aligned} \sum_{i=1}^n i &= \frac{n(n+1)}{2} \\ \sum_{i=1}^n i^2 &= \frac{n(n+1)(2n+1)}{6} \\ \sum_{i=1}^n i^3 &= \frac{n^2(n+1)^2}{4} = \left(\sum_{i=1}^n i \right)^2 \end{aligned}$$

9 Linear Algebra

9.1 Matrix Multiplication

Let $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$. The product $C = AB \in \mathbb{R}^{m \times p}$ is defined as:

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

for $i = 1, \dots, m$ and $j = 1, \dots, p$.

Matrix multiplication is associative and distributive, but not commutative: $AB \neq BA$ in general.

10 Probability Theory

Let X be a **discrete random** variable with probability $p_X(x)$ of assuming the value x .

It will then have an **expected value** (mean)

$$\mu = \mathbb{E}(X) = \sum_x x p_X(x)$$

Variance $\sigma^2 = V(X) = \mathbb{E}(X^2) - (\mathbb{E}(X))^2 = \sum_x (x - \mathbb{E}(X))^2 p_X(x)$ where σ is the standard deviation.

If X is instead continuous it will have a probability density function $f_X(x)$ and the sums above will instead be integrals with $p_X(x)$ replaced by $f_X(x)$.

Expectation is linear:

$$\mathbb{E}(aX + bY) = a\mathbb{E}(X) + b\mathbb{E}(Y)$$

For independent X and Y ,

$$V(aX + bY) = a^2 V(X) + b^2 V(Y).$$

10.1 Discrete distributions

10.1.1 Binomial distribution

The number of successes in n independent yes/no experiments, each which yields success with probability p is

$\text{Bin}(n, p)$, $n = 1, 2, \dots$, $0 \leq p \leq 1$.

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$\mu = np, \sigma^2 = np(1-p)$$

$\text{Bin}(n, p)$ is approximately $\text{Po}(np)$ for small p .

10.1.2 First success distribution

The number of trials needed to get the first success in independent yes/no experiments, each which yields success with probability p is $\text{Fs}(p)$, $0 \leq p \leq 1$.

$$p(k) = p(1-p)^{k-1}, k = 1, 2, \dots$$

$$\mu = \frac{1}{p}, \sigma^2 = \frac{1-p}{p^2}$$

10.1.3 Poisson distribution

The number of events occurring in a fixed period of time t if these events occur with a known average rate κ and independently of the time since the last event is $\text{Po}(\lambda)$, $\lambda = t\kappa$.

$$p(k) = e^{-\lambda} \frac{\lambda^k}{k!}, k = 0, 1, 2, \dots$$

$$\mu = \lambda, \sigma^2 = \lambda$$

10.2 Continuous distributions

10.2.1 Uniform distribution

If the probability density function is constant between a and b and 0 elsewhere it is $\text{U}(a, b)$, $a < b$.

$$f(x) = \begin{cases} \frac{1}{b-a} & a < x < b \\ 0 & \text{otherwise} \end{cases}$$

$$\mu = \frac{a+b}{2}, \sigma^2 = \frac{(b-a)^2}{12}$$

10.2.2 Exponential distribution

The time between events in a Poisson process is $\text{Exp}(\lambda)$, $\lambda > 0$.

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & x < 0 \end{cases}$$

$$\mu = \frac{1}{\lambda}, \sigma^2 = \frac{1}{\lambda^2}$$

10.2.3 Normal distribution

Most real random values with mean μ and variance σ^2 are well described by $N(\mu, \sigma^2)$, $\sigma > 0$.

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

If $X_1 \sim N(\mu_1, \sigma_1^2)$ and $X_2 \sim N(\mu_2, \sigma_2^2)$ then

$$aX_1 + bX_2 + c \sim N(\mu_1 + \mu_2 + c, a^2\sigma_1^2 + b^2\sigma_2^2)$$

10.3 Markov chains

A *Markov chain* is a discrete random process with the property that the next state depends only on the current state. Let X_1, X_2, \dots be a sequence of random variables generated by the Markov process. Then there is a transition matrix $\mathbf{P} = (p_{ij})$, with $p_{ij} = \Pr(X_n = i | X_{n-1} = j)$, and $\mathbf{p}^{(n)} = \mathbf{P}^n \mathbf{p}^{(0)}$ is the probability distribution for X_n (i.e., $p_i^{(n)} = \Pr(X_n = i)$), where $\mathbf{p}^{(0)}$ is the initial distribution.

10.3.1 Stationary distribution

is a stationary distribution if $\pi = \mathbf{P} \pi$. If the Markov chain is *irreducible* (it is possible to get to any state from any state), then $\pi_i = \frac{1}{\mathbb{E}(T_i)}$ where $\mathbb{E}(T_i)$ is the expected time between two

visits in state i . π_j/π_i is the expected number of visits in state j between two visits in state i .

For a connected, undirected and non-bipartite graph, where the transition probability is uniform among all neighbors, π_i is proportional to node i 's degree.

10.3.2 Ergodicity

A Markov chain is *ergodic* if the asymptotic distribution is independent of the initial distribution. A finite Markov chain is ergodic iff it is irreducible and *aperiodic* (i.e., the gcd of cycle lengths is 1). $\lim_{k \rightarrow \infty} \mathbf{P}^k = \mathbf{1}\pi$.

10.3.3 Absorption

A Markov chain is an A-chain if the states can be partitioned into two sets **A** and **G**, such that all states in **A** are absorbing ($p_{ii} = 1$), and all states in **G** leads to an absorbing state in **A**. The probability for absorption in state $i \in \mathbf{A}$, when the initial state is j , is $a_{ij} = p_{ij} + \sum_{k \in \mathbf{G}} a_{ik} p_{kj}$. The expected time until absorption, when the initial state is i , is $t_i = 1 + \sum_{k \in \mathbf{G}} p_{ki} t_k$.

11 Polynomial

11.1 Bhaskara

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad (48)$$

11.2 Pascal's Triangle

11.3 N-th first terms of P-th column in Pascal Triangle

$$\binom{p}{p} + \binom{p+1}{p} + \dots + \binom{p+n}{p} = \binom{p+n+1}{p+1} \quad (49)$$

11.4 Number of odd numbers in the N-th line of pascal triangle

There is a mathematical relation which gives the number of odd numbers in the N-th row of pascal's triangle. The theorem states that the number of odd numbers in N-th row is equal to 2 raised to the number of ones in the binary representation of N.

12 Trees

12.1 Heavy-Light Decomposition

Heavy-Light Decomposition (HLD) is a technique to decompose a tree into a set of disjoint paths. This technique is particularly useful to deal with problems which require us to do some path-queries in a tree which seemingly complicated but easy enough to be solved for a line-graph. The idea is to decompose the tree into several paths (line-graph) of disjoint vertices. Then, each path-query in the original tree might be able to be answered by queries in one or more of those paths.

An edge (a, b) is heavy if and only if $\text{size}(b) \geq \text{size}(a)/2$; otherwise, it is light

13 Bitwise

13.1 Binary to gray code

$$\begin{aligned} G_n &= B_n \\ G_{n-1} &= B_n \oplus B_{n-1} \\ G_i &= B_i \oplus B_{i-1} \\ &\dots \\ G_1 &= B_2 \oplus B_1 \end{aligned}$$

13.2 Gray code to binary

$$B_n = G_n$$

$$B_{n-1} = B_n \oplus G_{n-1} = G_n \oplus G_{n-1}$$

...

$$B_1 = B_2 \oplus G_1 = G_n \oplus G_1$$

14 Game Theory

14.1 Impartial Games

To be considered a impartial game following rules must be true:

1. The available moves win/lose depends only on the state of the game, in other words, the only difference between the two players is that one of them moves first
2. Additionally, we assume that the game has perfect information, i.e. no information is hidden from the players (they know the rules and the possible moves).
3. It is assumed that the game is finite, i.e. after a certain number of moves, one of the players will end up in a losing position — from which they can't move to another position. On the other side, the player who set up this position for the opponent wins. Understandably, there are no draws in this game.

Such games can be completely described by a directed acyclic graph: the vertices are game states and the edges are transitions (moves). A vertex without outgoing edges is a losing vertex (a player who must make a move from this vertex loses).

Since there are no draws, we can classify all game states as either winning or losing. Winning states are those from which there is a move that causes inevitable defeat of the other player, even with their best response. Losing states are those from which all moves lead to winning states for the other player. Summarizing, a state is winning if there is at least one transition to a losing state and is losing if there isn't at least one transition to a losing state.

Our task is to classify the states of a given game.

14.2 Sprague-Grundy Theorem

The Sprague-Grundy Theorem states that every impartial game is equivalent to a pile of a certain size in Nim. In other words, every impartial game can be solved as Nim by finding their corresponding game.

Basically, for a game situation A and its SG function value $g(A)$:

1. $g(A) = 0$ if and only if A is a must-lose situation.
Otherwise, $g(A) \in \mathbb{Z}^*$
2. If A can be divided into n sub-situations x_1, x_2, \dots, x_n ,
then $g(A) = g(x_1) \oplus g(x_2) \oplus \dots \oplus g(x_n)$
3. If A can be converted to situation B_1 or B_2 or ... or B_n by only one operation, then
 $g(A) = \text{mex}(g(B_1), g(B_2), \dots, g(B_n))$ where function $\text{mex}(S)$ is defined as the smallest non-negative integer that does not appear in S . For example,
 $\text{mex}(0, 1, 2, 4) = 3, \text{mex} = 0, \text{mex}(0, 1, 2, 4) = 3, \text{mex}() = 0$

14.3 Nim variation Subtract game

Work just like nim but instead remove any number of objects you can remove at most K , this game can be seen as a nim game but before computing the num-sum you need to take the size of each pile module $K + 1$, the optimal way to play it is by taking K at each turn.

15 Group Theory

15.1 Permutations

A permutation is an arrangement of elements. A permutation of N elements can be represented by an arrangement of the numbers $1, 2, \dots, N$ in some order. Eg. $5, 1, 4, 2, 3$.

15.1.1 Odd permutations

A permutation is called odd if it can be expressed as a product of odd number of transpositions.

A sorted permutation is an even permutation

15.1.2 Even permutations


A permutation is called even if it can be expressed as a product of even number of transpositions.

16 Others

16.1 Unimodal Functions

By unimodal function, we mean one of two behaviors of the function:

- The function strictly increases first, reaches a maximum (at a single point or over an interval), and then strictly decreases.
- The function strictly decreases first, reaches a minimum, and then strictly increases.



16.2 Critérios de divisibilidade

16.2.1 7

Para verificar a divisibilidade de um número por 7, siga a seguinte regra:

1. Pegue o número em questão.
2. Remova o último dígito (unidade) do número.
3. Dobre o valor removido no passo anterior.
4. Subtraia o valor dobrado do número restante.
5. Se o resultado da subtração for divisível por 7, o número original é divisível por 7.

Exemplo:

Suponha que desejamos verificar a divisibilidade do número 413 por 7.

1. Remova o último dígito (3) e dobre-o, obtendo 6.
2. Subtraia 6 do número restante ($41 - 6 = 35$).

16.2.2 11

n é divisível por 11 $\Leftrightarrow \sum_{i=1}^k a_{2i-1} - \sum_{i=1}^j a_{2i}$ é divisível por 11

onde a_i é o i -ésimo dígito do número n , k é a quantidade de dígitos ímpares, j é a quantidade de dígitos pares.

Exemplo:

Suponha que desejamos verificar a divisibilidade do número $n = 7923$ por 11.

$$k = 2, \quad j = 2$$

Soma dos dígitos ímpares: $7 + 3 = 10$

Soma dos dígitos pares: $9 + 2 = 11$

Subtração: $10 - 11 = -1$

Como -1 não é divisível por 11, o número 7923 não é divisível por 11.

16.2.3 13

$$13|x \equiv 13|4 \cdot (x\%10) + [x/10] \quad (50)$$

Em outras palavras 13 divide x se o quádruplo do último algarismo somado com o número sem este algarismo for divisível por 13.

16.2.4 17

$$17|x \equiv 17|[x/10] - 5 \cdot (x\%10) \quad (51)$$

Em outras palavras 17 divide x se o a diferença entre o quádruplo do último algarismo e o número sem este algarismo for divisível por 17.

16.2.5 19

$$19|x \equiv 19|[x/10] + 2 \cdot (x \bmod 10) \quad (52)$$

Em outras palavras 19 divide x se o dobro do último algarismo de x somado a o número restante de x é divisível por 19.

16.2.6 23

$$23|x \equiv 23|x/10 + 7 \cdot (x \bmod 10) \quad (53)$$