

A Digit-Structured Interval Decomposition for Lucas HIT Sets

Start Lattices, First-Strict Digits, Interval Lengths, and Exact Counting

IamLupo

13 September 2026

Abstract

We study the set of integers detected by a Lucas-theoretic divisibility predicate of the form

$$p \mid \binom{t}{m-t},$$

or, equivalently in the parameterization used throughout the experiments, the set of integers for which a binomial coefficient has positive p -adic valuation. The starting point is Lucas' theorem: divisibility is determined entirely by a comparison of base- p digits. From this digit predicate we experimentally discovered, and in the relevant elementary steps can derive exactly, a rigid interval structure.

For a family of parameters admitting the representation

$$m = s_0 + qp^e - 1,$$

the HIT set begins at lattice points

$$x_j = s_0 + jp^e, \quad j < q, \quad j \preceq_p q,$$

where $j \preceq_p q$ denotes digitwise domination in base p . If

$$r(j) = \min\{i : j_i < q_i\},$$

then the corresponding HIT interval has length

$$L_r = (p^r - (q \bmod p^r))p^e - s_0,$$

and terminates at

$$E_j = (\lfloor j/p^r \rfloor + 1)p^{e+r} - 1.$$

Thus all intervals having the same first-strict digit position r have exactly the same length, while the number of such intervals is

$$C_r = q_r \prod_{i>r} (q_i + 1).$$

The resulting weighted interval decomposition counts the complete HIT set exactly. Independently, the MISS set is counted directly by Lucas' digitwise condition, giving

$$\#H(1, \dots, m) = m + 1 - \prod_i (m_i + 1).$$

Exhaustive computational experiments over 24,896 parameter instances and more than 1.8 million individual intervals confirm all stated identities in the tested range. The paper separates statements proved directly from Lucas' theorem and elementary digit arguments from identities established by exhaustive and randomized computation. The resulting formulas give a complete experimentally validated interval decomposition and an exact algebraic collapse of its total cardinality.

1 Introduction

A recurring theme in the present line of research is that an apparently arithmetic condition becomes highly structured after passing to base- p digits. Lucas' theorem is a particularly clean example: divisibility of a binomial coefficient modulo a prime p is determined digitwise rather than by the full magnitude of the integers involved.

The purpose of this paper is to record a further structure discovered experimentally from that digitwise predicate. Instead of treating individual HIT and MISS values independently, we organize the HIT set into maximal consecutive intervals. The experiments show that those intervals are not arbitrary. Their starting points lie on a fixed arithmetic lattice, the lattice indices are precisely the digitwise subnumbers of a single integer q , and each interval is classified by the first base- p digit at which that lattice index is strictly smaller than q .

The resulting picture has several layers:

- (i) a fixed start lattice with spacing p^e ;
- (ii) a digitwise characterization of admissible start indices;
- (iii) a first-strict digit position r attached to each interval;
- (iv) an exact interval length determined only by r ;
- (v) an exact endpoint lying one unit below a multiple of p^{e+r} ;
- (vi) an exact count of intervals of each type r ; and
- (vii) an exact total HIT count, independently recoverable from the digits of m .

The central point is that a global-looking set of divisibility events collapses to a finite digit grammar.

2 Lucas' digit predicate

Let p be prime. Write nonnegative integers in base p as

$$x = \sum_{i \geq 0} x_i p^i, \quad y = \sum_{i \geq 0} y_i p^i, \quad 0 \leq x_i, y_i < p.$$

Definition 2.1 (Digitwise domination). We write

$$x \preceq_p y$$

if and only if

$$x_i \leq y_i \quad \text{for every } i.$$

Lucas' theorem gives

$$\binom{m}{t} \equiv \prod_i \binom{m_i}{t_i} \pmod{p}.$$

Consequently,

$$p \nmid \binom{m}{t} \iff t \preceq_p m,$$

and therefore

$$p \mid \binom{m}{t} \iff t \not\preceq_p m.$$

We call the latter values *HITs* and the former values *MISSes*:

$$H_p(m) = \{t : t \not\leq_p m\}, \quad M_p(m) = \{t : t \leq_p m\}.$$

The experiments in this paper use the natural HIT variable t together with the Lucas parameter m . No analytic approximation is involved: the underlying predicate is exactly digitwise.

3 A structured parameterization

The observed interval geometry occurs when m admits a decomposition

$$m = s_0 + qp^e - 1. \tag{1}$$

The experiments derive this representation from the parameter construction used in the earlier arithmetic-sequence setup. For the present paper, (1) is taken as the structural input.

The fixed quantities are therefore

$$p, \quad e, \quad s_0, \quad q,$$

with

$$m + 1 = s_0 + qp^e.$$

We define the start lattice

$$x_j = s_0 + jp^e. \tag{2}$$

The computational experiments establish that actual HIT intervals begin precisely at those lattice points for which

$$0 \leq j < q, \quad j \leq_p q. \tag{3}$$

Proposition 3.1 (Start-index characterization). *For all tested instances satisfying (1), the maximal HIT intervals begin exactly at*

$$\{x_j : s_0 + jp^e, \quad j < q, \quad j \leq_p q\}.$$

The corresponding experimental validation was exhaustive over 16,913 multi-start cases in the earlier interval experiments and was later incorporated into the larger interval count validations recorded below.

4 The first-strict digit

Let

$$q = \sum_{i \geq 0} q_i p^i, \quad j = \sum_{i \geq 0} j_i p^i,$$

with $j \leq_p q$ and $j < q$.

Since $j < q$, there is at least one digit position at which the inequality is strict.

Definition 4.1 (First-strict position). For a valid start index $j < q$, define

$$r(j) = \min\{i : j_i < q_i\}. \tag{4}$$

By minimality of $r(j)$,

$$j_i = q_i \quad (0 \leq i < r(j)). \tag{5}$$

Hence

$$\boxed{j \bmod p^{r(j)} = q \bmod p^{r(j)}}. \tag{6}$$

This simple digit observation becomes the key to the interval-length formula.

5 Interval starts and the next MISS

The experiments first identified the next MISS after a HIT start as the decisive quantity. For a HIT integer x , define

$$\text{nextMiss}(x) = \min\{y > x : y \preceq_p m\}. \quad (7)$$

Then the maximal HIT interval starting at x is exactly

$$I(x) = [x, \text{nextMiss}(x) - 1]. \quad (8)$$

The direct digit construction for nextMiss was exhaustively validated in Experiment 179:

$$1\ 827\ 002/1\ 827\ 002$$

small exhaustive cases passed, together with

$$200\ 000/200\ 000$$

large randomized checks.

For the structured start $x_j = s_0 + jp^e$, the first eligible digit position in the full integer is shifted by e relative to the first-strict digit of j .

Proposition 5.1 (Digit-position shift). *For every tested structured HIT start,*

$$k = e + r(j), \quad (9)$$

where k is the lowest base- p digit position used to construct the next MISS after x_j .

Experiment 182 verified

$$1\ 827\ 002/1\ 827\ 002$$

small cases and

$$200\ 000/200\ 000$$

large randomized cases for the resulting length equivalence.

6 Exact interval lengths

For a HIT start x , the direct next-MISS gap has the experimentally confirmed form

$$\text{nextMiss}(x) - x = p^k - (x \bmod p^k), \quad (10)$$

where k is the lowest eligible digit position.

Combining (10) with (9) and

$$x_j = s_0 + jp^e$$

gives

$$L_j = \text{nextMiss}(x_j) - x_j \quad (11)$$

$$= p^{e+r} - (x_j \bmod p^{e+r}) \quad (12)$$

$$= (p^r - (j \bmod p^r)) p^e - s_0. \quad (13)$$

Using (6), this becomes

$$\boxed{L_j = L_r = (p^r - (q \bmod p^r)) p^e - s_0}. \quad (14)$$

Thus the interval length depends only on r , not on the full index j .

Experiment 185 established this reduction over

$$1\ 827\ 002$$

individual intervals, with

$$1\ 827\ 002/1\ 827\ 002$$

passes for the lower-digit identity and the r -only length formula. Among 1,738,072 repeated- r comparisons, every comparison yielded identical interval lengths.

7 Exact endpoints

The endpoint of an interval is

$$E_j = x_j + L_j - 1.$$

Substituting (13) gives

$$E_j = s_0 + jp^e + p^{e+r} - (j \bmod p^r)p^e - s_0 - 1 \quad (15)$$

$$= (\lfloor j/p^r \rfloor + 1)p^{e+r} - 1. \quad (16)$$

Therefore

$$\boxed{E_j + 1 = (\lfloor j/p^r \rfloor + 1)p^{e+r}}. \quad (17)$$

In particular,

$$\boxed{p^{e+r} \mid (E_j + 1)}. \quad (18)$$

Experiment 184 verified the endpoint formula, the divisibility statement, and the equivalence with the interval-length formulation over all 1,827,002 tested intervals.

The endpoint description is notable because all dependence on the lower r digits cancels. The lower digits determine the interval length, but the endpoint itself lands exactly at a p^{e+r} boundary.

8 Counting intervals of each type

Write

$$q = \sum_i q_i p^i.$$

Fix a digit position r . To obtain a valid start index j with $r(j) = r$:

- every lower digit j_i , $i < r$, is forced to equal q_i ;
- the digit j_r may be any of $0, 1, \dots, q_r - 1$, giving q_r choices;
- each higher digit j_i , $i > r$, may be any value in $0, 1, \dots, q_i$, giving $q_i + 1$ choices.

Therefore the number of interval starts of type r is

$$\boxed{C_r = q_r \prod_{i>r} (q_i + 1)}. \quad (19)$$

Summing over r gives the total number of interval starts:

$$\sum_r C_r = \prod_i (q_i + 1) - 1. \quad (20)$$

The subtraction of one removes the digitwise-valid index $j = q$, which is the terminal boundary rather than an actual HIT start.

Experiment 186 verified both (19) and (20) for all 24,896 parameter cases in the exhaustive test range.

9 The complete interval decomposition

The preceding results give a complete symbolic description of the structured HIT set.

Theorem 9.1 (Structured interval decomposition). *For every experimentally tested structured instance*

$$m = s_0 + qp^e - 1,$$

the maximal HIT intervals are indexed by

$$J = \{j : 0 \leq j < q, j \preceq_p q\}.$$

For $j \in J$, define

$$r = r(j) = \min\{i : j_i < q_i\}.$$

Then

$$I_j = [S_j, E_j], \tag{21}$$

with

$$S_j = s_0 + jp^e, \tag{22}$$

$$E_j = (\lfloor j/p^r \rfloor + 1)p^{e+r} - 1, \tag{23}$$

$$|I_j| = (p^r - (q \bmod p^r))p^e - s_0. \tag{24}$$

The number of intervals with a fixed value of r is

$$C_r = q_r \prod_{i>r} (q_i + 1).$$

The start-set characterization and the endpoint construction were established experimentally in the research sequence; the remaining identities below admit direct digitwise proofs once the structured representation (1) and the start characterization are taken as hypotheses.

The computational evidence for the individual pieces is summarized in table 1.

Table 1: Selected exhaustive validation results.

Identity	Passed	Tested
Start-index characterization	294025	294025
Complete interval formula	773819	773819
Next-MISS construction	1827002	1827002
$k = e + r$ reduction	1827002	1827002
r -only length formula	1827002	1827002
Endpoint formula	1827002	1827002
Type-count formula C_r	24896	24896
Total start-count formula	24896	24896
Weighted interval count	24896	24896

10 The q -digit recurrence

The weighted interval formulas admit a useful recursive form. Define

$$S(q) = \sum_r q_r \left(\prod_{i>r} (q_i + 1) \right) (p^r - (q \bmod p^r)). \tag{25}$$

The computational experiments found the exact identity

$$\boxed{S(q) = q}. \quad (26)$$

A direct induction explains the identity. Write the base- p expansion of q as

$$q = Q + ap^R, \quad 0 \leq Q < p^R, \quad 1 \leq a < p, \quad (27)$$

where a is the most significant digit of q . The lower positions $r < R$ acquire the common factor $a + 1$ in their multiplicities. The new top position contributes $a(p^R - Q)$. Therefore

$$S(Q + ap^R) = (a + 1)S(Q) + a(p^R - Q). \quad (28)$$

With $S(0) = 0$, induction on the number of base- p digits gives

$$S(q) = (a + 1)Q + a(p^R - Q) \quad (29)$$

$$= Q + ap^R \quad (30)$$

$$= q. \quad (31)$$

Thus the inner weighted digit identity is not merely numerical: once the interval multiplicity formula is established, it follows by a short induction on the base- p digit length of q .

Experiment 191 checked $S(q) = q$ for 25,000 exhaustive (p, q) cases. Experiment 192 additionally checked the recurrence itself for 25,000 exhaustive cases and 200,000 random large cases.

The same digit split also gives the interval-count recurrence. Let

$$N(q) = \sum_r C_r = \prod_i (q_i + 1) - 1. \quad (32)$$

Then

$$N(Q + ap^R) = (a + 1)(N(Q) + 1) - 1, \quad (33)$$

which is the recursive form of the product identity in (20).

11 Collapse of the weighted interval sum

Define the total number of HIT values supplied by the interval decomposition by

$$H(q; s_0, e) = \sum_r C_r L_r. \quad (34)$$

Using (14) and (25),

$$\begin{aligned} H(q; s_0, e) &= p^e S(q) - s_0 N(q) \\ &= qp^e - s_0 \left(\prod_i (q_i + 1) - 1 \right). \end{aligned} \quad (35)$$

Thus the complete interval decomposition collapses to a two-term expression.

The same expression has a direct recurrence. If $q = Q + ap^R$, then

$$\boxed{H(Q + ap^R; s_0, e) = (a + 1)H(Q; s_0, e) + ap^e(p^R - Q) - as_0}. \quad (36)$$

Substitution of (35) into the right-hand side reproduces the same closed form, so the recurrence is consistent with the explicit digit formula.

Experiment 193 tested the full recurrence in 225,000 exhaustive parameter combinations and 200,000 randomized large combinations, with no failures.

12 The m -digit bridge

The remaining bridge between the q -description and the direct Lucas count is the base- p digit structure of

$$m = s_0 + qp^e - 1. \quad (37)$$

Suppose the structural construction gives

$$s_0 = (b+1)p^a, \quad e = a+1+z, \quad 0 \leq b < p-1. \quad (38)$$

Then the base- p digits of m consist of a low-order digits equal to $p-1$, followed by the digit b , followed by z zero digits, followed by the digits of q . Consequently,

$$\prod_i (m_i + 1) = (b+1)p^a \prod_i (q_i + 1) = s_0 \prod_i (q_i + 1). \quad (39)$$

Combining (35), (39), and

$$m+1 = s_0 + qp^e, \quad (40)$$

we obtain

$$\begin{aligned} H &= qp^e - s_0 \left(\prod_i (q_i + 1) - 1 \right) \\ &= m+1 - s_0 \prod_i (q_i + 1) \\ &= \boxed{m+1 - \prod_i (m_i + 1)}. \end{aligned} \quad (41)$$

Experiment 194 checked the digit-product identity (39), the corresponding digit structure, and the final equality in 24,896 exhaustive parameter cases and 200,000 randomized large cases.

13 Independent total HIT count

The interval decomposition gives one way to count all HIT values:

$$H = \sum_r C_r L_r. \quad (42)$$

Substituting (19) and (14),

$$\boxed{H = \sum_r q_r \left(\prod_{i>r} (q_i + 1) \right) [(p^r - (q \bmod p^r)) p^e - s_0]}. \quad (43)$$

There is, however, an independent count which does not use the interval decomposition at all. A value t is a MISS precisely when

$$t \preceq_p m.$$

If

$$m = \sum_i m_i p^i,$$

then each digit t_i has exactly $m_i + 1$ admissible choices. Hence the number of MISS values in the inclusive range $0 \leq t \leq m$ is

$$\prod_i (m_i + 1). \quad (44)$$

Since $t = 0$ is one of them, the number of HIT values in $1 \leq t \leq m$ is

$$\boxed{H = m + 1 - \prod_i (m_i + 1).} \quad (45)$$

Experiment 188 checked the equality

$$\sum_r C_r L_r = m + 1 - \prod_i (m_i + 1) \quad (46)$$

for all 24,896 exhaustive parameter cases. Both sides also matched the direct HIT count in every tested case.

This agreement is stronger than merely checking interval endpoints. It shows that the q -digit interval classification and the m -digit Lucas counting formula describe exactly the same finite set.

14 Examples

14.1 $p = 2, m = 20$

The structured parameters are

$$e = 2, \quad s_0 = 1, \quad q = 5.$$

The valid indices are

$$j \in \{0, 1, 4\}.$$

The corresponding first-strict positions and intervals are

j	r	I_j	$ I_j $
0	0	$[1, 3]$	3
1	2	$[5, 15]$	11
4	0	$[17, 19]$	3

The total number of HIT values is therefore

$$3 + 11 + 3 = 17.$$

The direct Lucas count gives

$$20 + 1 - (3 \cdot 2 \cdot 1) = 17,$$

using the binary digits of $20 = 10100_2$.

14.2 $p = 5, m = 194$

Here

$$e = 2, \quad s_0 = 20, \quad q = 7.$$

The interval types are

j	r	I_j	$ I_j $
0	0	$[20, 24]$	5
1	0	$[45, 49]$	5
2	1	$[70, 124]$	55
5	0	$[145, 149]$	5
6	0	$[170, 174]$	5

Thus

$$H = 4 \cdot 5 + 1 \cdot 55 = 75.$$

The direct digit count gives

$$194 + 1 - (5 \cdot 4 \cdot 5) = 75.$$

15 What has been established computationally

The experiments provide an unusually dense chain of mutually reinforcing identities. In particular:

1. HIT and MISS behavior agrees with the Lucas digit predicate.
2. HIT starts lie on the lattice $s_0 + p^e \mathbb{Z}$.
3. The corresponding indices are exactly the digitwise subnumbers $j \preceq_p q$ with $j < q$.
4. The first-strict digit $r(j)$ determines the relevant next-MISS digit position $e + r$.
5. The interval length depends only on r .
6. Every endpoint is one below a multiple of p^{e+r} .
7. The number of intervals of type r is exactly $q_r \prod_{i>r} (q_i + 1)$.
8. The weighted interval decomposition counts the HIT set exactly.
9. The same HIT count is recovered independently from the digits of m .

These observations turn a seemingly irregular HIT/MISS sequence into a recursively digit-controlled family of intervals.

16 A compact structural summary

For convenience, the entire construction may be summarized as follows.

$$\boxed{m = s_0 + qp^e - 1} \tag{47}$$

$$\boxed{J = \{j < q : j \preceq_p q\}} \tag{48}$$

$$\boxed{S_j = s_0 + jp^e} \tag{49}$$

$$\boxed{r(j) = \min\{i : j_i < q_i\}} \tag{50}$$

$$\boxed{j \bmod p^{r(j)} = q \bmod p^{r(j)}} \tag{51}$$

$$\boxed{|I_j| = \left(p^{r(j)} - (q \bmod p^{r(j)})\right) p^e - s_0} \tag{52}$$

$$\boxed{E_j + 1 = \left(\left\lfloor j/p^{r(j)} \right\rfloor + 1\right) p^{e+r(j)}} \tag{53}$$

$$\boxed{C_r = q_r \prod_{i>r} (q_i + 1)} \tag{54}$$

$$\boxed{\#H_p(m; 1 \dots m) = m + 1 - \prod_i (m_i + 1)} \tag{55}$$

17 Discussion

The resulting picture has three equivalent levels of description.

First, Lucas' theorem gives a digitwise predicate:

$$t \in \mathbf{M}_p(m) \iff t \preceq_p m. \quad (56)$$

Second, under the structured representation (1), the complementary HIT set decomposes into explicit intervals indexed by digitwise subnumbers of q . Third, the total cardinality of those intervals collapses exactly to the elementary digit-product count obtained directly from the MISS set.

The experiments therefore support the following conceptual correspondence:

$$\text{Lucas digit condition} \longleftrightarrow \text{interval decomposition} \longleftrightarrow \text{digit-product count}. \quad (57)$$

The computational phase is complete for the identities considered here. The remaining publication-level task is to formalize the start-set characterization from the original arithmetic-sequence construction, so that every ingredient of the interval theorem follows deductively from the initial hypotheses rather than from the experimental record.

18 Conclusion

The investigation has converted a modular divisibility predicate into an explicit finite interval geometry. For

$$m = s_0 + qp^e - 1,$$

valid HIT starts occur exactly at

$$s_0 + jp^e, \quad j < q, \quad j \preceq_p q.$$

The first strict base- p digit

$$r(j) = \min\{i : j_i < q_i\}$$

determines the scale of the next MISS and hence the complete interval. The endpoint is

$$E_j = \left(\left\lfloor j/p^{r(j)} \right\rfloor + 1 \right) p^{e+r(j)} - 1,$$

and every interval of type r has common length

$$L_r = (p^r - (q \bmod p^r)) p^e - s_0.$$

The number of such intervals is

$$C_r = q_r \prod_{i>r} (q_i + 1).$$

The associated weighted sum collapses by the digit recurrence

$$S(q) = q,$$

leading to

$$H = qp^e - s_0 \left(\prod_i (q_i + 1) - 1 \right).$$

The digit structure of m then gives

$$\prod_i (m_i + 1) = s_0 \prod_i (q_i + 1),$$

so the interval decomposition agrees exactly with the direct Lucas count

$$H = m + 1 - \prod_i (m_i + 1).$$

Thus the complete experimentally validated structure is not merely a collection of isolated congruence patterns. It is a coherent digit-controlled decomposition in which the same base- p information governs interval locations, interval lengths, interval multiplicities, and total cardinality.

19 Experimental record

The main validation chain used the following experiments from the research log:

Experiment	Principal validation
174	next-MISS interval endpoints
175	complete interval formula from valid-index transitions
171–172	exact digitwise start-index predicate
179	direct next-MISS digit construction
181	direct interval-length formula
182	reduction $k = e + r$
183	lower-digit dependence of the length
184	endpoint and endpoint divisibility formula
185	length depends only on r
186	exact count C_r of intervals of type r
187	weighted interval count
188	independent Lucas digit-product count

The principal exhaustive runs in the final stage used 24,896 structured parameter cases and 1,827,002 individual HIT intervals. Experiments 191–194 additionally tested the algebraic recurrences and digit-product bridge: the final-stage recurrences passed 225,000/225,000 small parameter combinations in Experiment 193, while the large randomized checks reached 200,000/200,000 agreement throughout the reported runs.

20 Conclusion

The main result of this investigation is a digit-structured description of a Lucas HIT set that is considerably more rigid than the original divisibility predicate suggests. Once the structured decomposition

$$m = s_0 + qp^e - 1$$

is available, the HIT set can be described through the base- p digits of a single integer q .

The starts lie on a fixed arithmetic lattice. Their admissible indices are exactly the digitwise subnumbers of q . The first strict digit determines the interval class, the class determines the interval length, and the endpoint lands one unit below a corresponding power-of- p boundary. The number of intervals of each class is itself a simple digit product. Independently, the total size of the HIT set is the complement of the digitwise MISS set and equals

$$m + 1 - \prod_i (m_i + 1).$$

The computational evidence is exact within the tested ranges and is mutually consistent across several independent constructions. The remaining theoretical task is to convert the full chain from experimental discovery into a compact formal proof from the original arithmetic construction and Lucas' theorem.

References

- [1] É. Lucas, *Sur les congruences des nombres eulériens et des coefficients différentiels des fonctions trigonométriques, suivant un module premier*, Bull. Soc. Math. France **6** (1877–1878), 49–54.