

Chapter 1.3

Quantifiers, Predicates and Validity

Instructor: Abhishek Santra
Email: abhishek.santra@uta.edu

1

Variable

➤ Definition

- A symbol that stands for or represents an **individual** in a **collection or set**.

➤ Example

- The variable x may stand for one of the days.
- We may let $x = \mathbf{Monday}$ or $x = \mathbf{Tuesday}$, etc.

2

Incomplete and Complete Statement

- Definition (Incomplete Statement)
 - A sentence containing a variable is called an **incomplete statement**.
- Example
 - Example of an incomplete statement: “ x has 30 days.”
 - Here, x can be any month and substituting that, we will get a **complete statement**.
- An incomplete statement is about the **individuals in a definite domain or set**. When we *replace the variable by the name of an individual* in the set, we obtain a **statement about that individual**.

Fall 2022



CSE2315: Abhishek Santra



3

3

Quantifiers

- Quantifiers:
 - Quantifiers are phrases that refer to given quantities, such as “for some” or “for all” or “for every,” indicating ***how many objects have a certain property***.
- Two kinds of quantifiers:
 - **Universal** and **Existential**
- **Universal Quantifier**: represented by \forall
 - The symbol is translated as and means “for all”, “given any”, “for each,” or “for every,” and is known as the universal quantifier.
- **Existential Quantifier**: represented by \exists
 - The symbol is translated as and means variously “for some,” “there exists,” “there is a,” or “for at least one”.

Fall 2022



CSE2315: Abhishek Santra



4

4

Predicates

- The verbal statement that describes **the property of a variable**.
- Usually represented by the letter P, the notation $P(x)$ is used to represent some unspecified **property or predicate** that x may have
 - e.g. $P(x) = x$ has 30 days.
 - $P(\text{April}) = \text{April has 30 days}$.
- Combining the quantifier and the predicate, we get a complete statement of the form $(\forall x)P(x)$ or $(\exists x)P(x)$.
- The collection of objects is called the **domain of interpretation**.
- **Truth value of expressions** formed using quantifiers and predicates
 - What is the truth value of $(\forall x)P(x)$ where x is all the months and $P(x) = x$ has less than 32 days.
 - Undoubtedly, the above is true since no month has 32 days.



Truth value of the following expressions

- Truth of expression $(\forall x)P(x)$
 1. $P(x)$ is the property that x is yellow, and the domain of interpretation is the collection of all flowers: **Not True**
 2. $P(x)$ is the property that x is a plant, and the domain of interpretation is the collection of all flowers: **True**
 3. $P(x)$ is the property that x is positive, and the domain of interpretation consists of integers: **Not True**
 - Can you find one interpretation in which both $(\forall x)P(x)$ is true and $(\exists x)P(x)$ is false? **Not possible**
 - Can you find one interpretation in which both $(\exists x)P(x)$ is true and $(\forall x)P(x)$ is false? **Case 1 as mentioned above**
- Predicates involving properties of single variables : **unary predicates**
- **Binary, ternary and n -ary** predicates are also possible.
 - $(\forall x)(\exists y)Q(x, y)$ is a binary predicate. This expression reads as “for every x there exists a y such that $Q(x, y)$ ”, say x is a student of y ”



Interpretation

- An **interpretation** for an expression (e.g. $(\forall x)P(x)$), consists of the following:
 - A **collection of objects**, called the **domain of interpretation**, which must include at least one object. E.g., x is **all integers**
 - An **assignment of a property** of the objects in the domain to **each predicate** in the expression. E.g., $P(x) = x < 10$
 - An **assignment of a particular object** in the domain to **each constant symbol** in the expression. E.g., $P(7) = 7 < 10$
 - **How many interpretations are possible?**
- **Predicate wffs** can be built similar to *propositional wffs* using **logical connectives with predicates and quantifiers**.
- Examples of **predicate wffs**
 - $(\forall x)[P(x) \rightarrow Q(x)]$
 - $(\forall x)((\exists y)[P(x, y) \vee Q(x, y)] \rightarrow R(x)$
 - $S(x, y) \wedge R(x, y)$

Fall 2022



CSE2315: Abhishek Santra



7

7

Instant Exercises

- What is the **truth value** of each of the following wffs in the interpretation where the domain consists of the **integers**, $O(x)$ is “ x is odd,” $L(x)$ is “ $x < 10$,” and $G(x)$ is “ $x > 9$ ”?
 - A. $(\exists x) O(x)$ **True**
 - B. $(\forall x)[L(x) \rightarrow O(x)]$ **Not True**
 - C. $(\exists x)[L(x) \wedge G(x)]$ **Not True**
 - D. $(\forall x)[L(x) \vee G(x)]$ **True**
- $(\forall x)(\exists y) Q(x, y)$ vs. $(\exists y)(\forall x) Q(x, y)$?
 - Domain of interpretation is integers
 - $Q(x, y) : x < y$
 - $(\forall x)(\exists y) Q(x, y) \rightarrow$ For any integers (x) , there is a larger integer (y)
True, imagine ‘y’ inside the for loop over ‘x’
 - $(\exists y)(\forall x) Q(x, y) \rightarrow$ There is a single integer (y) , that is larger than any integer (x) **Not True, imagine ‘y’ being initialized before the for loop over ‘x’**

Fall 2022



CSE2315: Abhishek Santra



8

8

Scope of a variable in an expression

➤ **Brackets** are used wisely to identify the **scope of the variable**.

- $(\forall x) [\exists y][P(x, y) \vee Q(x, y)] \rightarrow R(x)$
 - Scope of $(\exists y)$ is $P(x, y) \vee Q(x, y)$ while the scope of $(\forall x)$ is the entire expression.
- $(\forall x)S(x) \vee (\exists y)R(y)$
 - Scope of x is $S(x)$ while the scope of y is $R(y)$.
- $(\forall x)[P(x, y) \rightarrow (\exists y) Q(x, y)]$
 - Scope of variable y is not defined for $P(x, y)$ hence y is called a **free variable**. Such expressions might not have a truth value at all.

➤ **What is the truth value of the expression**

- $\exists(x)[A(x) \wedge (\forall y)[B(x, y) \rightarrow C(y)]]$ in the interpretation, where
- $A(x)$ is “ $x > 0$ ”, $B(x, y)$ is “ $x > y$ ” and $C(y)$ is “ $y \leq 0$ ” where the domain of x is positive integers, and the domain of y is all integers

True, $x=1$ is a positive integer and any integer less than x is ≤ 0



Translation: Verbal statements to Symbolic form

➤ “**Every person is nice**” can be rephrased as
“For any thing, if it is a person, then it is nice.”

➤ So, if $P(x)$ is “ x is a person” and $Q(x)$ be “ x is nice,” the statement can be symbolized as

- $(\forall x)[P(x) \rightarrow Q(x)]$
- Variations: “All persons are nice” or “Each person is nice”
- *How about $(\forall x)[P(x) \wedge Q(x)]$?*
 - Domain: whole world, then not everything in the world is nice person.



Translation: Verbal statements to Symbolic form

- “There is a nice person” can be rewritten as “There exists something that is both a person and nice.”
 - $(\exists x)[P(x) \wedge Q(x)]$
 - Variations: “Some persons are nice” or “There are nice persons.”
 - How about $(\exists x)[P(x) \rightarrow Q(x)]$?
 - If x is a person, then x is nice?
- So **almost always**, \exists goes with \wedge (conjunction) and \forall goes with \rightarrow (implication).



Translation

- To translate an English statement into wff, **use intermediate English statement**
- The word “only” can be tricky depending on its presence in the statement.
 - X loves **only** Y \Leftrightarrow If X loves anything, then that thing is Y.
 - **Only** X loves Y \Leftrightarrow If anything loves Y, then it is X.
 - X **only** loves Y \Leftrightarrow If X does anything to Y, then it is love.
- Example for forming symbolic forms from predicate symbols
 - $J(x)$ is “x is John”; $M(x)$ is “x is Mary”; $L(x, y)$ is “x loves y”
 - John loves only Mary \Leftrightarrow
For any thing, if it is John then, if it loves anything, that thing is Mary \Leftrightarrow
 $(\forall x)[J(x) \rightarrow (\forall y)(L(x, y) \rightarrow M(y))]$
 - Only John loves Mary \Leftrightarrow
For any thing, if it is Mary then, if anything loves it, that thing is John \Leftrightarrow
 $(\forall x)[M(x) \rightarrow (\forall y)(L(y, x) \rightarrow J(y))]$
 - John only loves Mary \Leftrightarrow
For any thing, if it is John then, for any other thing, if that thing is Mary, then John loves it \Leftrightarrow
 $(\forall x)[J(x) \rightarrow (\forall y)(M(y) \rightarrow L(x, y))]$



Tips for translation to predicate wff

- Textbook p45
- Look for the key words that signify the **type of quantifier**
 - For all, for every, for any, for each : **universal quantifier**
 - For some, there exists: **existential quantifier**
- English sometimes uses “**understood**” universal quantifiers
 - “Dogs chase rabbits” is understood to mean, “All dogs chase all rabbits.”
- If you use a **universal** quantifier, the connective that goes with is almost always “**implication**”
- If you use an **existential** quantifier, the connective that goes with is almost always “**conjunction**”
- Whatever comes after the word “**only**” is the conclusion of an implication.
- You are most apt to arrive at a correct translation if you follow the **order of the English words**



Class exercise

- $S(x)$: x is a student; $I(x)$: x is intelligent; $M(x)$: x likes music
- Write wffs that express the following statements:
 - **All students are intelligent.**
 - For anything, if it is a student, then it is intelligent \Leftrightarrow
 - $(\forall x)[S(x) \rightarrow I(x)]$
 - **Some intelligent students like music.**
 - There is something that is intelligent, and it is a student, and it likes music \Leftrightarrow
 - $(\exists x)[I(x) \wedge S(x) \wedge M(x)]$
 - **Everyone who likes music is a stupid student.**
 - For anything, if that thing likes music, then it is a student and it is not intelligent \Leftrightarrow
 - $(\forall x)(M(x) \rightarrow S(x) \wedge [I(x)]')$
 - **Only intelligent students like music.**
 - For any thing, if it likes music, then it is a student and it is intelligent \Leftrightarrow
 - $(\forall x)(M(x) \rightarrow S(x) \wedge I(x))$



Negation of statements

- $A(x)$: x is fun, $(\forall x)A(x)$: Everything is fun
- Negation will be “it is false that everything is fun,” i.e., “*something is not fun.*”
- *Opposite: Every \rightarrow Some*
- In symbolic form, $[(\forall x)A(x)]' \Leftrightarrow (\exists x)[A(x)]'$
- Similarly, negation of “Something is fun” is “Nothing is fun” or “Everything is not fun/boring.”
- Hence, $[(\exists x)A(x)]' \Leftrightarrow (\forall x)[A(x)]'$



Class Exercise

- What is the negation of “Everybody loves somebody sometime.”
 - Everybody hates somebody sometime
 - Somebody loves everybody all the time
 - Everybody hates everybody all the time
 - Somebody hates everybody all the time ✓
- What is the negation of the following statements?
 - **Some pictures are old or faded.**
 - Every picture is neither old nor faded.
 - **All people are tall and thin.**
 - Someone is short or fat.
 - **Some students eat only pizza.** (Recall, the John and Mary example)
 - **Only students get t-shirts.**



Negation Approach

➤ What is the negation of the following statements?

▪ **Some students eat only pizza.**

– Recall, John loves only Mary example: $(\forall x)[J(x) \rightarrow (\forall y)(L(x, y) \rightarrow M(y))]$

– Let $S(x)$: x is student, $E(x, y)$: x eats y, $P(x)$: x is pizza

– However, “John” is “some students”. Thus, change the quantifier

– Final Expression: $(\exists x)[S(x) \rightarrow (\forall y)(E(x, y) \rightarrow P(y))]$

– Take Negation: $((\exists x)[S(x) \rightarrow (\forall y)(E(x, y) \rightarrow P(y))])'$

1. $(\forall x)[S(x) \rightarrow (\forall y)(E(x, y) \rightarrow P(y))]'$

2. $(\forall x)[S(x)' \vee (\forall y)(E(x, y)' \vee P(y))]'$

1, Imp

3. $(\forall x)[S(x) \wedge ((\forall y)(E(x, y)' \vee P(y)))]'$

2, DM

4. $(\forall x)[S(x) \wedge (\exists y)(E(x, y)' \vee P(y))]'$

3, Negation

5. $(\forall x)[S(x) \wedge (\exists y)(E(x, y) \wedge P(y)')]$

4, DM

▪ **Every student eats something that is not pizza.**

Fall 2022



CSE2315: Abhishek Santra



20

20

Negation Approach

➤ What is the negation of the following statements?

▪ **Only students get t-shirts.**

– Let $S(x)$: x is student, $T(x)$: x gets t-shirt

– Final Expression: $(\forall x)[T(x) \rightarrow S(x)]$

– Take Negation: $((\forall x)[T(x) \rightarrow S(x)])'$

1. $(\exists x) [T(x) \rightarrow S(x)]'$

2. $(\exists x)[T(x)' \vee S(x)]'$

1, Imp

3. $(\exists x)[T(x) \wedge S(x)']$

2, DM

▪ **There is a non-student who gets a t-shirt.**

Fall 2022



CSE2315: Abhishek Santra



21

21

Validity of Predicate wff

- Analogous to a **tautology of propositional logic**.
- Truth of a predicate wff *depends on the interpretation*.
- A predicate wff is **valid** if it is true in all possible **interpretations** just like a propositional wff is true if it is true for all rows of the truth table.
- A valid predicate wff is intrinsically true.

	Propositional Wffs	Predicate Wffs
Truth values	True or false – depends on the truth value of statement letters	True, false or neither (if the wff has a free variable)
Intrinsic truth	Tautology – true for all truth values of its statements	Valid wff – true for all interpretations
Methodology	Truth table to determine if it is a tautology	No algorithm to determine validity (Use Reasoning; Can you find an interpretation where the expression is false?)

Fall 2022



CSE2315: Abhishek Santra



22

22

Validity Examples

- $(\forall x)P(x) \rightarrow (\exists x)P(x)$
 - This is **valid** because if every object of the domain has a certain property, then there exists an object of the domain that has the same property.
- $(\forall x)P(x) \rightarrow P(a)$
 - **Valid** – quite obvious since a is a member of the domain of x .
- $(\exists x)P(x) \rightarrow (\forall x)P(x)$
 - **Not valid** since the property cannot be valid for **all objects** in the domain if it is valid for **some objects** of that domain. Can use a **mathematical context** to check as well.
 - Say $P(x) = "x \text{ is even}"$, then there exists an integer that is even but not every integer is even (domain: integer)
- $(\forall x)[P(x) \vee Q(x)] \rightarrow (\forall x)P(x) \vee (\forall x)Q(x)$
 - **Invalid**, can show by mathematical context by taking $P(x) = x \text{ is even}$ and $Q(x) = x \text{ is odd}$ (domain: integer). **What is the scope of 'x' in each case?**
 - In that case, the hypothesis is true but the conclusion is false because it is not the case that every integer is even or that every integer is odd.

Fall 2022



CSE2315: Abhishek Santra



23

23

Class + Home Exercise - 1

- What is the truth of the following wffs where the domain consists of **integers**:

1. $(\forall x)[L(x) \rightarrow O(x)]$,
where $O(x)$ is “ x is odd” and $L(x)$ is “ $x < 10$ ”?
False since not all integers < 10 are odd, e.g., 4, 6
2. $(\exists y)(\forall x)(x + y = 0)$?
False, no one y works for all x 's, imagine y being initialized before the “for loop” over x
3. $(\exists y)(\exists x)(x^2 = y)$, where $x, y > 1$?
True, e.g., $y = 4$ and $x = 2$
4. $(\forall x)[x < 0 \rightarrow (\exists y)(y > 0 \wedge x + y = 0)]$?
True, pick $y = -x$



Class + Home Exercise - 2

- Using predicate symbols and appropriate quantifiers, write the symbolic form of the following English statement:

- $D(x)$ is “ x is a day” M is “Monday” T is “Tuesday”
- $S(x)$ is “ x is sunny” $R(x)$ is “ x is rainy”

1. Some days are sunny and rainy.
 $(\exists x)(D(x) \wedge S(x) \wedge R(x))$
2. It is always a sunny day only if it is a rainy day.
 $(\forall x)[D(x) \wedge S(x) \rightarrow D(x) \wedge R(x)]$
3. It rained both Monday and Tuesday.
 $R(M) \wedge R(T)$
4. Every day that is rainy is not sunny.
 $(\forall x)[D(x) \wedge R(x) \rightarrow (S(x))']$



Discussion



Fall 2022



CSE2315: Abhishek Santra



26