

# Effects of Aim Training Performance, Lifestyle Habits, and Environmental Factors on Fatigue and Stress

Ian Chen\*

Yan Yi Lance Du†

May 11, 2025

## 1 Abstract

**Purpose:** In this study, we investigated the effects of lifestyle, environment, and response time on students' mental state.

**Methods:** Participants were 145 students from an introductory statistics class at the University of Illinois, Urbana Champaign. Participants played an aim training game and completed a post-game survey that assessed personal and environmental factors that might affect aim training performance (i.e., reaction time). We operationalized mental state as students' stress and fatigue levels.

**Results:** We found that an increase in response time is associated with larger stress levels. Moreover, a distractive environment led to increased stress, but is counteracted by being active in sports. The only predictors associated with decreased fatigue is getting enough sleep. We also verified that aggregating the predictors led to an improved model.

**Implications:** Students tend to face high levels of stress and fatigue in academic settings. Our study implies that different interventions may be required for students to mitigate the negative effects of stress on performance. Indeed, future studies should consider all not only response times, but also lifestyle, environmental factors before proposing remedies for Fatigue and Stress.

## 2 Introduction

In this study, we explore which indicators are significantly correlated with reaction time, a proxy for their performance. We study both self-reported fatigue and stress levels, as well as their lifestyle habits and environment.

### 2.1 Research Questions

Here, we have three primary research questions (RQ):

1. Does aim training performance correlate to impaired mental state?
2. Are there environmental factors consistent with impaired mental state?
3. Are there lifestyle factors consistent with impaired mental state?

We hypothesize that worse performance on aim trainers would lead to more fatigue and stress levels. We expect poor environments to affect both stress and fatigue equally. We think that a poor lifestyle would lead to increased fatigue, and may not affect stress.

---

\*Writing, Analysis, EDA, Model Building

†Reviewing and Editing, Analysis, EDA, Model Building

### 3 Materials and Methods

We collect our dataset from introductory statistics students at the University of Illinois, Urbana Champaign. An online survey was posted that all students needed to fill in, and was graded (i.e. it was not anonymized). Students had a week to submit the form, and could do so asynchronously.

All students submitted their scores for their first three attempts. In this study, we will take their median response.

First, we only consider datapoints that have Stable Wifi (so we remove 2 points). Moreover, to make sure we are not overfitting, we partition into 90% training and 10% testing. Fitting models on the training set, we verify *a posteriori* whether the results are consistent on the testing set. Finally, we conduct a differential study (fitting different models) on each category.

### 4 Exploratory Data Analysis

Table 1: Fatigue and Stress counts on Training Dataset

Level	FatigueMouse	StressMouse	FatigueTrack	StressTrack
Very Low	7	8	14	16
Low	15	12	28	19
Moderate	18	19	19	20
High	7	8	7	11
Very High	1	1	1	3
Total	48	48	69	69

Table 2: Game and Sports counts on Training Dataset

Level	GameMouse	SportsMouse	GameTrack	SportsTrack
Daily	7	3	5	3
Several times a week	17	10	15	13
Once a week	2	2	9	5
Several times a month	5	7	15	11
Rarely	12	20	20	27
Never	5	6	5	10
Total	48	48	69	69

We see that there are very few datapoints for the *Very High* observations in our training dataset, and hence we remove these from the dataset (Table 1).

We see that the response variable is roughly normally distributed across both mouse and trackpad groups (Figure 1).

### 5 Research Question One

We investigate separately on the Mouse and Trackpad users.

#### 5.1 Mouse

Empirically, we see that for mouse users, the higher the aim training time level, the larger the stress. Indeed, an increase of 200ms would lead to half a unit increase of stress level. However, there seems to be no no




Figure 1: Distribution of Response Variable on the Training Dataset. The x-axis describes the median response time in milliseconds. The y-axis describes the number of results.




Figure 2: Correlation Matrix

Table 3: Descriptive Statistics

	mean	sd	median	trimmed	mad	min	max	range	skew	kurtosis	se
Time	559.99	107.72	565	561.06	91.92	298	868	570	-0.05	-0.04	10.22
StressN	2.47	1.02	3	2.46	1.48	1	4	3	-0.04	-1.14	0.10
FatigueN	2.36	0.92	2	2.33	1.48	1	4	3	0.13	-0.85	0.09
NoiseLevel	3.14	1.89	3	2.92	1.48	1	10	9	0.95	0.54	0.18
AvgHoursExercise	4.74	4.49	4	4.11	2.97	0	30	30	2.65	10.66	0.43
AvgSleepTime	7.30	1.03	7	7.23	1.48	5	11	6	0.74	1.31	0.10
LastNightSleep	7.28	1.93	7	7.24	1.48	1	17	16	0.64	5.64	0.18




Figure 3: Median response time based on Fatigue and Stress levels. The y-axis shows the median response time across three trials. Left: Increasing Fatigue. Right: Increasing Stress.

clear trend for fatigue (Figure 3).

There are no outliers, determined by studentized statistics ( $\alpha = 0.05$ , with Bonferroni correction). We also tested for normality using the Shapiro-Wilks test. The residuals for both stress and fatigue SLR models seems non-normal ( $p < 0.05$ ). We tested for homoscedasticity using the B-P test. The constant variance assumption held for both stress ( $p = 0.77$ ) and fatigue ( $p = 0.74$ ) SLR models. All Cook's Distances were less than 1, so there are no highly influential points.

## 5.2 Trackpad

On the Trackpad condition, the trend is a lot less clear (Figure 4). Indeed, neither *Stress* nor *Fatigue* are significant predictors now ( $p > 0.05$ ).

There are no outliers, determined by studentized statistics ( $\alpha = 0.05$ , with Bonferroni correction). We also tested for normality using the Shapiro-Wilks test. The residuals for both stress and fatigue SLR models seems normal ( $p > 0.05$ ). We tested for homoscedasticity using the B-P test. The constant variance assumption held for both stress ( $p = 0.64$ ) and fatigue ( $p = 0.14$ ) SLR models. All Cook's Distances were less than 1, so there are no highly influential points.




Figure 4: Median response time based on Fatigue and Stress levels for Trackpad. The y-axis shows the median response time across three trials. Left: Increasing Fatigue. Right: Increasing Stress.

## 6 Research Question Two

Now, we try to determine if there are correlations between Stress and Fatigue, based on environmental factors such as *Distraction*, *NoiseLevel*.

Being in a highly distractive environment led to an increase in stress ( $p = 0.0126$ ). Environmental factors did not affect one's fatigue ( $p > 0.05$ ).

## 7 Research Question Three

Now, let us now consider if there are correlations between Stress and Fatigue based on lifestyle habits, including sleep, exercise, sports, and gaming activity.

First, being active in sports decreases stress levels by half a unit ( $p = 0.084$ ). Indeed, this is the largest drop in stress levels out of any predictor.

Getting more sleep is also strongly related to decreases in fatigue ( $p = 0.0075$ ).

## 8 Model Building

We have studied three classes of predictors: aim training response time, lifestyle (Sports and Gaming), as well as environmental (Distraction). It is clear that fatigue is only related to lifestyle habits, but it is unknown which of the classes of predictors are most important for stress. Here, we synthesize the results and perform model comparison.

Running backwards variable selection using AIC, we find that we should use the predictors ActiveSports and Distraction. Our final model, using predictors ActiveSports and Distraction, is

$$Stress \sim \begin{cases} 2.463 & \text{if not active in sports and not distractive environment} \\ 2.018 & \text{if active in sports and not distractive environment} \\ 3.085 & \text{if not active in sports but distractive environment} \\ 2.640 & \text{if active in sports but distractive environment} \end{cases}$$

Running backwards variable selection using AIC, we find that we should use the predictors ActiveSports and Distraction to predict Stress. Our final model, using predictors ActiveSports and Distraction, is

$$Stress \sim \begin{cases} 2.463 & \text{if not active in sports and not distractive environment} \\ 2.018 & \text{if active in sports and not distractive environment} \\ 3.085 & \text{if not active in sports but distractive environment} \\ 2.640 & \text{if active in sports but distractive environment} \end{cases}$$

Running backwards variable selection using AIC, we find that we should use the predictors LastNightSleep and Distraction to predict Fatigue. Our final model, using predictors LastNightSleep and Distraction, is

$$Fatigue \sim \begin{cases} 3.160 - 0.118 * LastNightSleep & \text{If no distractions} \\ 3.485 - 0.118 * LastNightSleep & \text{if distractions} \end{cases}$$

## 8.1 Model Comparison

Table 4: AIC scores for different Stress models

	All	AimTraining	Environment	Lifestyle
AIC	313.8374	322.6045	316.2252	318.7321

Table 5: AIC scores for different Fatigue models

	All	AimTraining	Environment	Lifestyle
AIC	293.9788	300.0618	300.0618	294.2727

We can compare this aggregated model to the ones using predictors from each class individually. Indeed, the aggregated model has the lowest AIC score for both the Stress and Fatigue models, and is the best (Tables 4, 5).

## 8.2 a posteriori Validation

Table 6: RMSE between Training and Testing

	TrainStress	TrainFatigue	TestStress	TestFatigue
RMSE	0.9595483	1.020531	0.9244007	1.031643

We evaluate the root mean squared error (RMSE) of the stress and fatigue models on the training and testing dataset. We do not any inflated errors on the testing set, so we are probably not overfitting (Table 6).

## 9 Discussion and Conclusion

Trying to predict stress and fatigue is difficult, and our models only explained around 10% of the variation for each. However, we did gain valuable insights into which predictors are significant.

First, we determined that on trackpad, the aim training response time was too varied and there were no clear trends. For future studies using aim training time as a predictor for mental state, we recommend to

require mouse data in order to reduce the variation. Indeed, for mouse users, we were able to say that the faster response times were correlated with decreased stress.

It was surprising that the level of noise in the environment was insignificant towards both stress and fatigue, since we expected more fatigue with noisy environments. We verified that a highly distractive environment led to higher stress. Unsurprisingly, fatigue levels are strongly correlated with one's lifestyle, namely their sports activity and sleep.

Finally, we verified that using all three classes of predictors lead to an improved model for both Fatigue and Stress. Indeed, future studies should consider all not only response times, but also lifestyle, environmental factors before proposing remedies for Fatigue and Stress.