

Generalists link peaks in the shifting adaptive landscape
of Australia's dragon lizards

Ian G. Brennan^{1,2,3,*}, Natalie Cooper³, Joanna Sumner⁴, Sarin Tiatragul¹,
Leonardo G. Tedeschi⁵, Elizabeth S. Broady¹, Jane Melville⁴, J. Scott Keogh¹

Affiliations:

¹Division of Ecology & Evolution, The Australian National University, Australia

²Museum and Art Gallery of the Northern Territory, Australia

³Science Group, Natural History Museum, United Kingdom

⁴Museums Victoria Research Institute, Melbourne, Australia

⁵National Biodiversity DNA Library, Commonwealth Scientific and Industrial Research Organisation, Australia

*Corresponding author

Email Addresses:

ian.brennan@anu.edu.au

natalie.cooper@nhm.ac.uk

jsumner@museum.vic.gov.au

sarin.tiatragul@anu.edu.au

leonardo.goncalvestedeschi@csiro.au

elizabeth.broadly@anu.edu.au

jmelv@museum.vic.gov.au

scott.keogh@anu.edu.au

Statement of Authorship:

Conceptualization, IGB, JSK, NC; sampling, LGT, JS; data collection ESB, JT, IGB; data
curation, IGB, ESB; methodology, NC, IGB; analysis IGB, ST; writing - original draft, IGB;
writing - reviewing & editing, all authors; funding acquisition, JSK.

Data Accessibility: All data and code are available on GitHub at
[www.github.com/IanGBrennan/Amphibolurinae](https://github.com/IanGBrennan/Amphibolurinae)

Running Title: Generalists link adaptive peaks

Keywords: adaptive landscape, macroecology, macroevolution, phylogenomics

Article Type: Letter

Word Counts: abstract (139), main text (4934)

Number of: references (67), figures (5), tables (0), text boxes (0)

Abstract

The adaptive landscape is a flexible concept in biology which helps map entities onto an evolutionary topography. A common practice is clustering species to identify adaptive peaks but the pathways that allow movement into new adaptive zones remain poorly understood. Here we integrate new sequence-capture (>5,000 nuclear markers) and multivariate morphological data to investigate the adaptive landscape of Amphibolurinae dragon lizards (Agamidae). Our well-supported, time-calibrated phylogeny includes over 360 specimens from 176 species, including iconic taxa such as the frilled-neck lizard, thorny devil, and bearded dragon. We show that arrival in Australia was followed by rapid expansion of morphological diversity facilitated by **generalist habitat use**. Ancestral dragons were likely arboreal, but **species with broad niche breadths and intermediate morphologies** provided pathways to subsequent specialization. These findings reveal an evolutionary pathway for adaptive diversification in a major continental radiation.

Introduction

The adaptive landscape is an indispensable tool for considering and visualizing evolutionary spaces. In adapting Wright's (1932) genotypic fitness landscape to a phenotypic space, Simpson (1944) provided a shared vocabulary and conceptual bridge between micro- and macroevolution (Arnold et al. 2001). At local scales (species or populations), landscapes help us to map form to fitness. However, at broader scales (species, genera, families), landscapes (often called morphospaces or functional trait spaces) reveal the density and position of peaks, and how clades or communities fit in multidimensional space. This can help in determining the cost of moving between phenotypes or mapping ecology to morphology. Importantly, phenotypic landscapes are dynamic and can vary through time in their characteristics (Marshall 2014). This may dictate the persistence and evolution of the species they represent. Consequently, we can use data from those species to infer parameters of, and changes to, the landscape.

Despite how it is often visualized, the adaptive landscape was never limited to only two or three dimensions. Instead, landscapes can be multidimensional, which intersects with Hutchinson's (1957) niche hypervolume. In multidimensional space we can assess the volume and density of individual peaks, probe the depths of multidimensional valleys, or detect holes in the landscape hypervolume (Blonder 2016). Higher dimensionality may also provide paths between peaks that might not be identifiable with fewer dimensions. The complex topography of the landscape, including holes, dictates viable paths from existing morphologies into new ones. As such, the ideas of adaptive radiation and ecological opportunity have long been entwined with Simpson's view of movement across the landscape (Schluter 2000; Simpson 1944; Yoder et al. 2010). However, we still do not understand how new peaks form in disparate regions of the landscape. One hypothesis is that the center of morphological space acts as a hub/bridge to more distant adaptive zones (Dennis et al. 2011; Futuyma & Moreno, 1988). The expectation being that the morphological center is occupied by taxa with generalist morphologies that enable broad niche breadths, and this facilitates evolutionary movement into narrower specialist niches. With advances in computational power, development of more complex comparative methods, and larger phenotypic datasets, we can begin to unravel historical changes to the landscape and the evolution of new forms (Harmon et al. 2005). Importantly, we can ask which axes facilitate movement into new phenotypic space, and if changes to the landscape are associated with shifts in ecology, habitat, and/or geographic range.

While biologists have been visualizing the adaptive landscape and determining the relative heights of peaks since Simpson, comparatively little empirical attention has focused on how lineages populate new peaks and the landscape changes through time (but see Marshall 2014; Burin et al. 2023). This is partly due to the necessity of a high dimensional dataset, well-resolved phylogeny, and an appropriate organismal group. Here we investigate patterns of adaptive landscape change and the role of ecology and expanding ranges in the morphological evolution of Australo-Papuan agamid lizards. Amphibolurine agamids consist of ~130 species with varied and diverse ecologies, habitat preferences, and highly imbalanced biogeographic richness. The majority of species (~110 spp.) are endemic to Australia, where their distributions span the continent's driest deserts and wettest rainforests. Their morphologies reflect these diverse habitats, ranging from spiny forest dragons, squat desert pebble mimics, and include iconic species such as the frilled-neck lizard (*Chlamydosaurus*) and thorny devil (*Moloch*). To match, they show a range of ecologies from cryptic arboreal species to sand dune burrowers. These unique ecomorphologies provide an ideal opportunity to study the tempo and mode of evolution and the processes driving phenotypic change.

To address the evolution of the amphibolurine adaptive phenotypic landscape, we first reconstruct the phylogenetic relationships of Australo-Papuan dragon lizards. To investigate morphological diversity we collected an extensive phenotypic dataset that summarizes broad axes of variation across the head, body, limbs, and tail. We use phylogenetic comparative methods to assess the tempo and mode of trait evolution and from this present a picture of the evolution of amphibolurine dragon lizards. Across the dragon body, many traits show temporally and phylogenetically heterogeneous evolutionary histories. Those traits with evidence of varied evolutionary processes often exhibit large shifts in trait changes coincident with new morphologies. These morphological jumps are better described by biased directional changes which suggest selection, rather than increased evolutionary rates. Lastly, we identify that the diversity of forms of Australian dragons likely evolved from ancestors **with broad niches** that filled space in the adaptive **morphological** landscape and acted as intermediates connecting an otherwise patchy landscape.

Materials and Methods

Taxon Sampling

We assembled a sequence-capture dataset across 309 Amphibolurinae representing 120 of 136 currently recognized species, with a focus on Australian taxa (115 of 117 spp.). This sampling covers all 16 genera and many recognized subspecies (Table S1). We included outgroup

representatives from all agamid subfamilies and used tuatara (*Sphenodon punctatus*) and chicken (*Gallus gallus*) to root the tree (368 total samples).

Molecular Data Collection and Processing

Molecular sampling implemented the Squamate Conserved Loci (SqCL) kit (Singhal et al. 2017), which comprises ~5,000 ultra-conserved elements, ~400 anchored hybrid enrichment loci, and ~40 legacy genes commonly used in squamate molecular phylogenetics. Sample collection and laboratory preparation were completed as part of the Australian Amphibian and Reptile Genomics initiative (AusARG) following the protocol outlined in Tiatragul et al. (2023). Raw sequence reads are available from the Bioplatforms Australia Data Portal (<https://data.bioplatforms.com/organization/ausarg>). We processed raw sequence data using the *pipesnake* workflow (Brennan et al. 2024). A list of all software and versions used is included in the Supplement.

Phylogenetic Analyses

We estimated individual genealogies for our sequence-capture data (n=5441) under maximum-likelihood in IQ-TREE2 allowing the program to assign the best fitting substitution model using ModelFinder (Kalyaanamoorthy et al. 2017), then performed 1,000 ultrafast bootstraps (Minh et al. 2013). We then estimated the species tree using the coalescent-consistent methods wASTRAL-hybrid and ASTRAL-IV, using IQ-TREE2 gene trees as input. We used two ASTRAL methods to take advantage of the way wASTRAL-hybrid weights branch lengths and support values in estimating the species tree, and how ASTRAL-IV allows multiple individuals per species and estimates both internal and terminal branch lengths in substitutions-per-site (as opposed to coalescent units for internal branch lengths and static terminal branch lengths in wASTRAL). To quantify topological signal from individual gene trees we calculated gene concordance factors using IQ-TREE2 and focused on intergeneric relationships using a single representative for each genus.

Divergence Dating

To estimate divergence times among taxa we applied a series of fossil and secondary calibrations in MCMCTree (Rannala & Yang 2007) as outlined in the Supplement (Table S2). We started by trimming our ASTRAL tree down to a single representative for each ingroup species or subspecies for input. We reduced our molecular data to exonic markers in the AHE loci, then estimated raw genetic distances from alignments to use as a proxy for evolutionary rate. We removed the fastest 5% and slowest 5% of loci to avoid issues with extreme rate

heterogeneity, removed third codon positions from all loci, then partitioned the remaining loci into three partitions using AMAS (Borowiec 2016).

For each partition we ran MCMCTree with *usedata* = 3 to get the approximate likelihoods and branch lengths using *baseml* (dos Reis & Yang 2011), then concatenated the out.BV files. We then ran four replicate MCMCTree analyses on the gradient and Hessian (in.BV file; *usedata* = 2), each for 20k burn-in generations before collecting 20k samples at a sampling frequency of 100 generations (2,020,000 total generations). We compared mcmc files for stationarity and convergence (ESS of all parameters > 200), combined them using logCombiner, and used this combined mcmc file to summarize divergence times on our tree (*print* = -1 in .ctl file). To validate our priors we ran an additional analysis (*usedata* = 0) to run explicitly from the prior calibrations and determine our effective priors for comparison against our posterior age estimates (Fig.S1).

Morphological Data Collection

We collected 19 linear measurements for 128 amphibolurine lizard species from 482 museum samples ($\bar{x}$ = 5 per spp., min. = 1, max. = 8). Linear measurements were collected by one author (IGB) using Mitutoyo digital calipers (product 500-763-20). These measurements aimed to capture the gross morphology of the lizard body plan and are distributed across the head, body, limbs, and tail (Fig.S2). Definitions of trait measurements and adjustments are summarized in Table S3. Data processing and analyses used R (R Core Team 2024).

Defining Generalism through Habitat Use

To address the concepts of generalism and specialization, we focused on niche breadth. Specialization can occur on varied axes, with a given taxon being a generalist on one (e.g. habitat type), but a specialist on another (e.g. diet) (Poisot et al. 2011), like the thorny devil *Moloch horridus* that occurs throughout arid Australia but exclusively eats ants. We focus on the fundamental Eltonian niche of species morphologies and the realized niche through substrate usage (Devictor et al. 2010). We first discretized surface features and microhabitats used by dragon lizards into six categories which have different functional demands: aquatic, rock, terrestrial, low structure, high structure, and arboreal (see Supplement for details). We scored all species for these traits and treated the trait matrix in two ways. First, we generalized niche breadth by summing trait usage. Values of 1-6 indicate highly specialized (low values) to absolute generalist species (high values). Second, to investigate transitions among specialist and generalist states we classified taxa as either specialist (one or two substrates used) or generalist (≥ 3 substrates used) category. For both character codings we then fitted a series

of Markov models using *phytools* (Revell 2024) to compare theoretical pathways for the evolution of niche breadth and specialization in dragons. These included equal and independent transition rates among states, stepwise (increasing or decreasing breadth by one state) and loose stepwise models (multiple states can be gained or lost), and a strict model of increasing specialization. Importantly, we applied a derivative in which niche breadth acquisition and loss are allowed separate transition rates (see Supplement for more details) to all eligible models.

The Evolution of Morphological Diversity

To identify the mode of individual morphological traits we fit a series of models ranging from an unbiased random walk to rate-variable and trended. The most basic model—Brownian Motion (BM)—allows phenotypes to evolve through incremental random change. The most complex—*BayesTraits* V4’s fabric model (Pagel et al. 2022)—also follows a process of gradual change, but allows rates to shift at discrete points (changes in *evolvability*, v) on the tree and allows traits to change rapidly along individual branches through directed evolution (*trends*, β). Using *BayesTraits* V4 we fit these models and their derivatives (*fabric* with no v parameter, *fabric* with no β parameter) and estimated their marginal likelihoods using a stepping-stone sampler (500 stones each for 5,000 generations).

BayesTraits V4 implements a reversible-jump MCMC sampler, which we specified to run for 100 million generations with 10 million generations of burn-in. We summarized the output to check for convergence with a custom script *Scripts/processBTVarRates.R*, and verified that all parameters had ESS > 200. To verify model fit and consistency among runs we ran four separate runs for each trait. Individual model outputs (... *VarRates.txt*) were summarized using the *FabricPostProcessor* software and independent runs were combined using *MergeFabric*.

To track trait values at internal nodes we logged values from the MCMC chain using the *AddTag* and *AddMRCA* features in *BayesTraits* by generating custom control files using functions defined in the Supplement. We also used this script to generate priors on the root state and rate by fitting a single rate BM model using *GEIGER* (Pennell et al. 2014). We summarized posterior estimates of ancestral trait values under varied models using custom scripts included in the Supplement. To track the evolution of dragon morphospace through time we extrapolated trait values linearly along branches at 0.5 million year intervals given start and end values at nodes and constant estimated evolutionary rates. We then reduced observed and estimated ancestral values to three principal component axes and fit multidimensional hypervolumes using *hypervolume* (Blonder et al. 2018) to extract the volume.

To determine how the evolution of the empirical hypervolume might depart from expectations we simulated 100 morphological datasets under Brownian Motion and carried out the same exercise. To understand morphological distances between observed and estimated (ancestral) taxa/optima we measured Euclidean distances. We visualized functional morphological spaces using the package *funspace* (Pavanetto & Puglielli 2024).

To identify adaptive peaks across the multidimensional landscape (i.e., morphological regimes) we used an OU-based method that accounts for correlation among traits (*PhylogeneticEM*; Bastide et al. 2018). We used two datasets as input. The first included all 19 non-overlapping morphological traits. Due to the strong correlation of some traits (Fig.S3) we generated a second dataset by reducing dimensionality through principal components analysis (PCA) and choosing the top six principal component axes which accounted for >90% of the cumulative variance, where each component explained >2%. We fit the models across a number of shift values ($K=0-15$) and 30 alpha values ($\alpha=0-5$) to both datasets. For consistency, we fit several additional schemes to test the reliability of our results (see Supplement). After identifying the preferred number of optima (either 5 or 12; see Results; also called regimes) we used these as discrete characters for *RandomForest* (Liaw & Wiener, 2022) analysis to categorize ancestral taxa according to contemporary optima. Assigning species to ecological guilds is fraught with subjectivity so we relied on our morphological regimes to identify clusters of species that map loosely to existing ecological categorizations (e.g. arboreal, terrestrial) following discussion in Melville & Wilson (2019). RandomForest confusion (error) rates for almost all contemporary optima were sufficiently low (<10%), so we predicted the morphological regime of ancestral taxa (nodes) based on trait values estimated by the best fitting model for each trait. We further explored the possibility of bias in our inferences and model sensitivity through simulations (see Supplement).

Results

Molecular Data Collection and Processing

Sequence capture using the SqCL kit resulted in assembly of 5441 targets. New sampling recovered a median of 5118 loci, with a concatenated and aligned sequence length of >6 Mbp (Fig.S4). See Supplement for additional sequence statistics.

Phylogenetic Analyses

Coalescent-consistent topologies estimated by wASTRAL-hybrid and ASTRAL-IV are well resolved with high support across most branches, including good resolution among genera and

major clades (Fig.1; Fig.S5). Internal branches separating the Australian genera are short and consecutive rapid splits show high levels of incomplete lineage sorting resulting in low gene concordance factors (Fig.S6). This is notable in the positions of the *Moloch/Chelosania* and *Lophosaurus* clades and placement of *Tropicagama*. Additional areas of conflict are limited to interspecific placements of the *Ctenophorus fordi* group, the *Pogona minor* group, and the *Diporiphora pallida* group. In agreement with a recent study (Bruton et al. 2025), we find *Ctenophorus* (*Cryptagama*) *aurita*—a highly morphologically derived lineage—nested within *Ctenophorus*. Our phylogeny also supports the assignments of several frequently confused arid and savannah species, upholding the genera *Amphibolurus*, *Gowidon*, *Lophognathus*, and *Tropicagama*. We also provide strong evidence that unites *Rankinia*—the world’s highest latitude dragon—with *Pogona*. Beyond Amphibolurinae, we also provide a strongly supported estimate of subfamilial relationships in Agamidae. We recover a sister relationship between the Agaminae and Draconinae, which together are the sister to the Amphibolurinae. The next closest relatives are the Hydrosaurinae, then Uromastycinae, and lastly Leiolepidinae (Fig.1)

Divergence Dating

Amphibolurines appear in the late Cretaceous (~75 mya), separating from their sister the Agaminae/Draconinae. The crown split among amphibolurines does not occur until the Eocene/Oligocene boundary ~34 mya. Cladogenetic events occur rapidly at the base of the amphibolurine tree, with all major splits among Australian groups occurring between 30-25 mya. With the exception of the clade comprising *Chlamydosaurus*, *Lophognathus*, and *Amphibolurus*, all extant genera were established by the mid-Miocene ~15 mya. Rapid speciation events at the base of the amphibolurine tree obscure some relationships. This is likely the result of quick successive splits that occurred in a narrow temporal window, <600 ky (Fig.1).

Evolution of Niche Breadth

Discrete trait models of niche breadth favored a model in which transitions occur through neighboring ($n\pm 1$) or near-neighboring ($n\pm 2$) states and showed a faster transition rate towards decreasing niche breadth (increasing specialization) (Fig.S13). Model fitting of transitions among specialist (1 substrate) and generalist (3+ substrates) habitat usage favored a model where transition rates are faster between generalist and specialist states, and slower—but not absent—among specialist states. (Fig.2)

Phenotypic Evolution

Of 19 morphological traits, 10 were best fit by the rate-variable and trended model from *fabric*, including the summary trait encapsulating size (Table S4). The remaining nine traits were best fit by a random walk model (Brownian Motion). Traits with greater variance and disparity were more likely to fit the *fabric* model (Fig.S7). Ancestral trait values for the 10 traits best fit by *fabric* were extracted and summarized from the mcmc output files. Ancestral trait values for the remaining nine traits were estimated under Brownian Motion. This provided a complete trait matrix for living and ancestral species which we used to quantify ancestral morphologies. Morphospace showed a heterogeneous pattern of expansion with a steep increase from 30-20 ma, before a period of little expansion from 15-5 ma, and another increase in volume from 5 ma to the present (Fig.S8). Early (>20 ma) and middle (>15 ma) morphospace expansions departed from Brownian expectations.

Multi-optima OU models identified either five or 12 morphological regimes. We refer to these groups by their inferred regime identities (R1, R2 ... R5) or the clade to which they belong (*Tympanocryptis*, *Moloch*), and consider how these morphological assignments map to ecology and habitat use in the Discussion. Our multi-optima solutions were nested with the same five optima identified in the more conservative model and in the liberal 12 optima model. These optima correspond to arboreal taxa at the base of the tree (R1: *Hypsilurus*, *Lophosaurus*, *Chelosania*), two terrestrial clades (R2: *Ctenophorus*; R5: *Tympanocryptis*), an Australian semiarboreal group (R3: *Amphibolurus*, *Gowidon*, *Intellagama*, *Tropicagama*, *Lophognathus*, *et al.*), and the unique *Moloch* (R4) (Fig.3; Fig.S9). More liberal models added optima which largely corresponded to genera (e.g. *Diporiphora*, *Pogona* + *Rankinia*) and outlier species (*Chlamydosaurus kingii*, *Cryptagama aurita*, *Diporiphora superba*) and optimally identified 12 peaks (Fig.S10). These clusters were largely visible when looking at raw morphological distances among taxa (Fig.S11).

In both optima scenarios ($k = 5$, $k = 12$), RandomForest predictions suggested an arboreal-type ancestor for the Amphibolurinae with some uncertainty ($k = 5$, 66% assignment to arboreal; $k = 12$, 58%). The core Australian amphibolurine ancestor (most recent common ancestor [MRCA] of *Intellagama* and relatives) is estimated as regime R3 (semiarboreal generalist), however, its immediate ancestor is estimated as regime R1 (arboreal). The amphibolurine MRCA is estimated to have trait values most similar to *Chlamydosaurus*, *Intellagama*, and *Physignathus* (Fig.S11,S12). Simulations show our approach has reasonable accuracy (80%) and no identifiable bias in assigning ancestral morphologies to contemporary regimes, though accuracy varied across nodes and declined with increasing node age.

Discussion

Organismal forms in nature are not distributed evenly across the possible phenotypic space. Instead, observed phenotypes often cluster in discrete regions of morphospace, a result of genetic, functional, and/or developmental constraints (Deline et al. 2018). These clumps of morphologically similar organisms identify peaks in the adaptive landscape. Through time, the adaptive landscape may change as conditions vary and species go extinct, persist, and proliferate. The invasion of a new geographic region provides us the opportunity to observe how the adaptive landscape evolves, and to identify the tempo and mode with which new regions of the landscape become populated. Amphibolurinae dragon lizards are an ideal system to evaluate this because of their diversity of forms, age, and history. Our findings suggest that dispersal of dragon lizards into Australia facilitated a marked expansion in the morphological landscape. This exploration was driven first by a transition from an arboreal to a generalist ancestor and later into specialized ecomorphologies. Major morphological transitions were aided by focused evolution resulting in the rapid change of traits.

Phylogenomics of the Amphibolurinae

Dragon lizards (Agamidae) are a charismatic and diverse group of ~600 species found across Africa, Asia, and Oceania. In Australia and New Guinea they reach ecomorphological extremes that include burrowing species that live on salt-crusted desert lake-beds and arboreal species in tropical rainforests. Our new sequence-capture dataset extends our phylogenetic understanding of the group (Macey et al. 2000; Melville et al. 2001, Hugall et al. 2008; Shoo et al. 2008; Melville et al. 2018; Chaplin et al. 2020; Fenker et al. 2024) and explores their morphological diversification.

Phylogenetic relationships among agamid subfamilies have been estimated previously (e.g. Streicher et al. 2016; Burbrink et al. 2020; Welt & Raxworthy 2022), but only recently have they been discussed extensively (Scarpetta et al. 2025). Agamids likely originated in Asia, and their arrival in Australo-Papua mirrors biogeographic patterns in other Australian squamate groups (Keogh 1998; Esquerre et al. 2020; Brennan et al. 2021). While the Amphibolurinae as a group date back to the Cretaceous, their living diversity is less than half that age. The split between Asian (*Physignathus*) and Australo-Papuan species occurred around the Eocene/Oligocene transition and was closely followed by rapid diversification over the next ten million years. This is consistent with paleontological evidence from Australia which suggests fossils assignable to *Intellagama* are relatively common in the early-to-middle Miocene deposits of the Riversleigh World Heritage Area (Covacevich et al. 1990). Subsequent

diversification occurred rapidly following the divergence of Australian species from their forest-dwelling relatives in New Guinea and resulted in frequent splits over just a few hundred thousand years. This is coincident with a period of dramatic ecomorphological change. Through the late Miocene, diversification slowed and was dominated by three major groups (*Ctenophorus*, *Diporiphora*, *Tympanocryptis*), a pattern that is consistent with theory around adaptive radiations and dispersal into new regions (Harmon et al. 2010; Schluter 2000; Simpson 1944; Yoder et al. 2010).

Mapping Morphology to Ecology

The adaptive landscape of amphibolurine dragons is dispersed and patchy, with peaks that largely correspond to recognized genera. Obvious phenotypic peaks relate to strongly arboreal (*Hypsilurus*, *Lophosaurus*) and terrestrial species (*Tympanocryptis*, *Ctenophorus*). However, intermediate forms that use a variety of substrate types (*Amphibolurus*, *Gowidon*, *Lophognathus*, *Tropicagama*, et al.) fill the morphological space, uniting disparate peaks of the landscape and facilitating transitions to new ecologies. These taxa with broader niche breadths are either lumped under a single broad adaptive peak (when $k=5$ optima; Fig.S9) or two peaks (when $k=12$ optima; Fig.S10). This is important because these intermediate morphologies appear common (~50% of species richness), are distributed across the entire continent in all habitat types, and provide a theoretical link between more extreme morphological forms. Modelling niche breadth supports this by preferring pathways of narrowing substrate usage and favoring transitions between specialist and generalist states over transitions from one specialist to another (Fig.2). The idea of generalists as a bridge between morphotypes is not exclusively a reflection of how we generate morphospaces. In dragon lizards this process is informed by the branching pattern of the phylogeny. Specialist forms often descend from morphologically intermediate ancestors, and in some cases are nested deeply within generalist clades. For example the stout rock-dwelling *Diporiphora bennetti* and gracile hummock grass living *Diporiphora adductus* likely both evolved from a more typical amphibolurine morphology seen in their closest relatives.

Linking morphology to ecology however, is not always so clear. This may be the result of a many-to-one mapping where varied morphologies enable the same ecology (Wainwright et al. 2005; Moen 2019). For example, taxa such as *Lophosaurus*, *Chelosania*, and *Diporiphora superba* are all obligately arboreal and highly specialized dragons, however, these taxa map to either two or three discrete adaptive zones (Fig.3) and use arboreal spaces differently. In this situation, convergence on an arboreal lifestyle is met with only incomplete morphological similarity (Grossnickle et al. 2024). This may be due to a number of reasons including

phylogenetic inertia resisting selection (Blomberg & Garland 2002). This idea is relevant when considering the ecology of the ancestral amphibolurine. Our analyses suggest this ancestor was likely morphologically similar to the arboreal *Lophosaurus* and *Hypsilurus*, but also to *Physignathus* and *Intellagama*—two species that are considered semiaquatic (Fig.S12). These last two species are similar enough that they were considered congeneric for over 150 years and are often called “water dragons”. Despite their preferred habitat, they fit most morphological descriptors of an arboreal dragon. We flag this to note the imperfect mapping of morphology to ecology and uncertainty in estimating ancestral states.

The Evolving Adaptive Landscape

The transition from wet closed-canopy rainforests in Asia and New Guinea towards new and different environments in Australia presents a prime case for ecological opportunity (Collar et al. 2010; Gray et al. 2019a). Among the newly available habitats, dragons encountered sparse open forests, grasslands, sandy and stony deserts, and rocky escarpments, which are all but absent across Malesia. By expanding their geographical range, dragon lizards also extended the adaptive landscape to enable new forms that took advantage of these habitats (Yoder et al. 2010). This extension provided opportunity for new peaks located in distant regions of morphospace. To traverse that space however, required deterministic trends in trait evolution that suggest selection on individual or suites of traits (Figs.3,4). In dragon lizards, this may have first resulted in a transition from an arboreal form to a generalist form (Fig.2). Our analyses indicate that this occurred rapidly and dramatically, resulting in a multivariate shape change dozens of times greater than expected (Fig.4 highlight 5). The resulting form likely had shorter limbs and tail, a wider body, and shorter snout as it spent less time in the trees (Fig.5). From generalist forms amphibolurines were able to further explore morphological extremes. This includes repeated transitions towards small-bodied terrestrial forms (*Ctenophorus aurita*, *Rankinia*, *Tympanocryptis*), and moves back towards elongate arboreal forms (*Diporiphora superba*, *Chlamydosaurus*) (Fig.5). In this way, generalists negate the need to cross wide fitness valleys in the adaptive landscape. Instead, species can explore across ridges or broad fitness peaks to reach distant morphospaces. This provides more evidence that transitions across the landscape from specialist to specialist are uncommon and likely require an intermediate form (Fig.2) (Dennis et al. 2011).

In our investigation across the dragon lizard body plan, we identify varied modes of evolution among and within traits. The evolution of dragon morphology involved both a gradual diffusion process away from stable peaks and infrequent shifts in the evolutionary rate or trait mean. Interestingly, to explain the observed morphological landscape, fewer than 10% of branches


need multivariate trait change that exceeds Brownian expectations. **This is visible as discrete clumps across the morphospace.** These high density areas indicate slow evolutionary rates or constrained evolution (i.e. Ornstein Uhlenbeck), as has been previously identified in *Diporiphora* (Smith et al. 2010). Stable peaks populated by many closely-related species highlight diffusion as the most frequent pathway for change among species. In comparison, rare evolutionary shifts are concentrated around changes among adaptive zones. These shifts are nested and temporally staggered resulting in a dynamic adaptive landscape that does not reflect the archetypical tempo or mode of an adaptive radiation. This complex mosaic of evolutionary modes helps to reconcile Simpson's idea of quantum evolutionary change (few, extreme changes) with Darwin's ideas of gradualism (frequent, small changes) (Simpson 1944; Charlesworth et al. 1982).

The morphological evolution of dragon lizards has been a heterogeneous process, varying temporally, phylogenetically, and across traits. Broadly, dragons have gone through periods of early (30-25 my) and late expansion (5-0 my) in the morphological hypervolume, bookending a period of little innovation during much of the Miocene (Fig.S8). **Admittedly, this pattern lacks information provided by extinct species. However, if our expectation that ancestrally intermediate morphologies yield young derived morphologies is true, then we would not expect extinction to considerably affect our conclusions about the accumulation of morphological volume. This does establish further questions about how biased extinction of morphologically extreme species may drive the pattern that we see.** Looking across all species, we identify that the primary axes for variation are through change in absolute size and tail length (Fig.S7). This is immediately apparent when comparing large tree dragons (*Hypsilurus magnus*: SVL >200 mm, tail length >600 mm) to pebble-mimicking dragons (*Ctenophorus aurita*: SVL <40 mm, tail length <35 mm; *Tympanocryptis cephalus* group: SVL <60 mm, tail length <90 mm). Elaboration in size and tail length beyond other traits is a pattern seen in other squamate clades (Brennan et al. 2024; Meiri 2010), but dramatic changes are not limited to these axes. Instead, amphibolurines show exceptional variation in many traits and greater than expected diversity relative to other iguanians (Gray et al. 2019a; Gray et al. 2019b). Most trait trends appear associated with habitat type (Thompson & Withers 2005; Collar et al. 2010), e.g. wide bodies in terrestrial groups, but others are harder to quantify and likely associated with defense or communication, like the frill of *Chlamydosaurus*. Harder yet to explain is the novelty of the enigmatic thorny devil, *Moloch*. They likely pre-date the Australian deserts in which they wander (Pepper & Keogh, 2021). They exclusively eat ants. And they have undergone such a profound evolutionary modification that they are nearly unrecognizable as agamids. In expanding into this new adaptive zone, the thorny devil has stretched the morphological space.

Trends towards increasing specialization in agamids show how the adaptive landscape itself can change and grow, and how species with broad niches can act as bridges to stretch borders and alter the landscape topography in response to new opportunities.

Acknowledgements

This project represents an output of the Australian Amphibian and Reptile Genomics initiative (AusARG), generously funded by BioPlatforms Australia. We appreciate the provision of computing and data resources provided by the Australian BioCommons Leadership Share (ABLeS) program and Seqera Tower service. These programs are co-funded by Bioplatforms Australia (enabled by NCRIS), the National Computational Infrastructure and Pawsey Supercomputing Centre. Special thanks to Sophie Mazard and Ziad al Bkhetan. We thank our many Australian partner institutions (Australian Museum, Museum and Art Gallery of the Northern Territory, Queensland Museum, Museums Victoria, South Australian Museum, Western Australian Museum) and their associated curators and collections managers, who made this work possible through generous tissue loans and collections access. We declare no conflicts of interest.



494
495
496

Figure 1. Species-level timetree of the Amphibolurinae, color-coded by genus. The ASTRAL species tree was time-calibrated with MCMCTree. A yellow band highlights a series of rapid splits early in the history of this group. (A) Relationships among agamid subfamilies and color-coded dragon drawings at right (not to scale) highlight some of the morphological diversity of this group. (B) The spread of species in two-dimensional morphospace (also colored by genus) and primary axes of variation. *Diporiphora* (dark purple) and *Tympanocryptis* (light purple) highlight dense clusters of morphospace, contrasting with morphological outliers indicated by squares in the morphospace and on the tree (*D. superba*, *M. horridus*, *C. aurita*). The estimated root morphotype is indicated by a triangle. White circles on the tree denote local posterior probabilities <0.95 . (C) The evolution of the morphological hypervolume through time (orange) relative to null expectations (grey) shows periods of rapid expansion (25-15; 5-0 mya) and a period of relative stasis (15-5 mya).




Figure 2. The evolution of niche breadth favors transitions between niche generalists (high breadth) and specialists (low breadth). The same phylogeny is plotted mirrored (top and bottom) with preferred results of model fitting exercises. Top, niche breadth is characterized as the sum of substrates used by each taxon (darker colors, fewer substrates; lighter colors, more substrates) and indicates a stepping-stone pattern of substrate gain and loss, rather than rapid jumps towards specialization or generalism. Transition rates favor narrowing niche breadth, Bottom, transitions are favored between broad and narrow niche breadths, rather than between narrow niches (specialists). Outlined pie charts in both plots show that in both reconstructions the common ancestor of Amphibolurine likely had a narrow niche breadth (two substrates, or tree specialist), but niche breadth increased at the common ancestor of the Australian clade.





Figure 3. The adaptive morphological zones of amphibolurine dragons highlight a diversity of forms. Colored regimes were identified under a multi-optima OU model in *PhyloEM* and applied to ancestral taxa using *randomForests*. The tree shows the preferred 5 regime model estimated from all 19 size corrected morphological traits and likelihoods for ancestors as estimated by *randomForest*. Two regimes (blue, orange) map to obligately terrestrial groups. The red regime highlights primarily arboreal species and the morphologically similar *Physignathus*. Yellow indicates a diverse clade of primarily generalist species. The amphibolurine most recent common ancestor (MRCA) is estimated as a *Hypsilurus*/*Lophosaurus*-like arboreal dragon and the Australian MRCA is estimated as a semiarboreal generalist lizard (both indicated with black outlined pie). Illustrated species are (A) *Moloch horridus*, (B) *Diporiphora superba*, (C) *Chlamydosaurus kingii*, (D) *Ctenophorus aurita*, (E) *Rankinia diemensis*, (F) *Tympanocryptis pseudopsephos*, (G) *Ctenophorus reticulatus*, (H) *Hypsilurus modestus*, (I) *Chelosania brunnea*, (J) *Gowidon longirostris*, (K) *Lophosaurus boydii*.



① *Physignathus* ② *Moloch* ③ *C. aurita* ④ *Tympanocryptis* ⑤ MRCA Australia ⑥ *Ctenophorus*

Figure 4. Major shifts in morphologies occur as directional evolutionary changes that are greater than expected. Left, morphological innovation across the dragon tree, with warm colors indicating greater multivariate Euclidean distance from the amphibolurine most recent common ancestor, and cooler colors indicating greater similarity. Grey shading in outer circles delineates genera. Right, the position and strength of directional changes that exceed multivariate shape change under the null evolutionary model. Light grey branches follow expectations under the null Brownian model, and colored branches indicate greater than expected change. Circle number 5 indicates the branch leading to the Australian radiation of dragon lizards and denotes a 26-fold increase over the expected amount of morphological change. This very short branch accounts for a paradigm shift in the evolution of amphibolurine lizards away from an arboreal ancestor towards a more generalist morphology.




Figure 5. Movement among optima occurs via changes in different suites of traits. Plots in the left column visualize the shift from the immediately ancestral optimum (grey shading) to a new optimum (warm shading) represented in two dimensional space. Lizards at center illustrate the morphological extremes of new optima. Plots at right indicate how individual traits changed in evolving from the previous optima to current. Grey bars highlight the two most variable traits: size (a composite of the geometric mean of all trait measures) and tail length.

References

- Arnold, S. J., Pfrender, M. E., & Jones, A. G. (2001). The adaptive landscape as a conceptual bridge between micro-and macroevolution. *Genetica*, 112, 9-32.
- Bastide, P., Ané, C., Robin, S., & Mariadassou, M. (2018). Inference of adaptive shifts for multivariate correlated traits. *Systematic Biology*, 67(4), 662-680.
- Blomberg, S. P., & Garland Jr, T. (2002). Tempo and mode in evolution: phylogenetic inertia, adaptation and comparative methods. *Journal of Evolutionary Biology*, 15(6), 899-910.
- Blonder, B. (2016). Do hypervolumes have holes?. *The American Naturalist*, 187(4), E93-E105.
- Blonder, B., Morrow, C. B., Maitner, B., Harris, D. J., Lamanna, C., Violle, C., Enquist, B. J., & Kerkhoff, A. J. (2018). New approaches for delineating n-dimensional hypervolumes. *Methods in Ecology and Evolution*, 9(2), 305-319.
- Borowiec, M. L. (2016). AMAS: a fast tool for alignment manipulation and computing of summary statistics. *PeerJ*, 4, e1660.
- Brennan, I. G., Lemmon, A. R., Lemmon, E. M., Portik, D. M., Weijola, V., Welton, L., ... & Keogh, J. S. (2021). Phylogenomics of monitor lizards and the role of competition in dictating body size disparity. *Systematic Biology*, 70(1), 120-132.
- Brennan, I. G., Singhal, S., & Al Bkhetan, Z. (2024). *pipesnake*: generalized software for the assembly and analysis of phylogenomic datasets from conserved genomic loci. *Bioinformatics*, 40(5), btae195.
- Brennan, I. G., Chapple, D. G., Keogh, J. S., & Donnellan, S. (2024). Evolutionary bursts drive morphological novelty in the world's largest skinks. *Current Biology*, 34(17), 3905-3916.
- Bruton M., Melville J., Sumner J., Porter J., Mahony S. & Doughty P. (2025). The last piece of the puzzle: Detection of an extremely rare lizard in the Kimberley completes the phylogeny of Australian dragons (Agamidae). *Australian Journal of Taxonomy* 93: 1–15. doi: <https://doi.org/10.54102/ajt.zoa0>
- Burbrink, F. T., Grazziotin, F. G., Pyron, R. A., Cundall, D., Donnellan, S., Irish, F., ... & Zaher, H. (2020). Interrogating genomic-scale data for Squamata (lizards, snakes, and amphisbaenians) shows no support for key traditional morphological relationships. *Systematic biology*, 69(3), 502-520.
- Burin, G., Park, T., James, T. D., Slater, G. J., & Cooper, N. (2023). The dynamic adaptive landscape of cetacean body size. *Current Biology*, 33(9), 1787-1794.
- Bushnell, B. (2014). BBMap: A Fast, Accurate, Splice-Aware Aligner. Berkeley, CA: Lawrence Berkeley National Lab.
- Chaplin, K., Sumner, J., Hipsley, C. A., & Melville, J. (2020). An integrative approach using phylogenomics and high-resolution X-ray computed tomography for species delimitation in cryptic taxa. *Systematic Biology*, 69(2), 294-307.
- Charlesworth, B., Lande, R., & Slatkin, M. (1982). A neo-Darwinian commentary on macroevolution. *Evolution*, 36(3), 474-498.

- Collar, D. C., Schulte, J. A., O'meara, B. C., & Losos, J. B. (2010). Habitat use affects morphological diversification in dragon lizards. *Journal of Evolutionary Biology*, 23(5), 1033-1049.
- Covacevich, J., Couper, P., Molnar, R. E., Witten, G., & Young, W. (1990). Miocene dragons from Riversleigh: new data on the history of the family Agamidae (Reptilia: Squamata) in Australia. *Memoirs of the Queensland Museum*, 29(2), 339-360.
- Deline, B., Greenwood, J. M., Clark, J. W., Puttick, M. N., Peterson, K. J., & Donoghue, P. C. (2018). Evolution of metazoan morphological disparity. *Proceedings of the National Academy of Sciences*, 115(38), E8909-E8918.
- Dennis, R. L., Dapporto, L., Fattorini, S., & Cook, L. M. (2011). The generalism–specialism debate: the role of generalists in the life and death of species. *Biological Journal of the Linnean Society*, 104(4), 725-737.
- Devictor, V., Clavel, J., Julliard, R., Lavergne, S., Mouillot, D., Thuiller, W., Venail, P., Villegger, S. and Mouquet, N., (2010). Defining and measuring ecological specialization. *Journal of Applied Ecology*, 47(1), pp.15-25.
- dos Reis, M. and Yang, Z. (2011). Approximate likelihood calculation for Bayesian estimation of divergence times. *Molecular Biology and Evolution*, 28:2161-2172.
- Esquerré, D., Donnellan, S., Brennan, I. G., Lemmon, A. R., Moriarty Lemmon, E., Zaher, H., ... & Keogh, J. S. (2020). Phylogenomics, biogeography, and morphometrics reveal rapid phenotypic evolution in pythons after crossing Wallace's line. *Systematic Biology*, 69(6), 1039-1051.
- Fenker, J., Melville, J., & Moritz, C. (2024). Dragons in the tropics–Phylogeography and speciation in Diporiphora lizards and common geographic breaks in co-distributed taxa. *Molecular Phylogenetics and Evolution*, 197, 108090.
- Futuyma, D. J., & Moreno, G. (1988). The evolution of ecological specialization. *Annual review of Ecology and Systematics*, 207-233.
- Gray, J. A., Sherratt, E., Hutchinson, M. N., & Jones, M. E. (2019a). Evolution of cranial shape in a continental-scale evolutionary radiation of Australian lizards. *Evolution*, 73(11), 2216-2229.
- Gray, J. A., Hutchinson, M. N., & Jones, M. E. (2019b). Exceptional disparity in Australian agamid lizards is a possible result of arrival into vacant niche. *The Anatomical Record*, 302(9), 1536-1543.
- Grossnickle, D.M., Brightly, W.H., Weaver, L.N., Stanchak, K.E., Roston, R.A., Pevsner, S.K., Stayton, C.T., Polly, P.D. and Law, C.J., (2024). Challenges and advances in measuring phenotypic convergence. *Evolution*, 78(8), pp.1355-1371.
- Harmon, L. J., Kolbe, J. J., Cheverud, J. M., & Losos, J. B. (2005). Convergence and the multidimensional niche. *Evolution*, 59(2), 409-421.
- Harmon, L. J., Losos, J. B., Jonathan Davies, T., Gillespie, R. G., Gittleman, J. L., Bryan Jennings, W., Kozak, K.H., McPeck, M.A., Moreno-Roark, F., Near, T.J., Purvis, A., Ricklefs, R.E., Schluter, D., Schulte III, J.A., Seehausen, O., Sidlauskas, B.L., Torres-Carvajal, O., Weir, J.T., & Mooers, A. Ø. (2010). Early bursts of body size and shape evolution are rare in comparative data. *Evolution*, 64(8), 2385-2396.
- Hugall, A. F., Foster, R., Hutchinson, M., & Lee, M. S. (2008). Phylogeny of Australasian agamid lizards based on nuclear and mitochondrial genes: implications for

- morphological evolution and biogeography. *Biological Journal of the Linnean Society*, 93(2), 343-358.
- Hutchinson, G.E. (1957). Concluding Remarks. Cold Spring Harbor Symposia on Quantitative Biology, 22, 415-427.
- Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermini, L. S. (2017). ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature methods*, 14(6), 587-589.
- Keogh, J. S. (1998). Molecular phylogeny of elapid snakes and a consideration of their biogeographic history. *Biological Journal of the Linnean Society*, 63(2), 177-203.
- Liaw, A., & Wiener, M. (2002). Classification and regression by randomForest. *R news*, 2(3), 18-22.
- Marshall, C. R. (2014). The evolution of morphogenetic fitness landscapes: conceptualising the interplay between the developmental and ecological drivers of morphological innovation. *Australian Journal of Zoology*, 62(1), 3-17.
- Macey, J. Robert, James A. Schulte, Allan Larson, Natalia B. Ananjeva, Yuezhao Wang, Rohan Pethiyagoda, Nasrullah Rastegar-Pouyani, and Theodore J. Papenfuss. Evaluating trans-Tethys migration: an example using acrodont lizard phylogenetics. *Systematic Biology* 49, no. 2 (2000): 233-256.
- Meiri, S. (2010). Length–weight allometries in lizards. *Journal of Zoology*, 281(3), 218-226.
- Melville, J., Schulte, J. A., & Larson, A. (2001). A molecular phylogenetic study of ecological diversification in the Australian lizard genus *Ctenophorus*. *Journal of Experimental Zoology*, 291(4), 339-353.
- Melville, J., Ritchie, E. G., Chapple, S. N., Glor, R. E., & Schulte II, J. A. (2018). Diversity in Australia's tropical savannas: An integrative taxonomic revision of agamid lizards from the genera *Amphibolurus* and *Lophognathus* (Lacertilia: Agamidae). *Memoirs of Museum Victoria*, 77, 41-61.
- Melville, J., Date, K. S., Horner, P., & Doughty, P. (2019). Taxonomic revision of dragon lizards in the genus *Diporiphora* (Reptilia: Agamidae) from the Australian monsoonal tropics. *Memoirs of Museum Victoria*, 78, 23-55.
- Melville, J., & Wilson, S. K. (2019). Dragon Lizards of Australia: Evolution, ecology and a comprehensive field guide. Museums Victoria Publishing.
- Minh, B. Q., Nguyen, M. A. T., & von Haeseler, A. (2013). Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution*, 30(5), 1188-1195.
- Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution*, 37(5), 1530-1534.
- Moen, D. S. (2019). What determines the distinct morphology of species with a particular ecology? The roles of many-to-one mapping and trade-offs in the evolution of frog ecomorphology and performance. *The American Naturalist*, 194(4), E81-E95.
- Pagel, M., O'Donovan, C., & Meade, A. (2022). General statistical model shows that macroevolutionary patterns and processes are consistent with Darwinian gradualism. *Nature communications*, 13(1), 1113.
- Pavanetto N, Puglielli G (2024). *funspace*: Creating and Representing Functional Trait Spaces. R package version 0.2.2, <https://CRAN.R-project.org/package=funspace>.

- Pepper, M., & Keogh, J. S. (2021). Life in the “dead heart” of Australia: The geohistory of the Australian deserts and its impact on genetic diversity of arid zone lizards. *Journal of Biogeography*, 48(4), 716-746.
- Pennell, M. W., Eastman, J. M., Slater, G. J., Brown, J. W., Uyeda, J. C., FitzJohn, R. G., ... & Harmon, L. J. (2014). geiger v2. 0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. *Bioinformatics*, 30(15), 2216-2218.
- Poisot, T., Bever, J. D., Nemri, A., Thrall, P. H., & Hochberg, M. E. (2011). A conceptual framework for the evolution of ecological specialisation. *Ecology letters*, 14(9), 841-851.
- R Core Team. (2024). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. www.R-project.org
- Rannala, B., Yang, Z. (2007) Inferring speciation times under an episodic molecular clock. *Systematic Biology*, 56:453-466.
- Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and other things). *PeerJ*, 12, e16505.
- Scarpetta, S. G., Karin, B. R., Corl, A., & McGuire, J. A. (2025). The Evolutionary History of Dragon Lizards (Squamata: Agamidae) Revealed by Phylogenomics. *Systematic Biology*, syaf088.
- Schluter, D. (2000). The ecology of adaptive radiation. OUP Oxford.
- Shoo, L. P., Rose, R., Doughty, P., Austin, J. J., & Melville, J. (2008). Diversification patterns of pebble-mimic dragons are consistent with historical disruption of important habitat corridors in arid Australia. *Molecular Phylogenetics and Evolution*, 48(2), 528-542.
- Simpson, G. G. (1944). Tempo and mode in evolution. Columbia University Press.
- Singhal, S., Grundler, M., Colli, G., & Rabosky, D. L. (2017). Squamate conserved loci (SqCL): a unified set of conserved loci for phylogenomics and population genetics of squamate reptiles. *Molecular ecology resources*, 17(6), e12-e24.
- Smith, K. L., Harmon, L. J., Shoo, L. P., & Melville, J. (2011). Evidence of constrained phenotypic evolution in a cryptic species complex of agamid lizards. *Evolution*, 65(4), 976-992.
- Streicher, J. W., Schulte, J. A., & Wiens, J. J. (2016). How should genes and taxa be sampled for phylogenomic analyses with missing data? An empirical study in iguanian lizards. *Systematic biology*, 65(1), 128-145.
- Thompson, G., & Withers, P. (2005). Shape of western Australian dragon lizards (Agamidae). *Amphibia-Reptilia*, 26(1), 73-85.
- Tiatragul, S., Brennan, I. G., Broady, E. S., & Keogh, J. S. (2023). Australia’s hidden radiation: Phylogenomics analysis reveals rapid Miocene radiation of blindsnakes. *Molecular Phylogenetics and Evolution*, 185, 107812.
- Title, P. O., Singhal, S., Grundler, M. C., Costa, G. C., Pyron, R. A., Colston, T. J., ... & Rabosky, D. L. (2024). The macroevolutionary singularity of snakes. *Science*, 383(6685), 918-92.
- Welt, R. S., & Raxworthy, C. J. (2022). Dispersal, not vicariance, explains the biogeographic origin of iguanas on Madagascar. *Molecular Phylogenetics and Evolution*, 167, 107345.

735 Wainwright, P. C., Alfaro, M. E., Bolnick, D. I., & Hulsey, C. D. (2005). Many-to-one
736 mapping of form to function: a general principle in organismal design?. *Integrative*
737 *and Comparative Biology*, 45(2), 256-262.

738 Wright, S. (1932). The roles of mutation, inbreeding, crossbreeding and selection in
739 evolution. In: Proceedings of the sixth international congress of genetics, pp 356–366.

740 Yoder, J. B., Clancey, E., Des Roches, S., Eastman, J. M., Gentry, L., Godsoe, W., ... &
741 Harmon, L. J. (2010). Ecological opportunity and the origin of adaptive radiations.
742 *Journal of Evolutionary Biology*, 23(8), 1581-1596.

743 Zhang, C., & Mirarab, S. (2022). Weighting by gene tree uncertainty improves accuracy of
744 quartet-based species trees. *Molecular Biology and Evolution*, 39(12), msac215