

# Integrating macroecology into macroevolutionary models: Evolution in Australian reptiles

*NSF Postdoc Outline*

*October 17, 2018*

## Contents

<b>Project Summary (1pg max)</b>	<b>2</b>
Overview (include sponsoring scientists and institutions): . . . . .	2
Intellectual Merit . . . . .	2
Broader Impacts . . . . .	2
<b>Project Description (6pg max)</b>	<b>3</b>
Introduction . . . . .	3
Research Plan . . . . .	3
Training Objectives . . . . .	5
Career Development . . . . .	5
Sponsoring Scientist and Institution Justification . . . . .	5
Yearly Timetable (goals, outcomes) . . . . .	5
Broader Impacts Narrative . . . . .	5
<b>References</b>	<b>5</b>

*NSF Postdoctoral Research Fellowships in Biology Solicitation*

## Project Summary (1pg max)

### Overview (include sponsoring scientists and institutions):

Ecosystems are defined by both the organisms inhabiting them, and the interactions among these organisms. Quantifying the influence of these interactions on the evolution of the participants should therefore be a priority in evolutionary biology, but remains difficult. Instead, common comparative methods account for phylogeny, but ignore how cohabiting lineages influence one another phenotypically and spatially. My research goal is to extend our existing toolkit to include methods which more accurately account for macroecology in macroevolution. This includes incorporating (1) inter-lineage interactions, (2) intra-lineage variation, and (3) historical biogeography, to better explain observed patterns in trait diversity.

### Intellectual Merit

Organismal interactions provide an important selective force for evolution. Predator-prey, plant-pollinator, and host-parasite relationships form the basis for ecosystems, and drive the accumulation and distribution of diversity. For a while now, both community ecologists and comparative evolutionary biologists have sought to understand how this diversity is built on local and phylogenetic scales. However, they have approached these questions from very different angles. Community ecology has been largely agnostic of evolution along phylogenies, and evolutionary biologists have relied on ecology naïve comparative methods. Now, we are on the edge of being able to appropriately incorporate community interactions into phylogenetic comparative methods of macroevolution. My specific aims are to improve our ability to identify the influence of interlineage interactions on community assembly and trait evolution, and better estimate the strength of these interactions. By accounting for intralinear trait variation and biogeographic histories, we can make these methods more biologically meaningful. This research intends to blend together the fields of macroecology and macroevolution, and in the process, bring together ecologists with evolutionary biologists, and museum curators with computer scientists.

### Broader Impacts

For this project, I will create a number of resources for both the macroecological/macroevolutionary and general science communities. This includes curating an open-source ecological database of Australian reptiles (morphology, dietary, distributional data), designing new methods for modelling trait evolution in a phylogenetic community framework, and communicating my results through scientific publications and popular science engagements. My computational and methodological mentor Dr. Brian O'Meara (*University of Tennessee, Knoxville*) is a world expert in comparative methods, bringing together mathematical advances with empirical evolutionary studies. My curatorial and biological mentor Dr. Mike Lee (*South Australian Museum*) has been actively researching macroevolution and the complex origins of reptiles for over two decades. Dr. Lee works to actively incorporate the often disparate fields of morphological and molecular evolution. Together, these sponsors (and their institutions) provide the opportunity to expand my theoretical understanding of evolutionary biology, and the computational and methodological abilities to apply it. In turn, I aim to enrich the experience of others by collaborating with domestic and international colleagues, including undergraduate and postgraduate students. I have also outlined a plans for a workshop on phylogenetic comparative methods (see appendix). Finally, I enjoy outreach through social media, so will continue to use my position in the community to broadcast results in an entertaining and digestible “PopSci” manner. *workshop development*

# Project Description (6pg max)

## Introduction

Organismal phenotypes are the result of genetic *and* environmental forces, and the interaction between them. When explaining patterns of phenotypic evolution, we can account for some genetic influence by incorporating a phylogeny and drift, recognizing that phenotypes tend to be more similar between closely related species. Descent with modification thus provides us with some expectations as to how phenotypes should evolve. But how do we account for influence of extrinsic biotic factors on observed traits? In response to these questions, and in an attempt to explain the accumulation of organismal diversity, the field of phylogenetic comparative methods was developed. Unfortunately, nearly all commonly used models for describing phenotypic evolution are ecologically neutral, meaning they fail to account for species interactions with one another, and their environment. This results in lifeless models that mathematically describe the *pattern* of phenotypic evolution, but lack a biological explanation of the *process*. **My overarching research goal is to extend our existing toolkit to include a more diverse set of models which account for macroecology in macroevolution.** This includes incorporating inter-lineage interactions, intra-lineage variation, and historical biogeography, to better explain observed patterns in trait diversity. To develop these methods, I intend to use Australian reptiles as an empirical example.

**But, why Australian reptiles?** As an island continent, Australia is a natural laboratory for studies of evolution. With more than 1,000 species of squamate reptiles (lizards and snakes), extensive museum collections, and curated occurrence records, Australia is a prime opportunity for investigating macroecological models of phenotypic evolution. Since parting ways with Antarctica, the Australian continent has been isolated for around 35 million years. This means that with exception a single preexisting Gondwanan clade (Pygopodoid geckos), all other Australian reptile groups have arrived from Asia and radiated in situ. This includes iconic radiations of pythons, monitor lizards, elapid snakes, skinks, agamid lizards, and blind snakes.

## Research Plan

### Research Objectives

The overarching goal of this project is to better account for macroecological processes in macroevolutionary methods. This includes designing new tools *and* implementing them to tackle empirical evolutionary questions. We can classify the broad aims of this study into more specific questions:

1. Does including interspecific interactions improve our inferences of phenotypic evolution?
2. How do interspecific interactions affect the spatial evolution and distribution of species?
3. Can we build an extensive ecological dataset of Australian reptiles on a continental scale?
4. Is competitive signal identifiable from phenotypic and spatial data of Australian reptiles?

## Methods

This research depends directly on the development of novel methods and the collection and curation of a species-level ecological database (of Australian reptiles). The ultimate goal is to integrate these two aspects, to directly address our questions of interlineage interactions using empirical data. We expect that the approaches used to collect the data and develop the comparative methods will also be of direct interest for those hoping to follow a similar methodology.

### **Objective 1: Does including interspecific interactions improve our inferences of phenotypic evolution?**

Interspecific interactions act as a strong selective force on organismal phenotypes. Interactions may be competitive (vying for resources), mutualistic (beneficial to both parties), commensal, parasitic, or predatory, and dictate a variety of dynamic responses. This has led to theories of character displacement, competitive exclusion, and phenotypic matching, among others. In a classic anecdote, Darwin and Wallace each considered a Madagascan orchid with a foot long nectary. They then independently determined there must be a moth pollinator with an equally long proboscis to match. The later discovery of just such a moth was ultimately less of a surprise, but fulfilled this expectation that organismal interactions can dictate phenotypic evolution, even to absurd extremes.

### **Objective 2: How do interspecific interactions affect the spatial evolution and distribution of species?**

The distribution of species across a landscape can be dictated by both intrinsic and extrinsic influences. In maybe the most obvious examples, habitats may “filter” for species with similar ecologies, or instead, competition might result in exclusion from a given ecosystem.

### **Objective 3: Can we build an extensive ecological dataset of Australian reptiles on a continental scale?**

Existing eco-morphological databases cover much of the Australian vertebrate fauna, including mammals (Pantheria) and birds (Garnett et al., 2014; EltonTraits), but reptile taxa remain poorly represented (Meiri, 2018). This is surprising given the number of studies of Australian reptile phylogenetics, ecology, and morphology, and remarkable museum collections. In association with the **OpenTraits** initiative, I intend to collate existing, undatabased ecological and morphological information with newly collected data, to span the entirety of Australian reptile taxa (>1000 species). In a departure from traditional databasing methods, I hope to include measurements, observations, and metadata for all viewed specimens. My hope is that instead of summarizing this information to species means, I can contribute a database which is more flexible to taxonomic changes and better describes intraspecific variation. As a current research associate of the Western Australian Museum (130,000+ reptile records) and a proposed associate of the South Australian Museum (55,000+ reptile records), I would have access to sampling for the entirety of the continent, making this type of data collection feasible. A resource of this kind would be a world-first, providing invaluable information to future researchers, and inalienable data to museums about their own collections.

### **Objective 4: Is competitive signal identifiable from phenotypic and spatial data of Australian reptiles?**

This question represents the synthesis of the three previous objectives. Broadly, my intention is to develop methods that may be of interest to other researchers studying macroevolution and macroecology. However, it is also important to ask these questions empirically, employing existing biological collections. Australian reptiles are particularly well suited for these questions because they represent a number of species rich, eco-morphologically diverse, endemic radiations. Building on phylogenomic research from my PhD and others, we now have strong species-level time-calibrated phylogenetic hypotheses for nearly all of these reptile groups (*citations*), which make macroevolutionary inferences tractable. The Atlas of Living Australia also provides an unparalleled resource for species distributions based on museum occurrence records, for all described taxa. Given existing data and resources, and the access to material, I honestly believe that this is the best possible system to focus this research on.

## Significance

## Training Objectives

## Career Development

## Sponsoring Scientist and Institution Justification

I have chosen Drs. Lee and O'Meara specifically for their skill sets, research interests, and proven track records of pushing the evolutionary biology community forward. They are both creative and judicious thinkers, who I admire for their contributions to science and broader education.

**Dr. Brian O'Meara** (*Methods/Biology*) is an Associate Professor in the Department of Ecology & Evolutionary Biology at the University of Tennessee, Knoxville.

**Dr. Michael Lee** (*Theory/Biology*) is the curator of Palaeontology at the South Australian Museum, and holds a joint appointment in the Department of Biological Sciences at Flinders University of South Australia.

## Yearly Timetable (goals, outcomes)

## Broader Impacts Narrative

Accounting for these interactions in a phylogenetic framework is integral to understanding the evolution of both simple and complex traits, and fits with our existing knowledge of character displacement.

[1]

## References

1. Brennan IG, Oliver PM. 2017 Mass turnover and recovery dynamics of a diverse Australian continental radiation. *Evolution* **71**, 1352–1365. (doi:[10.1111/evo.13207](https://doi.org/10.1111/evo.13207))