Math6450 Assignment2: Multiple Linear Regression

Ian Tai Ahn September 19, 2025

1 Data Exploration

(a) Descriptive Statistics for Continuous Variables

Comprehensive Descriptive Statistics: Mean Median Std Dev Minimum u →Maximum Skewness Kurtosis claims 18.049 17.845 6.448 0.72 41. **⇔**39 0.254 0.095 2.490 1.905 deductible 0.51 10. 1.942 1.542 2.351 coverage 189.014 186.750 72.169 50.00 424. ⇔50 0.145 -0.292 15.438 11.000 14.227 1.00 85. **⇔**00 1.869 4.496 premium 2.969 2.945 0.822 0.50 ⇔78 0.245 0.030

(b) Correlation Matrix for Continuous Variables

Correlation Matrix:


	claims	deductible	coverage	age	ш
\hookrightarrow premium					
claims	1.000	-0.265	0.761	0.199	0.
deductible ⇔059	-0.265	1.000	-0.066	0.006	-0.
coverage ⇔723	0.761	-0.066	1.000	-0.015	0.
age	0.199	0.006	-0.015	1.000	0.
premium ⇔000	0.793	-0.059	0.723	0.314	1.

Variable with strongest linear relationship with $_{\sqcup}$ $_{\hookrightarrow}$ 'claims':

Variable: premium

Correlation coefficient: 0.793

Correlation Matrix Heatmap - Continuous Variables



Skewness Assessment:

Rule of thumb: |skewness| > 1 indicates highly $skewed_{\square}$ G

Rule of thumb: 0.5 < |skewness| < 1 indicates⊔

⇔moderately skewed distribution

claims:

Skewness: 0.254

Assessment: Approximately symmetric

deductible:

Skewness: 1.542

Assessment: Highly skewed

Log transformation skewness: 0.134
Improvement from log transformation: 1.408
Recommendation: Log transformation would improve

→normality

coverage:

Skewness: 0.145

Assessment: Approximately symmetric

age:

Skewness: 1.869

Assessment: Highly skewed

Log transformation skewness: -0.347 Improvement from log transformation: 1.523


Recommendation: Log transformation would improve⊔

→normality

premium:

Skewness: 0.245

Assessment: Approximately symmetric



Summary of Findings:

Variables with skewed distributions: deductible, age Variable most strongly correlated with claims: \Box \Box premium (r = 0.793)

Data Overview:

Total observations: 1,340 Variables analyzed: 5 Missing values: 0

2 Simple Linear Regression

Dataset Information:

Total observations: 1,340

Observations used in regression: 1,340

Missing values removed: 0

(a) Simple Linear Regression Model Fitting

Model Coefficients:

Intercept (β_0) : 5.2054 Slope (β_1) : 0.0679

Slope (β_1) : 0.0679

Fitted Regression Equation: Claims = $5.2054 + 0.0679 \times Coverage$

In mathematical notation:

 $\hat{y} = 5.2054 + 0.0679x$

where \hat{y} = predicted claims, x = coverage

(b) Interpretation of Slope Coefficient

Slope coefficient: 0.0679

Practical Interpretation:

- For every 1-unit increase in coverage, claims are ⊔ ⇔expected to increase by
- 0.0679 units, on average.

Alternative interpretation:

• For every 100-unit increase in coverage, claims_change by 6.79 units, on average.

Example predictions:

- Coverage = 100: Predicted Claims = 12.00
- Coverage = 150: Predicted Claims = 15.40
- Coverage = 200: Predicted Claims = 18.80
- Coverage = 250: Predicted Claims = 22.19

(c) Coefficient of Determination (R2) Analysis

Model Performance Metrics:

 ${\tt R}^2$ (Coefficient of Determination): 0.5784

R² as percentage: 57.84%

Correlation coefficient (r): 0.7605 Root Mean Square Error (RMSE): 4.1850

Interpretation of \mathbb{R}^2 :


- 57.84% of the variation in claims is explained by $_{\!\sqcup}$ $_{\!\hookrightarrow}$ coverage.
- 42.16% of the variation in claims is due to other $_{\mbox{$\sqcup$}}$ -factors not included in the

model.

• The linear relationship between coverage and claims $_{\mbox{\sc u}}$ $_{\mbox{\sc oth}}$ is moderate (R^2 = 0.5784).

Statistical Significance:

- t-statistic: 42.8442
- p-value: 0.0000
- Degrees of freedom: 1338
- The relationship is statistically significant at $_{\sqcup}$ $_{\hookrightarrow}$ the 5% level.



Summary Table:

Metric Value

 \hookrightarrow Interpretation
Intercept (β_0) 5.2054 Expected claims

Intercept (β_0) 5.2054 \rightarrow when coverage = 0

Slope (β_1) 0.0679 Change in claims per unit

⇔increase in coverage

 R^2 0.5784 57.8% of

→variance explained

Correlation (r) 0.7605 Linear

⇒association strength

RMSE 4.1850 Average

→prediction error

Observations 1340

Sample size

Key Findings Summary:

- Regression equation: Claims = $5.2054 + 0.0679 \times_{\square}$ \rightarrow Coverage
- Slope interpretation: Each additional unit of $_{\sqcup}$ $_{\hookrightarrow} coverage$ is associated with a
- 0.0679 unit change in claims
- Model explains 57.8% of the variation in claims
- The relationship is statistically significant (p = $_{\sqcup}$ 0.0000)

3 Multiple Regression Model

Dependent Variable: claims

Explanatory Variables: deductible, coverage, age, uprior_claims, premium

Dataset Information:

Total observations: 1,340 Complete cases used: 1,340

Observations removed (missing data): 0 Number of explanatory variables: 5

(a) Fitted Regression Equation

G. (6) data Batimata (a.m.) data 2 data 1 data 2

Coefficient Estimates (rounded to 3 decimal places): Intercept (β_0): 3.208

 β _1 (deductible): -0.728

β_{-2} (coverage): 0.062 β_{-3} (age): 0.091 β_{-4} (prior_claims): 2.580 β_{-5} (premium): 0.495
Fitted Regression Equation: Claims = 3.208 - 0.728 × deductible + 0.062 × $_{\Box}$ $_{\ominus}$ coverage + 0.091 × age + 2.580 × prior_claims + 0.495 × premium
Compact Mathematical Form: $ \hat{\mathbf{y}} = \beta_0 + \beta_1 \mathbf{x}_1 + \beta_2 \mathbf{x}_2 + \beta_3 \mathbf{x}_3 + \beta_4 \mathbf{x}_4 + \beta_5 \mathbf{x}_5 \\ \hat{\mathbf{y}} = 3.208 + -0.728 \mathbf{x}_1 + 0.062 \mathbf{x}_2 + 0.091 \mathbf{x}_3 + 2.580 \mathbf{x}_4 + \mathbf{y}_5 \mathbf{x}_5 $

(b) Standard Errors for Each Coefficient

 \rightarrow x₄=prior_claims, x₅=premium

where x_1 =deductible, x_2 =coverage, x_3 =age, \sqcup

Standard Errors:

Intercept (β_0) : 0.3172 β_-1 (deductible): 0.0394 β_-2 (coverage): 0.0020 β_-3 (age): 0.0068 β_-4 (prior_claims): 0.1210 β_-5 (premium): 0.2118

Additional Statistics (t-statistics and p-values): Estimate Std Error t-stat Coefficient →p-value Significance 0.3172 10.113 Ω Intercept 3.208 **⇔**0000 deductible -0.728 0.0394 -18.459 **⇔**0000 0.062 coverage 0.0020 30.624 0. **⇔**0000 age 0.091 0.0068 13.401 **⇔**0000 prior_claims 2.580 0.1210 21.316 Ο. **⇔**0000 premium 0.495 0.2118 2.338

Significance codes: *** p<0.001, ** p<0.01, * p<0.05

(c) Model Performance Statistics

 R^2 (Coefficient of Determination): 0.8130

Adjusted R^2 : 0.8123

⇔0195

Residual Standard Deviation: 2.7938

Additional Model Statistics:

Multiple R (Correlation): 0.9016

Residual Sum of Squares (RSS): 10412.1409

Mean Squared Error (MSE): 7.8052

F-statistic: 1159.6202

F-statistic p-value: 0.000000

Overall model significance: Yes (α = 0.05)


Degrees of Freedom:

Model: 5 Residual: 1334 Total: 1339

Summary Results Table:


Variable Coefficient Std_Error ⊔

GCoefficient_Rounded



Model Performance Table:

 $\begin{array}{ccc} Statistic & Value \\ R^2 & 0.8130 \\ Adjusted R^2 & 0.8123 \\ Residual Std Deviation & 2.7938 \\ F-statistic & 1159.6202 \\ p-value & (F-test) & 0.000000 \\ Observations & 1340 \\ Variables & 5 \end{array}$



Key Results Summary:

- \checkmark Multiple regression equation fitted with 5_{\sqcup}
- ⇔explanatory variables
- \checkmark Model explains 81.3% of variance in claims (R² = 0. 8130)
- \checkmark Adjusted R^2 = 0.8123 (accounts for number of $_{\sqcup}$ $_{\hookrightarrow}$ variables)
- √ Residual standard deviation = 2.7938
- $\sqrt{}$ Overall model is significant (F-test p-value = 0. \rightarrow 000000)
- \checkmark Standard errors calculated for all 6 coefficients

4 Statistical Inference

Multiple Linear Regression Model: Claims vs⊔

Goductible, Coverage, Age,

Prior_Claims, Premium)

Model Summary: Observations: 1340

Variables: 5

Degrees of freedom (residual): 1334

 R^2 : 0.8130 MSE: 7.8052

Coefficient Estimates:

 $\begin{array}{lll} \mbox{Variable} & \mbox{Coefficient Std Error} & \mbox{t-statistic}_t \\ \hookrightarrow \mbox{p-value} \end{array}$

deductible → 0.0000	-0.7278	0.0394	-18.4591	L
coverage	0.0621	0.0020	30.6239	ı
age ⇔ 0.0000	0.0906	0.0068	13.4010	L
prior_claims	2.5797	0.1210	21.3156	L
premium	0.4953	0.2118	2.3382	L

(a) Testing Significance of Age Coefficient Hypothesis Test for Age Coefficient:

Null Hypothesis (H_0): β _age = 0 Alternative Hypothesis (H_1): β _age \neq 0 Significance level (α): 0.05 Test type: Two-tailed t-test

Test Statistics:

Age coefficient (β _age): 0.0906 Standard error (SE): 0.0068 t-statistic: 13.4010 Degrees of freedom: 1334

p-value: 0.0000

Critical value (±): 1.9617

Decision Rule:

Reject H_0 if |t-statistic| > 1.9617 OR if p-value < 0.

Conclusion:

 \checkmark REJECT H_0 : The coefficient for age IS $_{\mbox{\tiny LI}}$ ⇔statistically significant at the 5%

|t-statistic| = 13.4010 > 1.9617p-value = 0.0000 < 0.05

(b) 95% Confidence Interval for Prior Claims $_{\mbox{\scriptsize L}}$ Goefficient

Confidence Interval Calculation: Coefficient (β _prior_claims): 2.5797 Standard error: 0.1210

Degrees of freedom: 1334 Confidence level: 95%

Confidence Interval Formula:

CI = $\beta \pm t_{\alpha/2,df} \times SE(\beta)$

 $CI = 2.5797 \pm 1.9617 \times 0.1210$

 $CI = 2.5797 \pm 0.2374$

95% Confidence Interval for Prior Claims Coefficient: [2.3423, 2.8171]

Practical Interpretation:

• We are 95% confident that the true effect of having ⇒prior claims on current

claims

is between 2.3423 and 2.8171 units.

 \bullet Since the entire interval is positive, prior claims $_{\sqcup}$ ${\scriptstyle \hookrightarrow} {\sf consistently\ INCREASE}$

current claims.

→higher current claims than

those without.

- The width of the interval (0.4748) indicates the ⇒precision of our estimate.
- (c) Overall F-test for Model Significance Overall F-test for Regression Model:

Null Hypothesis (H₀): β_1 = β_2 = β_3 = β_4 = β_5 = 0 (All explanatory variables have no effect on claims) Alternative Hypothesis (H₁): At least one $\beta_i \neq 0$ (At least one explanatory variable has a ⇒significant effect)

Significance level (α): 0.05

Test Statistics:

Total Sum of Squares (TSS): 55667.4953 Explained Sum of Squares (ESS): 45255.3543 Residual Sum of Squares (RSS): 10412.1409 Mean Square Regression (MSR): 9051.0709 Mean Square Error (MSE): 7.8052

F-statistic: 1159.6202 Degrees of freedom: (5, 1334)

p-value: 0.000000

Critical F-value (α = 0.05): 2.2208

Decision Rule:

Reject ${\rm H}_0$ if F-statistic > 2.2208 OR if p-value < 0.05

Conclusion:


 \checkmark REJECT H_0 : The regression model IS statistically \hookrightarrow significant at the 5% level. F-statistic = 1159.6202 > 2.2208 p-value = 0.000000 < 0.05

At least one explanatory variable has a significant \hookrightarrow effect on claims.

⇔variation in claims.

Model Performance Context:

 R^2 = 0.8130 (81.3% of variance explained) The model performs well in predicting claims.



Summary of All Statistical Tests:

Test Statistic _ →p-value Conclusion Age Coefficient (t-test) t = 13.4010 0. Significant Prior Claims CI CI = [2.3423, 2.8171]N/ →A Does not contain 0

```
F = 1159.6202 0.
  Overall Model (F-test)
                                                          Detailed Mathematical Form:
 ⇔000000 Model Significant
                                                          Claims = 3.027 + -0.713 \times \text{deductible} + 0.058 \times \text{coverage}
                                                                    + 0.077 \times age + 2.392 \times prior_claims + 1.
LaTeX Summary Table:
                                                            ⇒019×premium
\begin{table}
                                                                    + -1.419 \times type + 0.859 \times location
\caption{Summary of Statistical Tests}
                                                           (b) Interpretation of Type Coefficient
\label{tab:hypothesis_tests}
                                                           Type Coefficient Analysis:
\begin{tabular}{1111}
                                                          Coefficient (\beta_type): -1.419
\toprule
                                                          Standard Error: 0.1699
Test & Statistic & p-value & Conclusion \\
                                                          t-statistic: -8.355
                                                          p-value: 0.0000
Age Coefficient (t-test) & t = 13.4010 & 0.0000 &
 ⇔Significant \\
                                                           Type variable coding: [np.int64(0), np.int64(1)]
Prior Claims CI & CI = [2.3423, 2.8171] & N/A & Does_
 ⇔not contain 0 \\
                                                          Practical Interpretation:
Overall Model (F-test) & F = 1159.6202 & 0.000000 & _{\sqcup}
                                                           • Properties with type = 1 have claims that are 1.419

→Model Significant \\
                                                            \hookrightarrowunits LOWER than
\bottomrule
                                                           properties with type = 0,
\end{tabular}
                                                            holding all other variables constant.
\end{table}
                                                          Assuming standard coding (0 = Commercial, 1 =_{\sqcup}
5 Binary Variables and Model Interpretation
                                                           ⇔Residential):
                                                           • Residential properties have claims that are 1.419
Adding 'type' and 'location' to the original model
                                                            ⇔units lower than commercial
Dependent Variable: claims
                                                          properties.
Original Variables: deductible, coverage, age, u
                                                           →prior_claims, premium
                                                            ⇒with higher insurance
New Variables: type, location
                                                          claims.
Data Summary:
                                                          Statistical Significance:
Original model observations: 1,340
                                                           Extended model observations: 1,340
                                                           \Rightarrow (p = 0.0000 < 0.05)
                                                           • We can be confident that property type has a real_
Extended Model Summary:
Observations: 1340
                                                           ⇔effect on claims.
Variables: 7
                                                           (c) Partial F-test for Model Improvement
R^2: 0.8263
                                                          Model Comparison (same sample size: 1340):
Adjusted R^2: 0.8254
                                                          Model
                                                                                \mathbb{R}^2
                                                                                              Adj R^2
Residual Standard Error: 2.6939
                                                           ⇔Variables RSS
(a) Extended Regression Model Equation
                                                           Original
                                                                                0.8130
                                                                                              0.8123
                                                                                                           5
Coefficient Estimates:
                                                           → 10412.1409
Variable
                Coefficient Std Error
                                           t-stat
                                                          Extended
                                                                                0.8263
                                                                                              0.8254
                                                                                                           7
→p-value
                                                                 9666.7444
Intercept
                3.027
                              0.3171
                                                          R<sup>2</sup> Improvement: 0.0134 (1.34 percentage points)
deductible
                -0.713
                              0.0381
                                           -18.706
 ⇔0.0000
                                                          Partial F-test:
coverage
                0.058
                              0.0022
                                           26.539
                                                          {\rm H_0}\colon\ \beta\_{\rm type} = \beta\_{\rm location} = 0 (binary variables add {\rm no_{\sqcup}}
 ⇔0.0000
                                                           →explanatory power)
                0.077
                              0.0070
                                           10.935
age
                                                          H_1: At least one of \beta_type or \beta_location \neq 0 (binary_
 40.0000
                                                           ⇔variables improve the
prior_claims
                2.392
                              0.1254
                                           19.077
                                                          model)
 ⇔0.0000
premium
                1.019
                              0.2378
                                           4.284
                                                          Partial F-test Calculations:
 ⇔0.0000
                                                          RSS(original): 10412.1409
type
                -1.419
                              0.1699
                                           -8.355
                                                          RSS(extended): 9666.7444
 ⇔0.0000
                                                          Reduction in RSS: 745.3965
location
                0.859
                              0.1731
                                           4.959
                                                          Additional variables (q): 2
40.0000
                                                          DF residual (extended): 1332
Fitted Regression Equation:
                                                          F-statistic: 51.3548
Claims = 3.027 - 0.713 \times deductible + 0.058 \times_{\square}
                                                          Degrees of freedom: (2, 1332)
 \hookrightarrowcoverage + 0.077 	imes age + 2.392 	imes
                                                          p-value: 0.0000
prior_claims + 1.019 \times premium - 1.419 \times type + 0.
                                                          Critical F-value (\alpha = 0.05): 3.0025
 ⇔859 × location
                                                          Conclusion:
```


√ REJECT H₀: Adding type and location SIGNIFICANTLY

___ ⇒improves the model F = 51.3548 > 3.0025p-value = 0.0000 < 0.05

The binary variables provide significant additional \Box ⇔explanatory power.

Model Improvement Assessment:

- R^2 improved by 0.0134 (1.34 percentage points) -→this is modest
- Extended model explains 82.6% vs 81.3% of variance
- \bullet Adjusted R^2 increased from 0.8123 to 0.8254
- The improvement in adjusted R² suggests the added_□ \hookrightarrow variables are worthwhile



Executive Summary:

Aspect Finding Extended Model Equation Claims = 3.027 + ... + -1. \hookrightarrow 419×type + 0.859×location Type Coefficient -1.419 Type Effect Type=1 ⇔has 1.419 lower claims Statistical Significance \hookrightarrow Significant (p = 0.0000) R² Improvement 0.0134 →(1.34 percentage points) Partial F-test Result Significant_

6 Interaction Effects

⇔improvement (p = 0.0000)

Regression Model with Interaction Term: Deductible \times_{\sqcup} Type

Model Features: deductible, type, coverage, age, ⊔ →prior_claims, premium

Interaction Term: deductible \times type

Data Summary:

Total observations: 1,340 Complete cases used: 1,340 Missing values removed: 0

Type variable coding: [np.int64(0), np.int64(1)]

Interaction Term (deductible \times type) Statistics: Mean: 1.5335

Std Dev: 1.9042

Range: [0.0000, 10.0000]

Model Summary: R^2 : 0.8233

Adjusted R^2 : 0.8224

Residual Standard Error: 2.7172

F-statistic: 886.8341

Coefficient Estimates:

Variable → p-value	Coefficient Sig	Std Error	t-stat L
→ p varue	nig		
Intercept	3.2856	0.3300	
deductible	-0.6729	0.0596	-11.2894
→ 0.0000	***		
type	-1.2573	0.2598	-4.8392
→ 0.0000	***		
coverage	0.0553	0.0021	25.9580
→ 0.0000	***		
age	0.0703	0.0070	10.1034
→ 0.0000	***		
prior_claims	2.2568	0.1234	18.2905
→ 0.0000	***		
premium	1.3647	0.2290	5.9595
→ 0.0000	***		
deductible_x_t	ype -0.0946	0.0779	-1.2151
→ 0.2245			

Significance codes: *** p<0.001, ** p<0.01, * p<0.05

(a) Regression Function with Interaction Term General Form:

 $\texttt{Claims} = \beta_0 \ + \ \beta_1 \times \texttt{deductible} \ + \ \beta_2 \times \texttt{type} \ + \ \beta_3 \times \texttt{coverage}_{\square}$ \hookrightarrow + $\beta_4 \times \text{age}$ + $\beta_5 \times \text{prior_claims}$ + $\beta_6 \times \text{premium} + \beta_7 \times (\text{deductible} \times \text{type}) + \varepsilon$

Fitted Regression Equation:

Claims = $3.2856 - 0.6729 \times \text{deductible} - 1.2573 \times \text{type} + 1.2573 \times \text{type}$ \hookrightarrow 0.0553 \times coverage + 0.0703 \times age

+ 2.2568×prior_claims + 1.3647×premium - 0. ⇔0946×(deductible×type)

With Coefficient Values:

Claims = $3.2856 + -0.6729 \times \text{deductible} + -1.2573 \times \text{type}$ + 0.0553×coverage + 0.0703×age + 2.

⇒2568×prior_claims

+ $1.3647 \times \text{premium} + -0$.

 \hookrightarrow 0946 \times (deductible \times type)

(b) Interpretation of Deductible Effect by Property⊔

Key Coefficients:

 β_1 (deductible): -0.6729

 β_2 (type): -1.2573

 β_7 (deductible×type): -0.0946

Interpretation of Interaction Effect:

The interaction model allows the effect of deductible →to differ by property

For Commercial Properties (type = 0):

 $\partial \text{Claims}/\partial \text{deductible} = \beta_1 + \beta_7 \times 0 = \beta_1 = -0.6729$

• A 1-unit increase in deductible changes claims by \hookrightarrow -0.6729 units for commercial

properties.

For Residential Properties (type = 1):

 $\partial {\rm Claims}/\partial {\rm deductible}$ = β_1 + $\beta_7 \times 1$ = β_1 + β_7 = -0.6729 $_{\Box}$ + -0.0946 = -0.7675

residential properties.

Comparison:

Difference in deductible effect: -0.0946

• The deductible effect is 0.0946 units MORE NEGATIVE

defor residential

properties.

• Deductible increases have a stronger negative deffect on residential claims

than commercial claims.

Practical Business Interpretation:

- \bullet This association is STRONGER for residential $_{\square}$ -properties
- (c) Statistical Significance Test for Interaction Term Hypothesis Test for Interaction Term:

 $\mathbf{H}_0 \colon \beta_7$ = 0 (no interaction between deductible and utype)

 ${\rm H_1}\colon\,\beta_7\neq$ 0 (significant interaction exists) Significance level: α = 0.05

Test Statistics:

Interaction coefficient (β_7): -0.0946

Standard error: 0.0779 t-statistic: -1.2151 Degrees of freedom: 1332

p-value: 0.2245

Critical value (\pm): 1.9617

Decision Rule:

Reject ${\rm H}_0$ if |t-statistic| > 1.9617 OR if p-value < 0. ${}_{\rm \hookrightarrow}05$

Conclusion:

FAIL TO REJECT $\mathrm{H}_0\colon$ The interaction term is NOT_{\square} -statistically significant at the

5% level.

|t-statistic| = 1.2151 \le 1.9617


p-value = 0.2245 \geq 0.05

property types.

The interaction term may not be necessary.

95% Confidence Interval for Interaction Coefficient: [-0.2473, 0.0581]

uncertain



Executive Summary:

Aspect

Result

Model Specification Claims ~ deductible + type + $_{\!\!\!\!\square}$ +coverage + age +

prior_claims + premium + deductiblextype

Interaction Coefficient

-0.0946 (SE = 0.0779)

Commercial Effect

-0.6729 per unit deductible

Residential Effect

-0.7675 per unit deductible

Difference

-0.0946

Statistical Significance Not significant (p = 0.2245)

Model \mathbb{R}^2

0.8233

Model Interpretation:

similar across commercial and residential properties

• A simpler model without interaction may be adequate

7 Residual Analysis

Extended Multiple Linear Regression Model Variables: deductible, coverage, age, prior_claims,_u

→premium, type, location

Model Summary:

Observations: 1,340

Variables: 7

 R^2 : 0.8263

Residual Standard Error: 2.6939

(a) Residuals vs Fitted Values Analysis Residuals vs Fitted Values Analysis: Residual range: [-3.376, 15.203] Fitted values range: [0.792, 39.985]

Pattern Analysis:

- Variance appears roughly constant
- Correlation magnitude suggests homoscedasticity⊔

 ⇔(constant variance)


Linearity Assessment:

Mean residuals by fitted value terciles:

- Low tercile: -0.0800 • Middle tercile: 0.0229 • High tercile: 0.0572
- Maximum deviation from zero: 0.0800 (suggests⊔

→linear relationship is

appropriate)



(b) Q-Q Plot and Normality Analysis

Normality Test Results: Shapiro-Wilk Test:

Statistic: 0.8106 p-value: 0.0000

REJECT normality at $\alpha \text{=0.05}$

Jarque-Bera Test:

Statistic: 2188.1490 p-value: 0.0000

REJECT normality at α =0.05

Kolmogorov-Smirnov Test:

Statistic: 0.1468 p-value: 0.0000

REJECT normality at $\alpha \text{=0.05}$

Descriptive Statistics for Normality:

Skewness: 1.9531 (Normal \approx 0) Kurtosis: 4.8921 (Normal \approx 0)

Skewness interpretation: highly skewed Kurtosis interpretation: heavy-tailed

Overall Normality Assessment: Assumption appears to $_{\sqcup}$

⇒be violated

(c) Outliers and Influential Points Analysis Diagnostic Thresholds:

Outlier threshold (standardized residuals): ± 3

High leverage threshold: 0.0119

High Cook's distance threshold: 0.0030

Outliers and Influential Points:

Observations with |standardized residuals| > 3: 31

Observations with |studentized residuals| > 3: 31

High leverage points: 73

High Cook's distance points: 74

Most Extreme Observations:

Highest Residual: Observation 315

Fitted value: 23.547 Actual value: 38.750 Standardized residual: 5.643

Leverage: 0.0072

Cook's distance: 0.0331

Highest Leverage: Observation 262

Fitted value: 34.070 Actual value: 36.160

Standardized residual: 0.776

Leverage: 0.0305

Cook's distance: 0.0027 Highest Cooks: Observation 315

Fitted value: 23.547 Actual value: 38.750

Standardized residual: 5.643

Leverage: 0.0072

Cook's distance: 0.0331

<Figure size 640x480 with 0 Axes>

Detailed Analysis of Problematic Observations:

	ual Std_Residual	Leverage	Cooks_D	ш
	670 3.412	0.0032	0.0054 _L	
2 5.711 3. → High Leve	340 -0.880	0.0122	0.0014	П
14 20.959 20. → High Leve	000 -0.356	0.0128	0.0002	П
36 10.929 8. → High Leve	700 -0.827	0.0142	0.0014	ш
70 13.967 11.	670 -0.852	0.0130	0.0014	ш
71 20.337 24.	990 1.727	0.0074	0.0032	ш
→ High Cool 73 30.965 29.	670 -0.481	0.0141	0.0005	П
→ High Level 118 22.728 22.	290 -0.163	0.0193	0.0001	П
→ High Level 122 5.247 10.	1.805	0.0072	0.0034	ш
→ High Cool129 31.861 36.→ High Cool	730 1.807	0.0092	0.0043	ш

... and 124 more observations with issues.

Diagnostic Summary:

- 1. Linearity: suggests linear relationship is ⊔ →appropriate
- ⇔(constant variance)
- 3. Normality: Assumption appears to be violated
- 4. Outliers: 31 potential outliers identified
- 5. Influential Points: 74 high Cook's distance ⇔observations

Recommendations:

- \bullet Consider transformation of variables or \mathtt{robust}_\sqcup \hookrightarrow regression methods
- ullet Examine influential points consider their impact ⇔on coefficient estimates

8 Model Comparison and Selection

Comparing three different model specifications: Model A: claims ~ deductible + coverage + age + →prior_claims + premium

Model B: claims ~ deductible + coverage + age + $_{\sqcup}$ →prior_claims + premium + type +

```
location
                                                         Nested Model Comparisons (F-tests):
Model C: claims ~ deductible + coverage +⊔
                                                         Model A vs Model B:

→prior_claims + premium + type

                                                           F-statistic: 51.3548
                                                           p-value: 0.0000
                                                           Model B significantly better
Data Summary:
                                                           Note: Model A vs C and Model B vs C are not nested_{\sqcup}
Original dataset size: 1,340
Complete cases for all models: 1,340
                                                          ⇔comparisons
Cases removed due to missing data: 0
                                                         (b) Model Recommendation and Analysis
                                                         Statistical Criteria Analysis:
----- Model A -----
Variables: deductible, coverage, age, prior_claims, __
                                                         1. Goodness of Fit:
 →premium
                                                            • \mathbb{R}^2 ranking: Model B > others
• Adjusted \mathbb{R}^2 ranking: Model B > others
Number of variables: 5
R^2: 0.8130
                                                            \bullet R^2 improvement from A to B: 0.0134
Adjusted R^2: 0.8123
                                                             \bullet Adjusted R^2 change from A to B: 0.0132
Residual Standard Deviation: 2.7938
AIC: 6566.17
                                                         2. Model Parsimony:
BIC: 6592.18
                                                             • AIC favors: Model B (AIC = 6472.65)
Significant coefficients (p < 0.05): 5/5
                                                             • BIC favors: Model B (BIC = 6509.05)
                                                            • BIC penalizes complexity more heavily than AIC
----- Model B -----
Variables: deductible, coverage, age, prior_claims, u
                                                         3. Coefficient Significance:

→premium, type, location
                                                             • Model A: 5/5 coefficients significant (100.0%)
Number of variables: 7
                                                            • Model B: 7/7 coefficients significant (100.0%)
R^2: 0.8263
                                                            • Model C: 5/5 coefficients significant (100.0%)
Adjusted R^2: 0.8254
Residual Standard Deviation: 2.6939
                                                         4. Prediction Accuracy:
AIC: 6472.65
                                                             • Lowest prediction error: Model B (SD = 2.6939)
BIC: 6509.05
Significant coefficients (p < 0.05): 7/7
                                                         Practical Interpretability Analysis:
----- Model C -----
                                                         1. Variable Inclusion Logic:
Variables: deductible, coverage, prior_claims,_
                                                            • Model A: Core financial variables (deductible, u
→premium, type
                                                          ⇔coverage, premium) + risk
Number of variables: 5
                                                         factors (age, prior_claims)
R^2: 0.8095
                                                             • Model B: Model A + property characteristics ⊔
Adjusted R^2: 0.8088
                                                          ⇔(type, location)
Residual Standard Deviation: 2.8197
                                                            • Model C: Simplified version with key variables +
ATC: 6590.93
                                                          ⇔property type
BIC: 6616.94
Significant coefficients (p < 0.05): 5/5
                                                         2. Business Relevance:
                                                            • Age variable: Present in A, Present in B, Absent
                                                          ⇔in C
(a) Model Comparison Table
                                                            \bullet Property type: Absent in A, Present in B, _{\sqcup}
Primary Comparison Metrics:
 \hookrightarrowPresent in C
Model A 5 vars 0.8130 0.8123
Model B 7 vars 0.8263 0.8254
                                       2.7938
                                                            • Location: Absent in A, Present in B, Absent in C
                                       2.6939
Model C 5 vars 0.8095 0.8088
                                       2.8197
                                                         3. Marginal Contribution Analysis:
                                                            • Adding type + location (B vs A): R<sup>2</sup> improves by
Additional Model Selection Criteria:
                                                           ⇔0.0134
 Model AIC BIC F_statistic Sig_Coefs
                                                            • Adjusted R<sup>2</sup> change: 0.0132 (improvement)
Model A 6566.17 6592.18 1159.62
                                           5/5
Model B 6472.65 6509.05
                             905.51
                                           7/7
                                                         Recommendation Framework:
Model C 6590.93 6616.94
                           1133.51
                                           5/5
                                                         Composite Scoring (weighted combination of criteria):
Best Model by Criterion:
                                                            • Model B: 1.000
• Highest R<sup>2</sup>: Model B (0.8263)
                                                            • Model A: 0.700
• Highest Adjusted R<sup>2</sup>: Model B (0.8254)
                                                            • Model C: 0.400
• Lowest Residual SD: Model B (2.6939)
• Lowest AIC: Model B (6472.65)
                                                          RECOMMENDED MODEL: Model B
• Lowest BIC: Model B (6509.05)
                                                         Justification for Model B:
                                                            \checkmark Highest predictive power (R^2 = 0.8263)
Model Complexity Analysis:
Model A: 5 variables, R^2/var = 0.1626
                                                            \checkmark Includes important property characteristics
Model B: 7 variables, R^2/var = 0.1180
                                                            \checkmark Comprehensive variable coverage
Model C: 5 variables, R^2/var = 0.1619


√ Best for prediction accuracy
```

Limitations of Model B:

More complex with potential overfitting risk May have multicollinearity issues

Alternative Recommendations by Use Case:

- For prediction accuracy: Model B
- For model parsimony: Model B
- For balanced approach: Model B
- For regulatory reporting: Model A (simplest, ⊔ →most interpretable)



9 Practical Application

Features shape: (1340, 7) Target shape: (1340,)

=== MODEL RESULTS === R-squared: 0.8263

Adjusted R-squared: 0.8254

Model Coefficients:

	Feature	Coefficient
0	Intercept	3.026950
1	deductible	-0.713391
2	coverage	0.058017
3	age	0.076546
4	prior_claims	2.391648
5	premium	1.018707
6	type	-1.419290
7	location	0.858614


Statistical Significance:

Duau	btatibtical bignificance.				
	Feature	Coefficient	Std_Error	t_statistic 📙	
\hookrightarrow	p_{value}	\			
0	Intercept	3.026950	0.316198	9.572968 👝	
⇔ 0.	000000e+00				
1	${\tt deductible}$	-0.713391	0.038024	-18.761697 👝	
⇔ 0.	000000e+00				
2	coverage	0.058017	0.002180	26.618766 👝	
⇔ 0.	000000e+00				
3	age	0.076546	0.006979	10.967841 👝	
⇔ 0.	000000e+00				
4 p	rior_claims	2.391648	0.124993	19.134246 🔟	
⇔ 0.	000000e+00				
5	premium	1.018707	0.237095	4.296623 _⊔	
⇔1 .	860187e-05				
6	type	-1.419290	0.169370	-8.379819 🔟	
⇔ 0.	000000e+00				


7 location 0.858614 0.172627 4.973820 \hookrightarrow 7.420302e-07

	Significan
0	Tru
1	Tru
2	Tru
3	Tru
4	True
5	True
6	Tru
7	Tru

=== MODEL DIAGNOSTICS ===
Mean Squared Error: 7.2140
Root Mean Squared Error: 2.6859



=== CORRELATION ANALYSIS ===



age 0.198837 location 0.105441 type -0.061114 Name: claims, dtype: float64

=== PART (a): PREDICTION ===

Understanding categorical variables:

Type values: [1 0]
Location values: [1 0]
Type value counts: type

1 835 0 505

Name: count, dtype: int64 Location value counts: location

968
 372

Name: count, dtype: int64

Prediction for the given property: Expected claims amount: 19.49

Sensitivity analysis for categorical variables:

Type=Commercial, Location=Rural: 20.05 Type=Commercial, Location=Urban: 20.91 Type=Residential, Location=Rural: 18.63 Type=Residential, Location=Urban: 19.49

PART (b): BUSINESS IMPLICATIONS AND RECOMMENDATIONS

Feature Importance (by absolute coefficient value):

	Feature	Coefficient	Abs_Coefficient
3	prior_claims	2.391648	2.391648
5	type	-1.419290	1.419290
4	premium	1.018707	1.018707
6	location	0.858614	0.858614
0	deductible	-0.713391	0.713391
2	age	0.076546	0.076546
1	coverage	0.058017	0.058017

Prediction Confidence Interval (95.0%):

Expected claims: 19.49 Lower bound: 14.22 Upper bound: 24.76

BUSINESS RECOMMENDATIONS:

1. PRICING STRATEGY:

- Most significant factors should drive $\mathtt{premium}_{\square}$ -calculations

2. RISK FACTORS ANALYSIS:

Based on the coefficients, focus on:

- Variables with largest absolute coefficients $% \left(1\right) =\left(1\right) \left(1\right)$
- Statistically significant predictors (p < 0.05)
- $\mbox{\sc High}$ correlation factors with claims

3. UNDERWRITING GUIDELINES:

- Properties with high predicted claims may need:
 - * Higher premiums
 - * Additional risk assessment
 - * Different deductible structures
- Consider segmented pricing models

4. PORTFOLIO MANAGEMENT:

- Monitor actual vs predicted claims regularly

5. OPERATIONAL INSIGHTS:

- Use model predictions for:
 - * Reserve allocation
 - * Risk-based pricing
 - * Customer segmentation
 - * Fraud detection (outliers in residuals)

OUTLIER ANALYSIS:

Properties with unusually high/low claims (>2 std_devs): 66

These may require special investigation for:

- Fraud detection
- Model improvement opportunities
- Special risk factors not captured in current model

10 Critical Thinking

PART (a) MULTIPLE LINEAR REGRESSION ASSUMPTIONS


→ ANALYSIS

The key assumptions of multiple linear regression are:

- 1. LINEARITY: The relationship between predictors and $_{\!\!\!\sqcup}$ -response is linear
- 2. INDEPENDENCE: Observations are independent of each $_$ other
- ${\tt 4.\ NORMALITY:\ Residuals\ are\ normally\ distributed}$
- 5. NO MULTICOLLINEARITY: Predictors are not highly occrrelated with each other

Let's test each assumption:

1. LINEARITY ASSUMPTION



LINEARITY ASSESSMENT:

- Examine scatter plots for linear patterns

Correlations with claims:

deductible: -0.265 coverage: 0.761 age: 0.199 premium: 0.793

INSURANCE CONTEXT IMPLICATIONS:

- Insurance claims may have non-linear relationships $_{\sqcup}$ $_{\hookrightarrow}$ (e.g., coverage thresholds)
- Age effects might be non-linear (newer vs very old $_{\!\sqcup}$ -properties)
- Premium-claims relationship might be non-linear \mathtt{due}_{\sqcup} -to risk-based pricing
- 2. INDEPENDENCE ASSUMPTION

INDEPENDENCE ASSESSMENT:

- Cannot be fully tested without knowing \mathtt{data}_{\sqcup}
- ${ \hookrightarrow } \texttt{collection} \ \texttt{method}$
- Check for patterns in residuals order




Durbin-Watson statistic: 2.038

(Values near 2.0 suggest independence, <1.5 or >2.5 $_{\mbox{\tiny \mbox{\tiny \mbox{\tiny \mbox{\tiny }}}}}$ suggest correlation)

INSURANCE CONTEXT IMPLICATIONS:

- Properties in same area might have correlated risks_□ ⇔(floods, earthquakes)
- Temporal clustering if data spans multiple years
- \hookrightarrow with economic changes
- Policy renewals might create dependencies
- 3. HOMOSCEDASTICITY (CONSTANT VARIANCE) ASSUMPTION




Breusch-Pagan test:

p-value: 0.1076


LM statistic: 11.7914

Heteroscedasticity detected: No



INSURANCE CONTEXT IMPLICATIONS:

- Heteroscedasticity common in insurance data
- May need weighted regression or transformation
- 4. NORMALITY OF RESIDUALS ASSUMPTION



NORMALITY TESTS:

Shapiro-Wilk test:

Statistic: 0.8106, p-value: 0.0000

Normal distribution: No

Jarque-Bera test:

Statistic: 2188.1490, p-value: 0.0000


Normal distribution: No

Descriptive statistics:

Skewness: 1.9531 Kurtosis: 4.8921

INSURANCE CONTEXT IMPLICATIONS:

- Insurance claims often right-skewed (many small, $_{\mbox{\sc d}}$ -few large claims)
- May need log transformation or robust $regression_{\sqcup}$ \hookrightarrow methods
- Non-normality affects confidence intervals and $_{\!\sqcup}$ hypothesis tests
- 5. NO MULTICOLLINEARITY ASSUMPTION



HIGH CORRELATIONS (|r| > 0.7): coverage - premium: 0.723

VARIANCE INFLATION FACTORS:

	Variable	VIF
0	deductible	2.302515
1	coverage	36.103529
2	age	3.980363
3	prior_claims	3.974976
4	premium	86.843712
5	type	3.287249
6	location	3.743009

VIF > 5: Moderate multicollinearity VIF > 10: High multicollinearity

Variables with high VIF:


Variable VIF

- 1 coverage 36.103529
- 4 premium 86.843712

INSURANCE CONTEXT IMPLICATIONS:

- Premium and coverage likely correlated (higher ∪ coverage = higher premium)
- Deductible and coverage might be related
- 6. NO OUTLIERS/INFLUENTIAL POINTS ASSUMPTION OUTLIER DETECTION:

Observations with $|standardized\ residuals| > 2.5$: 48 Observations with high Cook's distance: 13



Outlier observations (standardized residuals > 2.5): claims deductible coverage age premium

0	22.67	1.44	165.7	2	2.23
182	18.12	5.67	131.1	30	2.05
203	35.65	5.08	378.5	8	4.39
247	18.22	7.30	214.5	1	2.29
269	28.41	1.20	209.1	13	2.89

INSURANCE CONTEXT IMPLICATIONS:

- Large claims are natural in insurance (catastrophic $_{\sqcup}$ $_{\square}$ events)
- Outliers might represent legitimate extreme events, ⊔ ⇔not errors

OVERALL ASSUMPTION ASSESSMENT FOR INSURANCE CLAIMS

LIKELY VIOLATED ASSUMPTIONS:

- 1. Linearity: Insurance relationships often non-linear
- 2. Normality: Claims typically right-skewed
- 3. Homoscedasticity: Variance often increases with calaim size
- 4. Independence: Geographic/temporal clustering ⊔
 →possible

RECOMMENDED SOLUTIONS:

- 1. Log transformation of claims (handle skewness)
- 2. Robust regression methods
- 3. Polynomial or interaction terms
- 4. Weighted least squares (address heteroscedasticity)
- 5. Consider GLM (Gamma or Poisson regression)
- 6. Outlier-robust methods

PART (b) Additional Useful Variables

- 1. PROPERTY-SPECIFIC VARIABLES
- Construction: Building materials, roof type/age, ∪ year built, size, stories
- Condition: Recent renovations, security features,

2. ENVIRONMENTAL & GEOGRAPHIC

- Climate: Climate zones, precipitation, natural

 disaster scores
- Location: Crime rates, distance to fire station/ water, building codes

3. ECONOMIC & DEMOGRAPHIC

- Economic: Local income, property appreciation, ⊔
 →unemployment rate
- Demographics: Owner vs tenant occupied, primary $vs_{\mbox{\tiny \square}}$ -secondary residence

4. USAGE & BEHAVIORAL

- Property Use: Home business, rental income, vacancy $\underline{\mbox{\ }}$
- Claims History: Previous claim types, time \mathtt{since}_{\square} ${\hookrightarrow}\mathtt{last}$ claim
- Behavior: Payment history, policy shopping, service

 →interactions

5. ADVANCED MODELING

- Interaction Effects: Age×Construction, __ __Location×Weather, Coverage×Deductible
- External Data: Credit scores, satellite imagery, ⊔ →weather APIs

6. IMPLEMENTATION PRIORITY

- HIGH: Natural disaster scores, construction ⊔ details, claims history
- \hookrightarrow patterns
- LOW: Credit indicators, satellite analysis, ⇔economic metrics

7. EXPECTED OUTCOMES

- Model Accuracy: 60-80% → 85-95% predictive accuracy
- Benefits: Better risk selection, fraud detection, →dynamic pricing

8. KEY CONSIDERATIONS

- Data availability varies by property
- Quality validation required for third-party data
- Regulatory compliance (fair housing laws)
 Cost-benefit analysis essential