

Практикум E1

73 marks, 20 blocks, nine techniques. The whole topic is one question asked over and over: what does this expression approach? You answer it by substituting. When substituting works, you are finished in one line. When it gives $\frac{0}{0}$ or $\frac{\infty}{\infty}$ you have learned nothing yet — and everything else in this practicum is what to do next.

О практикуме

Пятый практикум серии на английском, устроенный как B2–B5: ноутбук, карточка приёмов и сообщения проверок английские, документация репозитория остаётся русской.

Заданий

0

Раздел решений

включён

Интерактивная версия

kaggle.com/code/artemkonukhov/ib-math-praktikum-e1-limits

Собрано из ноутбука репозитория `ib-math-aa-hl-past-papers`. Проверки в PDF не работают — они для интерактивной версии.

Practicum E1: limits, indeterminate forms, l'Hôpital's rule

73 marks, 20 blocks, nine techniques. The whole topic is one question asked over and over: *what does this expression approach?* You answer it by substituting. When substituting works, you are finished in one line. When it gives $\frac{0}{0}$ or $\frac{\infty}{\infty}$ you have learned nothing yet — and everything else in this practicum is what to do next.

Material. All of `calculus.limits` from the AA HL archive, sessions May 2021 — November 2025. Twenty-two blocks in the corpus, of which two are duplicates: November 2023 Paper 3 is one paper filed under both zones, and the November 2023 Paper 1 question on $\sin^2(kx)/x^2$ was set in TZ1 and TZ2 word for word.

This practicum is in English, like B2 to B5. The checks speak whichever

language the notebook asks them to, and this one asks for English in the setup cell.

The one thing to carry out of here.

Substitute **first**, always. If it gives a number, that is the limit and you are done. If it gives $\frac{0}{0}$ or $\frac{\infty}{\infty}$, you have a licence to differentiate — and not before.

That order is not pedantry, it is where the marks are. May 2025 TZ2 is worth five marks for a limit whose value is 0, and the first of them is for rewriting $(3x+1)e^{-3x}$ as $\frac{3x+1}{e^{3x}}$ — because a product is not a quotient, and l'Hôpital's rule applies to quotients. The markscheme says so in as many words: *"This first A1 must be seen."*

Where the calculator sits, and here it barely matters. 44% of the marks carry one, and almost none of them use it. The answers in this topic are $\frac{2}{3}$, -3 , $-\frac{1}{4}$, $\frac{\pi}{4}$, $-\frac{n}{2}$, $\frac{1}{2}n(n+1)$ — exact things. The two

questions that do want a number want it from an integral, not from the limit.

How the checks work here. Two of them are new, and the first one is the reason this practicum exists.

- `verify_limit` does not know the answer. It takes the expression from the question, walks in towards the point along a ladder — $10^{-1}, 10^{-2}, \dots, 10^{-8}$, from both sides — and asks whether the number you named is what those values are settling on. There is no stored constant anywhere: it cannot be wrong in the same way you are, because there is nothing for it to be wrong about.

It also catches the two standard losses. An answer with x still in it is rejected on sight — a limit is a number, not an expression. And an answer read off after one round of l'Hôpital when the form was still $\frac{0}{0}$ simply does not converge to what you wrote.

- `verify_indeterminate` is for *"show that the limit is in indeterminate form"*, which is worth its own mark twice in this archive. It looks at the numerator and at the denominator separately and tells you which way each one is actually going. Claiming $\frac{0}{0}$ where the truth is $\frac{\infty}{\infty}$ does not pass.

How to work

1. Read the map of techniques first. It is arranged by **what substituting gave you**.
2. Work **on paper**. Every limit in this topic is a Paper 1 answer even when it appears on Paper 2.
3. Exact answers throughout: `Rational(2, 3)`, `-pi/4`, `-n/2`, `n*(n+1)/2`. `oo` for an infinite limit.
4. The value of a limit is entered as a plain expression — no `lim`, no `x` in it.
5. A form is entered as a string: `'0/0'` or `'oo/oo'`.

6. The last three blocks are a whole Paper 3 part, a recognition trainer, and one question on a timer.

Difficulty marks: ● the technique on its own · ● the technique in a wrapper · ● several techniques, or a whole exam question.

```
import sys
sys.path.append('..')          # from
practicum/calculus to practicum/kit.py
import sympy as sp             # the escape
hatch: anything not in kit is in sp
from kit import *              # checks +
Interval, Rational, oo, pi, limit-free tools

language('en')                 # this
notebook is in English, and so are the checks

b, c, m, n = symbols('b c m n')
alpha = symbols('alpha')

print('ready; sympy', sp.__version__)
print('an exact limit:      ', Rational(2,
3))
print('a limit in terms of n:', -n/2)
print('an infinite limit:   ', oo)
print("a form, as a string: ", '0/0')
```

Map of techniques

#	Technique	Trigger in the question	First move
1	Substitute, and cancel	the point is in the domain, or $\frac{0}{0}$ from a shared factor	put the number in; factorise if it stops you
2	Highest power	$x \rightarrow \infty$ and no exponentials	divide top and bottom by it
3	l'Hôpital once	$\frac{0}{0}$ or $\frac{\infty}{\infty}$, nothing to cancel	differentiate top and bottom separately
4	l'Hôpital again	the denominator is x^3, x^4 , a high power	check the form again before repeating
5	Maclaurin instead	$e, \sin, \cos, \ln$ around 0	replace each factor by its series
6	Choose the constant	a letter in the question, «a finite limit only exists»	force the numerator to vanish
7	Limit in a parameter	$n \rightarrow \infty$ or $\alpha \rightarrow \frac{\pi}{2}$, with x held fixed	ask what survives
8	The rule with a letter	$x^{n+2}, S_n(x)$ — a parameter	differentiate in x , carrying n

#	Technique	Trigger in the question	First move
	in it	inside the expression	along
9	What it means	«state a reason», «explain the significance»	answer about the model, not the algebra

The ladder goes by what substituting gave you.

Rungs 1-2 — substituting settled it.

Either the point is in the domain and the limit is the value, or the $\frac{0}{0}$ came from a factor you can cancel, or the point is at infinity and only the highest powers matter. None of these needs calculus. Do not reach for l'Hôpital's rule here: it would give the right answer, but slower, and on a question worth one mark that is the whole of the time you had.

Rungs 3-5 — substituting gave $\frac{0}{0}$ and there is nothing to cancel. This is the middle of the topic and nine of the twenty blocks. Rung 3 is the rule itself. Rung 4 is the discipline around it: after

differentiating, *look again* — the form is often still $\frac{0}{0}$, and the markscheme for May 2024 will not give you the mark unless the second application is justified. Rung 5 is the alternative the Paper 1 questions are really built for: when the expression is made of e^x , $\sin$, $\cos$ around zero, its Maclaurin series answers in one line what l'Hôpital answers in three.

Rungs 6-8 — the limit is not quite the question. Rung 6 hands you a letter and tells you the limit is finite; you work backwards to the letter. Rung 7 takes the limit in something other than x — in n , in α — while x sits there unmoved, and the answer is an expression in the letters that stayed. Rung 8 keeps the limit in x but leaves a parameter inside the expression, so the differentiation carries n along and the answer is a formula: $\frac{1}{2}n(n+1)$, or simply 1 for every n at once.

Rung 9 — the sentence at the end. Two marks in the archive are for saying what the limit means: that a sprinter's

speed never reaches 8.14 m s^{-1} because the race ends first, that a parabola looks straight from far enough away. They are the easiest marks in the topic and the ones most often left blank.

What saves the most time. Write $\lim_{x \rightarrow a}$ on every line. It looks like decoration and it is not: May 2021 caps the question at **M1A1A0M1A1** — four marks out of five — for a perfect solution with the symbol missing, and November 2022 refuses the mark for the indeterminate form outright if $\lim$ is not written.

Part I — when substituting is enough

Theory 1. What the notation asks, and why substituting comes first

$$\lim_{x \rightarrow a} f(x) = L$$

says: *as x gets close to a , $f(x)$ gets close to L* — and it says nothing whatever about $f(a)$. That gap is the whole subject. $f(a)$ may be a different number, or may not exist at all, and the limit is fine either way.

But when the function is continuous at a , the gap closes and $\lim_{x \rightarrow a} f(x) = f(a)$. Every function you meet in this course — polynomials, e^x , $\sin$, $\cos$, $\ln$ on its domain, and sums, products and quotients of them where the denominator is not zero — is continuous. So *substitute first* is not a shortcut. It is the theorem.

The $\frac{0}{0}$ you get is not an answer, and not a failure either. It is a diagnosis: the numerator and denominator are both dying at a , and the limit is a statement about *which of them dies faster*. Nothing about $\frac{0}{0}$ tells you the answer — it can come out to 0, to $\frac{2}{3}$, to -3 , or to ∞ . That is exactly why it is called *indeterminate*.

The first cure: they are dying because of a shared factor. Take

$$g(x) = \frac{x^3 - 8}{x^2 - 4} \quad \text{at } x = 2.$$

Substituting gives $\frac{0}{0}$. But $x^3 - 8 = (x - 2)(x^2 + 2x + 4)$ and $x^2 - 4 = (x - 2)(x + 2)$, so

$$g(x) = \frac{x^2 + 2x + 4}{x + 2} \quad \text{for } x \neq 2,$$

and now substituting works: the limit is $\frac{12}{4} = 3$. The cancelled factor is exactly the shared cause of death. Note the phrase *for $x \neq 2$* : the two functions are not equal at 2, and that is precisely why the limit exists where the value does not.

Factorising the top is the whole of the work, and it is the work the next task asks for before it asks for anything else.

Task 1 — **substitute, and cancel what stops you**

Consider the polynomials

$$P(x) = 3x^3 + 5x^2 + x - 1,$$

$$Q(x) = (x + 1)(2x + 1),$$

and the function

$$f(x) = \frac{P(x)}{(x + 1)Q(x)}, \quad x \neq -1, x \neq -\frac{1}{2}.$$

(a) Express $P(x)$ as a product of three linear factors.

(b) Find $\lim_{x \rightarrow -1} f(x)$.

The check for (b) has no stored answer. It takes f as the question defines it, walks in towards -1 from both sides, and asks whether your number is what the values are settling on.

May 2024 TZ1 Paper 1 Q11(b) and Q11(f)(i) — 3 marks and part of 3.

```
FX = (3*x**3 + 5*x**2 + x - 1) / ((x + 1) *  
(x + 1) * (2*x + 1)) # f as given
```

```
qla = ... # the cubic as a product of  
three linear factors
```

```
qlb = ... # the limit as x -> -1
```

```
verify_factored('la', qla, 3*x**3 + 5*x**2 +  
x - 1, n=3)
```

```
verify_limit('lb', qlb, FX, point=-1)
```

Theory 2. At infinity: only the highest power is alive

$x \rightarrow \infty$ is not a point you can substitute into, so the rule is different — but it is just as mechanical. **Divide the numerator and the denominator by the highest power of x that appears,** and then use $\frac{1}{x^k} \rightarrow 0$.

$$\frac{5x + 2}{4x - 7} = \frac{5 + \frac{2}{x}}{4 - \frac{7}{x}} \longrightarrow \frac{5 + 0}{4 - 0} = \frac{5}{4}.$$

Everything except the leading terms has died. That is the whole of the technique, and it gives the three cases at once:

top vs bottom	example	limit
same degree	$\frac{5x+2}{4x-7}$	ratio of leading coefficients, $\frac{5}{4}$
bottom wins	$\frac{2x+5}{x^2}$	0
top wins	$\frac{x^2}{2x+5}$	∞ — no limit

Roots count as powers. A square root halves the degree under it:

$$\frac{6u}{\sqrt{4u^2 + 9}}.$$

Under the root, $4u^2 + 9$ behaves like $4u^2$, so the root behaves like $2u$: top and bottom are both degree one, and dividing through by u makes it visible,

$$\frac{6u}{\sqrt{4u^2 + 9}} = \frac{6}{\sqrt{4 + \frac{9}{u^2}}} \longrightarrow \frac{6}{2} = 3.$$

A model built out of a root behaves the same way, and the number that comes out is the value the model creeps towards but never reaches.

Exponentials are not powers, and they beat every power. e^x outgrows x^{100}

; e^{-x} dies faster than $\frac{1}{x^{100}}$. So $\frac{x^5}{e^x} \rightarrow 0$ and $\frac{e^x + x^2}{e^x} \rightarrow 1$ — but neither of those is settled by dividing by a power, and both belong to Part II.

Task 2 ● — the far end

(a) With f as in Task 1, find $\lim_{x \rightarrow \infty} f(x)$.

Two athletes run a 200 m race. Fiona's velocity, in m s^{-1} , is modelled by

$$v(t) = \frac{8.14t}{\sqrt{t^2 + 0.2}}, \quad t \geq 0.$$

(b) Write down the limit of $v(t)$ as t approaches infinity.

May 2024 TZ1 Paper 1 Q11(f)(ii) — part of 3 marks. May 2025 TZ3 Paper 2 Q10(c)(i) — part of 3 marks.

```
VT = Rational(814, 100)*t / sqrt(t**2 +
Rational(2, 10))
```

```
q2a = ...           # the limit of f as x -> oo
```

```
q2b = ...           # the limit of v as t -> oo
```

```
verify_limit('2a', q2a, FX, point=oo)
```

```
verify_limit('2b', q2b, VT, var=t, point=oo)
```

Theory 3. The sentence at the end: what a limit means

Two of the archive's marks are not for a number at all. They are for one sentence about what the number means, and both are worth as much as a line of algebra.

A limit the model never reaches. A cup of coffee cooling in a room is modelled by $T(t) = 18 + 62e^{-0.06t}$ degrees, and $T(t) \rightarrow 18$ as $t \rightarrow \infty$. Asked *why is this value not attained*, the answer is not about the exponential: the coffee is drunk, or thrown away, long before then, and in any case it can never be colder than the room it is standing in. The limit describes the formula; the formula stops describing the coffee at some point the formula does not know about.

A limit that says what a shape looks like from far away. The gradient of $y = \sqrt{x^2 + 1}$ is

$$\frac{dy}{dx} = \frac{x}{\sqrt{x^2 + 1}} \longrightarrow 1,$$

and the sentence that earns the second mark is what that means about the picture: far out to the right the curve has settled into the direction of the line $y = x$, which is its asymptote. The number is one mark; the sentence about the shape is the other.

How to write these. Say what happens to the *thing being modelled*, not to the algebra. "The denominator grows without bound" earns nothing. "The race is over long before then" earns the mark.

Task 3 — the number and the sentence

Fiona's velocity is $v(t) = \frac{8.14t}{\sqrt{t^2 + 0.2}}$, as in Task 2, in a 200 m race.

(a) State a reason why the value in Task 2(b) is not valid in the context of this question.

For a quadratic $h(x) = ax^2 + bx + c$ with $a \neq 0$, the curvature is

$$k(x) = \frac{2|a|}{(1 + (2ax + b)^2)^{3/2}}.$$

(b) State the value of $\lim_{x \rightarrow \infty} k(x)$.

(c) Explain briefly the significance of this result.

Parts (a) and (c) are sentences, and no cell can mark a sentence. Write them in the markdown cell below in one line each, then compare with the solution — the point is whether you talked about the runner and the parabola or about the algebra.

May 2025 TZ3 Paper 2 Q10(c)(ii) — part of 3 marks. May 2025 TZ1 Paper 3 Q2(b)(iii) — 2 marks.

```
a = symbols('a')
KX = 2*abs(a) / (1 + (2*a*x +
b)**2)**Rational(3, 2)

q3b = ...           # the limit of the curvature
as x -> oo

verify_limit('3b', q3b, KX, point=oo, params=
{a: (2, -3), b: (5, 0)})
```

(a) *your sentence here*

(c) *your sentence here*

Part II — when substituting gives $\frac{0}{0}$

Theory 4. l'Hôpital's rule, and the licence it needs

If $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$ has the form $\frac{0}{0}$ or $\frac{\infty}{\infty}$, then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

whenever the limit on the right exists.

Three things about it that cost marks.

It is not the quotient rule. You differentiate the top, you differentiate the bottom, and you put one over the other. Nothing is subtracted, nothing is squared.

The licence has to be shown. The rule is only true for those two forms. Substituting first is what earns the right to use it, and the markschemes are explicit: November 2022 gives a mark

purely for writing down the substitution that produced the $\frac{0}{0}$, with the $\lim$ symbol in front of it, and refuses the mark when the symbol is missing.

It applies to quotients only. A *product* of the form $\infty \cdot 0$ is not covered until it has been turned into a quotient:

$$x \sin \frac{1}{x} = \frac{\sin \frac{1}{x}}{\frac{1}{x}} \quad \text{— now it is } \frac{0}{0} \text{ as } x \rightarrow \infty,$$

and in questions of this shape the markscheme awards its first mark for that rewrite alone.

A worked one, in full.

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\sin 5x}{\arcsin 2x} &= \frac{\sin 0}{\arcsin 0} \\ &= \frac{0}{0} \checkmark \\ &= \lim_{x \rightarrow 0} \frac{5 \cos 5x}{\frac{2}{\sqrt{1-4x^2}}} \\ &= \frac{5 \cdot 1}{2/1} \\ &= \frac{5}{2}. \end{aligned}$$

Five marks in a question of this size: one for attempting to differentiate top and bottom, one for each derivative, one for substituting, one for the number. Write the lim on every line or the maximum drops to four.

Task 4 — the rule, once

(a) Show that $\lim_{x \rightarrow 0} \frac{\arctan 2x}{\tan 3x}$ is in indeterminate form, and name the form.

(b) Use l'Hôpital's rule to find $\lim_{x \rightarrow 0} \frac{\arctan 2x}{\tan 3x}$.

For (a) the form is entered as a string, '0/0' or 'oo/oo'. The check looks at the numerator and the denominator separately and will tell you which way each one is really going.

May 2021 TZ1 Paper 1 Q8 — 5 marks.

```
q4a = ...           # the form, as a string:
'0/0' or 'oo/oo'
q4b = ...           # the limit

verify_indeterminate('4a', q4a, atan(2*x),
tan(3*x))
verify_limit('4b', q4b, atan(2*x)/tan(3*x))
```

Theory 5. Applying it again — and knowing when to stop

After one round you have a new quotient, and the first thing to do with it is the first thing you did with the old one: **substitute**. Three outcomes.

- **A number.** That is the limit. Stop.
- $\frac{0}{0}$ or $\frac{\infty}{\infty}$ **again.** New licence, apply the rule again.
- **Something like $\frac{-2}{-2}$ or $\frac{3}{0}$.** Not indeterminate. Stop, and read the answer off. Applying the rule here is not merely wasteful — it gives the wrong number.

Watch the third outcome happen. For

$$\lim_{x \rightarrow 0} \frac{1 - \cos 2x}{x^2}$$

one round gives $\frac{2 \sin 2x}{2x}$, still $\frac{0}{0}$; a second gives $\frac{4 \cos 2x}{2}$, and the denominator is now the number 2. That is the signal to stop and read off $\frac{4}{2} = 2$. A third round would differentiate that 2 into 0 and produce nonsense.

The markschemes say so in as many words: *M1 for second use of l'Hôpital's rule provided expression is in indeterminate $\frac{0}{0}$ form and no third attempt at using the rule.* A round too many costs the mark that the round before it earned.

How many rounds to expect, at a glance. Look at the denominator. A denominator of x^2 usually takes two rounds, x^3 three, x^4 four — each round lowers the power by one, and the limit becomes visible when the denominator finally survives substitution. So a numerator over x^3 is a three-round question before you start, and if three

rounds of differentiating a product of transcendental functions sounds like a lot — it is, and Theory 6 is the way round it.

Task 5 — twice, and stop

Use l'Hôpital's rule to find

$$\lim_{x \rightarrow 0} \frac{\sec^4 x - \cos^2 x}{x^4 - x^2}.$$

Six marks, no calculator. Differentiate carefully — $\sec^4 x$ needs the chain rule, and the whole question is lost in that one derivative.

May 2024 TZ2 Paper 1 Q8 — 6 marks.

```
q5 = ...           # the limit

verify_limit('5', q5, (sec(x)**4 -
cos(x)**2)/(x**4 - x**2))
```

Task 6 — three rounds, or one line

The function g is defined by $g(x) = e^x \cos x$, $x \in \mathbb{R}$.

(a) Find the Maclaurin series for $g(x)$ up to and including the x^4 term.

(b) Hence, or otherwise, determine the value of

$$\lim_{x \rightarrow 0} \frac{e^x \cos x - 1 - x}{x^3}.$$

Do (b) both ways if you have time — first with l'Hôpital three times, then with the series from (a). The second way is four lines and the first is a page, and knowing that in the exam is worth more than the answer.

May 2022 TZ1 Paper 1 Q12(d) and Q12(e) — 5 marks and 3 marks.

```
q6a = ...          # the Maclaurin series of
e^x cos x up to the x^4 term
q6b = ...          # the limit

check_series('6a', q6a, '6a49f54210cb')
verify_limit('6b', q6b, (exp(x)*cos(x) - 1 -
x)/x**3)
```

Theory 6. Maclaurin instead: read the leading term

Every function in this topic has a series you already know:

$$\begin{aligned} e^x &= 1 \\ &+ x \\ &+ \frac{x^2}{2!} \\ &+ \frac{x^3}{3!} \\ &+ \dots, \\ \sin x &= x - \frac{x^3}{3!} + \dots, \\ \cos x &= 1 \\ &- \frac{x^2}{2!} \\ &+ \frac{x^4}{4!} \\ &- \dots \end{aligned}$$

Near 0, a series *is* the function, and a quotient of series is settled by looking at the lowest power that survives on each side.

Why it wins. l'Hôpital differentiates the whole expression, again and again, and

every round doubles the mess. The series is built once and then you read it:

$$\begin{aligned}
 \ln(1+x) &= x \\
 &\quad - \frac{x^2}{2} \\
 &\quad + \frac{x^3}{3} \\
 &\quad - \dots \\
 \implies &\frac{\ln(1+x) - x}{x^2} \\
 = &\frac{-\frac{x^2}{2} + \dots}{x^2} \longrightarrow -\frac{1}{2}.
 \end{aligned}$$

Notice what the subtracted x was for: it cleared the term that would have survived the division and hidden the answer. Whenever a question subtracts the first few terms of a familiar series, that is what it is doing, and it tells you which power of x you are about to divide by.

The version that looks impossible and is not.

$$\lim_{x \rightarrow 0} \frac{(\sin x - x)^2}{x^6}.$$

A sixth power. But $x^6 = (x^3)^2$, so the whole thing is a square:

$$\begin{aligned}\frac{(\sin x - x)^2}{x^6} &= \left(\frac{\sin x - x}{x^3} \right)^2 \longrightarrow \left(-\frac{1}{6} \right)^2 \\ &= \frac{1}{36},\end{aligned}$$

because $\sin x - x = -\frac{x^3}{6} + \dots$ and dividing by x^3 leaves $-\frac{1}{6} + \dots$. Pull the power out first, always: one small limit raised to a power beats one enormous limit every time.

With a parameter in the exponent. A letter in a power is handled by the binomial series and nothing else. For

$\lim_{x \rightarrow 0} \frac{(1+x)^n - 1}{x}$ raise the bracket:

$$\begin{aligned}(1+x)^n &= 1 \\ &+ nx \\ &+ \frac{n(n-1)}{2}x^2 \\ &+ \dots \\ \implies \frac{(1+x)^n - 1}{x} &\longrightarrow n.\end{aligned}$$

Only the first two terms matter, because everything after them still carries an x

once the division is done. The same counting decides how many terms you need when the bracket is a series rather than $1 + x$.

Task 7 — pull the power out, keep the parameter

(a) Determine the value of $\lim_{x \rightarrow 0} \frac{(x^2 e^x - x^2)^3}{x^9}$.

Let $f_n(x) = \cos^n x$, where $x \in \mathbb{R}$ and $n \in \mathbb{N}$.

(b) Find the Maclaurin series of $f_n(x)$ up to the term in x^2 .

(c) Hence or otherwise, find $\lim_{x \rightarrow 0} \frac{f_n(x) - 1}{x^2}$ in terms of n .

The check for (c) takes your answer, puts $n = 1, 2, 5, 9$ into it in turn, and walks in on $\cos^n x$ each time. An answer that happens to work for one n will not survive the others.

November 2021 Paper 1 Q11(c) — 4 marks. May 2025 TZ1 Paper 1 Q12(e) — 5 marks.

```

q7a = ...          # the limit of (x^2 e^x -
x^2)^3 / x^9
q7b = ...          # the Maclaurin series of
cos^n x up to x^2, in terms of n
q7c = ...          # the limit of (cos^n x -
1)/x^2, in terms of n

verify_limit('7a', q7a, (x**2*exp(x) -
x**2)**3 / x**9)
check_series('7b', q7b, '281b04dc414d')
verify_limit('7c', q7c, (cos(x)**n - 1)/x**2,
params={n: (1, 2, 5, 9)})

```

Theory 7. $\frac{\infty}{\infty}$, and the forms that must be rearranged

The rule takes $\frac{\infty}{\infty}$ on exactly the same terms as $\frac{0}{0}$, and at infinity that is usually the form you meet.

A worked one. Both parts of

$$\lim_{x \rightarrow \infty} \frac{x^2}{e^x}$$

run away to infinity, so the rule applies, and it applies again to what it produces:

$$\lim_{x \rightarrow \infty} \frac{x^2}{e^x} = \lim_{x \rightarrow \infty} \frac{2x}{e^x} = \lim_{x \rightarrow \infty} \frac{2}{e^x} = 0.$$

Two rounds, and the third would be the mistake: the last quotient is $\frac{2}{\infty}$, which is not indeterminate at all.

The rearrangements you must know.

The rule reads quotients. Anything else is rewritten first, and the rewrite is where the mark is.

what you have	form	rewrite as
$x^2 e^{-x}$	$\infty \cdot 0$	$\frac{x^2}{e^x}$
$x \ln x, x \rightarrow 0^+$	$0 \cdot (-\infty)$	$\frac{\ln x}{1/x}$
$\frac{1}{x} - \frac{1}{\sin x}$	$\infty - \infty$	one fraction

Put the factor that dies into the denominator, upside down. That is the whole of the trick.

And what the answer is then used for.

The limit is often not the point of the question but a step inside it. A probability density has to integrate to 1 over its whole range, so the constant in front of it is found by integrating up to a , letting $a \rightarrow \infty$, and setting the result equal to 1: the limit is what turns an

integral with a movable end into a number that an equation can use. Expect the limit to be worth two marks of five, and the rest to be the equation around it.

Task 8 ● — the other indeterminate form

(a) Use l'Hôpital's rule to find $\lim_{b \rightarrow \infty} \frac{e^b - b - 1}{e^b}$. You may assume that the condition for applying the rule has been met.

The time T , in minutes, that a spinning top is in motion is modelled by the probability density function

$$f(t) = \begin{cases} kte^{-3t}, & t \geq 0 \\ 0, & \text{otherwise} \end{cases} \quad k \in \mathbb{Z}^+,$$

and it is given that $\int_0^a f(t) dt = \frac{k}{9} [1 - (3a + 1)e^{-3a}]$.

(b) Use l'Hôpital's rule to find $\lim_{x \rightarrow \infty} (3x + 1)e^{-3x}$.

(c) Hence, by considering $\lim_{a \rightarrow \infty} \int_0^a f(t) dt$, find the value of k .

May 2023 TZ1 Paper 3 Q1(c)(i) — 2 marks. May 2025 TZ2 Paper 2 Q11(c) — 5 marks.

```
q8a = ...          # the limit as b -> oo
q8b = ...          # the limit as x -> oo
q8c = ...          # the value of k

verify_limit('8a', q8a, (exp(b) - b -
1)/exp(b), var=b, point=oo)
verify_limit('8b', q8b, (3*x + 1)*exp(-3*x),
point=oo)
check_num('8c', q8c, 1, '19581e27de7c')
```

Part III — when the limit is not the question

Theory 8. Choosing the constant that makes a limit exist

Sometimes the letter is the answer. The question hands you an unknown constant and tells you the limit is finite; you work backwards.

The reasoning, in one move. Consider

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x} - c}{x}, \quad c \in \mathbb{R}.$$

The denominator goes to 0. If the numerator went to anything other than 0, the quotient would run away to $\pm\infty$ and there would be no finite limit. So a finite limit **requires** $\frac{0}{0}$:

$$\begin{aligned}
\lim_{x \rightarrow 0} (\sqrt{1+x} - c) &= 0 \\
&\implies 1 - c \\
&= 0 \\
&\implies c \\
&= 1,
\end{aligned}$$

and only then is there a limit to compute — one round of the rule gives $\frac{1}{2\sqrt{1+x}} \rightarrow \frac{1}{2}$. Two marks for the constant, and the second of them is for evaluating the numerator at 0 rather than for the reasoning. The markscheme's warning is worth reading twice: *award M1A0 for using the value of the constant to show the limit is $\frac{0}{0}$* — that is the argument backwards, and it earns half.

The same idea from the other end.

Sometimes the form is $\frac{0}{0}$ whatever the letter is, and it is the **value** of the limit that pins the letter down. Given that

$$\lim_{x \rightarrow 0} \frac{1 - \cos(px)}{x^2} = 18, \quad p > 0,$$

two rounds (or the series $\cos u = 1 - \frac{u^2}{2} + \dots$) turn the left side into $\frac{p^2}{2}$, so $p^2 = 36$

and, since $p > 0$, $p = 6$. The negative root is there to be discarded, and discarding it out loud is part of the answer.

Task 9 — solve for the constant

Consider $\lim_{x \rightarrow 0} \frac{\arctan(\cos x) - k}{x^2}$, where $k \in \mathbb{R}$.

(a) Show that a finite limit only exists for $k = \frac{\pi}{4}$.

(b) Using l'Hôpital's rule, find the value of the limit when $k = \frac{\pi}{4}$.

Now consider $f(x) = \frac{\sin^2(kx)}{x^2}$, where $x \neq 0$ and $k \in \mathbb{R}^+$.

(c) Given that $\lim_{x \rightarrow 0} f(x) = 16$, find the value of k .

*May 2022 TZ2 Paper 2 Q7 — 8 marks.
November 2023 TZ1 Paper 1 Q9(b) — 6 marks.*

```

q9a = ...          # the value of k that makes
the limit finite
q9b = ...          # the value of the limit at
that k
q9c = ...          # the value of k in
sin^2(kx)/x^2

verify_exact('9a', q9a, pi/4)
verify_limit('9b', q9b, (atan(cos(x)) -
pi/4)/x**2)
check_num('9c', q9c, 1, '4b227777d4dd')

```

Theory 9. The limit taken somewhere other than x

x is not sacred. The archive takes limits in n , in b , in α , in m^n — and when it does, everything that is not the limiting variable simply stays where it is.

The limit taken in a second variable.

The oldest example in the course is the definition of the derivative, where the limit runs in h and x is a spectator:

$$\begin{aligned}
 \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h} &= \lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2 + h^3}{h} \\
 &= \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2) \\
 &= 3x^2.
 \end{aligned}$$

Every x survives untouched; only the h 's go to zero, and the terms that still carry one disappear with it. The answer therefore has an x in it, and that is not a failure — it is the point. Substituting a number for x before taking the limit answers a different question.

A parameter going to a finite place. A regular n -gon inscribed in a circle of radius 1 has perimeter $2n \sin \frac{\pi}{n}$. As $n \rightarrow \infty$ the polygon becomes the circle, so the perimeter must tend to 2π — and it does, because with $u = \frac{\pi}{n} \rightarrow 0$,

$$2n \sin \frac{\pi}{n} = 2\pi \cdot \frac{\sin u}{u} \longrightarrow 2\pi.$$

The rewrite is the whole trick: put the vanishing quantity in one place and read the standard limit $\frac{\sin u}{u} \rightarrow 1$ off it. Markschemes here accept 2π and refuse 6.28 — an *exact* value was asked for.

A parameter running to a place where a function blows up. When the moving letter is the one that becomes

unbounded, divide top and bottom by it and let its reciprocal go to zero instead:

$$h(x) = \frac{1 + \lambda x}{\lambda - x} = \frac{\frac{1}{\lambda} + x}{1 - \frac{x}{\lambda}} \xrightarrow{\lambda \rightarrow \infty} \frac{0 + x}{1 - 0} = x.$$

Nothing was substituted into infinity; the expression was rearranged until the only moving part was something that goes to 0. That rearrangement is the mark, and the rest is substitution.

Task 10 — let the parameter go

Consider $f^n(x) = m^n x + c \left(\frac{1 - m^n}{1 - m} \right)$, where $-1 < m < 1$.

(a) As $n \rightarrow \infty$, the family of graphs $y = f^n(x)$ approaches the graph of a straight line L . Determine the equation of L , giving your answer in terms of c and m .

Consider the vectors $\mathbf{u} = \mathbf{i} + \mathbf{j}$ and $\mathbf{v} = (\cos \frac{1}{n})\mathbf{i} + (\sin \frac{1}{n})\mathbf{j}$, where $n \in \mathbb{Z}^+$, and let θ be the angle between them.

(b) Find an expression for $\cos \theta$ in terms of n .

(c) Find the exact value of the limit approached by θ as $n \rightarrow \infty$.

For (a) the answer is the right-hand side of $y = \dots$ — an expression in c and m , with no x and no n in it.

May 2025 TZ3 Paper 3 Q1(d) — 4 marks.
November 2022 Paper 2 Q7 — 6 marks.

```
FN = m**n*x + c*(1 - m**n)/(1 - m)
COSTHETA = (cos(1/n) + sin(1/n))/sqrt(2)

q10a = ...          # the right-hand side of the
                    # equation of L
q10b = ...          # cos(theta) in terms of n
q10c = ...          # the exact limit of theta

verify_limit('l0a', q10a, FN, var=n,
             point=oo,
             params={m: (Rational(1, 2), -
Rational(3, 4)),
                    c: (5, -2), x: (3, 7)})
verify_identity('l0b', q10b, COSTHETA, var=n,
               samples=(1, 2, 3, 5))
check_num('l0c', q10c, 6, '0e0202e1d463')
```

Task 11 — a whole Paper 3 part

Consider $f(x) = 1 + x + x^2 + \cdots + x^n$, where $n \in \mathbb{Z}^+$, and let $f_1(x) = x f'(x)$. It has been shown that, for $x \neq 1$,

$$f_1(x) = \frac{nx^{n+2} - (n+1)x^{n+1} + x}{(x-1)^2}.$$

(a) Show that $\lim_{x \rightarrow 1} f_1(x)$ is in indeterminate form, and name the form.

(b) Hence, by applying l'Hôpital's rule, show that

$$\lim_{x \rightarrow 1} f_1(x) = \frac{1}{2}n(n+1).$$

Two rounds of the rule, with n sitting in the exponents the whole time — differentiate x^{n+2} as $(n+2)x^{n+1}$ and keep going. The answer is a formula you have met before: $f_1(1) = 1 + 2 + \cdots + n$, which is what the whole question was built to produce.

November 2022 Paper 3 Q1(g) — 6 marks.

```
TOP = n*x**(n + 2) - (n + 1)*x**(n + 1) + x
F1 = TOP/(x - 1)**2
```

```
q11a = ...          # the form, as a string
q11b = ...          # the limit, in terms of n
```

```
verify_indeterminate('11a', q11a, TOP, (x -
1)**2, point=1, params={n: (2, 5, 8)})
verify_limit('11b', q11b, F1, point=1,
params={n: (2, 3, 7)})
```

Task 12 — two limits taken in something else

Two families of curves F and G meet at an acute angle α . The gradient of F is $f(x, y)$, the gradient of G is $g(x, y)$, and

$$g(x, y) = \frac{f(x, y) + \tan \alpha}{1 - f(x, y) \tan \alpha}.$$

(a) By considering $\lim_{\alpha \rightarrow \pi/2} \tan \alpha$, show that for all finite $f(x, y)$,

$$\lim_{\alpha \rightarrow \pi/2} g(x, y) = -\frac{1}{f(x, y)}.$$

Let $S_1(x) = \sin x$ and $S_n(x) = \sin(S_{n-1}(x))$ for $n \geq 2$, so $S_3(x) = \sin(\sin(\sin x))$. It has been proved that

$$S'_n(x) = \cos(S_{n-1}(x)) \cos(S_{n-2}(x)) \cdots \cos(S_1(x)) \cos x.$$

(b) Use l'Hôpital's rule to show that $\lim_{x \rightarrow 0} \frac{S_n(x)}{x} = 1$ for $n \in \mathbb{Z}^+$.

For (a), enter the limit as an expression in the letter **F** standing for $f(x, y)$; the check substitutes several values of it. For (b), enter the value once — the check tries it on S_3 and on S_6 .

November 2023 Paper 3 Q2(f) — 2 marks. November 2025 TZ3 Paper 3 Q1(g) — 3 marks.

```

F = symbols('F')                                # f(x,
y), which does not move
G = (F + tan(alpha))/(1 - F*tan(alpha))

S3 = sin(sin(sin(x)))
S6 = sin(sin(sin(sin(sin(sin(x))))))

q12a = ...          # the limit of g as alpha ->
pi/2, in terms of F
q12b = ...          # the limit of S_n(x)/x as x
-> 0

verify_limit('12a', q12a, G, var=alpha,
point=pi/2,
              params={F: (2, -3, Rational(1,
5))})
verify_limit('12b (n = 3)', q12b, S3/x)
verify_limit('12b (n = 6)', q12b, S6/x)

```

Trainer: name the technique in five seconds

Twelve openings. For each one, name the technique — **do not solve anything**. On the exam the difference between a solved limit and an unsolved one is usually the first ten seconds, because every technique here is short once chosen and hopeless once mis-chosen.

Use these codes:

`substitute` · put the number in, cancelling first if you must | `infinity` · $x \rightarrow \infty$, highest power | `lhospital` · $\frac{0}{0}$ or $\frac{\infty}{\infty}$, differentiate | `again` · expect several rounds | `series` · Maclaurin instead | `finite` · find the constant that makes it exist | `parameter` · the limit is in something other than x | `symbolic` · a parameter sits inside the expression | `context` · say what it means

1. $\lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3}$

2. $\lim_{x \rightarrow \infty} \frac{5x^2 - x}{2x^2 + 7}$

3. A population is modelled by $P(t) = \frac{800}{1 + 9e^{-0.4t}}$. State $\lim_{t \rightarrow \infty} P(t)$ and explain what it says about the population.

4. $\lim_{x \rightarrow 0} \frac{e^{2x} - 1}{\sin 5x}$

5. $\lim_{x \rightarrow 0} \frac{\sin x - x \cos x}{x^3}$

6. $\lim_{x \rightarrow 0} \frac{\ln(1+x) - x + \frac{x^2}{2}}{x^3}$, given the series for $\ln(1+x)$

7. $\lim_{x \rightarrow 0} \frac{\sqrt{4+x} - a}{x}$ is finite. Find a .
8. $\lim_{n \rightarrow \infty} (m^n a + b)$, where $|m| < 1$ and a, b are fixed
9. $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\sin x}{1 + \cos x}$
10. $\lim_{x \rightarrow \infty} x \ln \left(1 + \frac{1}{x} \right)$
1. $\lim_{x \rightarrow -\infty} \frac{3x^3 + x}{x^3 - 4x^2}$
2. $\lim_{x \rightarrow 1} \frac{x^n - 1}{x - 1}$, where $n \in \mathbb{Z}^+$

```
answers = {
    1: '', 2: '', 3: '', 4: '', 5: '', 6: '',
    7: '', 8: '', 9: '', 10: '', 11: '', 12:
    '',
}
```

```
trigger_check(answers, {1: '75519932722f', 2:
'd79987051a25', 3: 'ea7792a26f40', 4:
'ff4c5d09f6a9', 5: 'b4c9e14061c2', 6:
'9cda38ea3a2b', 7: 'd4f568b800c6', 8:
'6390361dfdff', 9: '75519932722f', 10:
'ff4c5d09f6a9', 11: 'd79987051a25', 12:
'21e2db94d100'})
```

On the timer

May 2025 TZ3 Paper 1 Q9 — 6 marks.

Target: 9 minutes.

Close everything above. No calculator.

Determine the value of

$$\lim_{x \rightarrow 0} \left(\frac{x \sin x}{1 - \cos x} \right).$$

Six marks for one limit. Substituting gives $\frac{0}{0}$; the denominator is second order in x , so expect two rounds, and check the form again after the first. There is a one-line route through the series as well — if you see it, take it, and then do it the long way afterwards to check you would have got there anyway.

Attempt log. Copy this line and add a date and a time each pass.

2026-08-30 —

```
qt = ...           # the limit

verify_limit('timed', qt, x*sin(x)/(1 -
cos(x)))
```



Solutions

Read one only after you have an answer of your own — even a wrong one. Each solution names its source and the marks it carries.

Task 1 — substitute, and cancel what stops you

May 2024 TZ1 P1 Q11(b), 3 marks · Q11(f)(i), part of 3 marks

(a) $P(-1) = -3 + 5 - 1 - 1 = 0$, so $(x + 1)$ is a factor. Dividing,

$$\begin{aligned} P(x) &= (x + 1)(3x^2 + 2x - 1) \\ &= (x + 1)(3x - 1)(x + 1) \\ &= (x + 1)^2(3x - 1). \end{aligned}$$

(b) Substituting $x = -1$ into f gives $\frac{0}{0}$, so look for the shared factor. With $Q(x) = (x + 1)(2x + 1)$,

$$\begin{aligned} f(x) &= \frac{(x + 1)^2(3x - 1)}{(x + 1) \cdot (x + 1)(2x + 1)} = \frac{3x - 1}{2x + 1} \\ &\quad (x \neq -1), \end{aligned}$$

and now

$$\lim_{x \rightarrow -1} f(x) = \frac{-3 - 1}{-2 + 1} = \frac{-4}{-1} = 4.$$

No calculus needed, and none wanted: the whole question is the factorisation from (a) being used again.

Task 2 — the far end

May 2024 TZ1 P1 Q11(f)(ii) · May 2025 TZ3 P2 Q10(c)(i)

(a) Using the cancelled form, divide by x :

$$\lim_{x \rightarrow \infty} \frac{3x - 1}{2x + 1} = \lim_{x \rightarrow \infty} \frac{3 - \frac{1}{x}}{2 + \frac{1}{x}} = \frac{3}{2}.$$

(b) $\sqrt{t^2 + 0.2}$ behaves like t , so

$$v(t) = \frac{8.14t}{\sqrt{t^2 + 0.2}} = \frac{8.14}{\sqrt{1 + \frac{0.2}{t^2}}} \rightarrow 8.14.$$

"Write down" means one mark and no working — but the working above is what tells you it is 8.14 and not 0 or ∞ .

Task 3 — the number and the sentence

May 2025 TZ3 P2 Q10(c)(ii) · May 2025 TZ1 P3 Q2(b)(iii), 2 marks

(a) The race is 200 metres long. Fiona finishes in about 26 seconds, so t never approaches infinity and her velocity never reaches 8.14 m s^{-1} — it is a bound the model tends to, not a speed anybody runs at.

(b) As $x \rightarrow \infty$, $(2ax + b)^2 \rightarrow \infty$, so the denominator grows without bound while the numerator $2|a|$ is fixed:

$$\lim_{x \rightarrow \infty} k(x) = 0.$$

(c) The curvature of a quadratic tends to zero far from the vertex: for large x a parabola is very nearly a straight line. The markscheme takes "*for large positive values of x , a quadratic function behaves like a linear function*" word for word.

Task 4 — the rule, once

May 2021 TZ1 P1 Q8, 5 marks

(a) $\arctan(2 \cdot 0) = 0$ and $\tan(3 \cdot 0) = 0$, so

$$\lim_{x \rightarrow 0} \frac{\arctan 2x}{\tan 3x} = \frac{0}{0},$$

which is indeterminate and gives the licence for the rule.

(b) Differentiating top and bottom separately,

$$\begin{aligned}\frac{d}{dx} \arctan 2x &= \frac{2}{1 + 4x^2}, \\ \frac{d}{dx} \tan 3x &= 3 \sec^2 3x,\end{aligned}$$

so

$$\lim_{x \rightarrow 0} \frac{\arctan 2x}{\tan 3x} = \lim_{x \rightarrow 0} \frac{\frac{2}{1+4x^2}}{3 \sec^2 3x} = \frac{2}{3}.$$

M1 for attempting to differentiate both, **A1A1** for the two derivatives, **(M1)** for substituting $x = 0$, **A1** for $\frac{2}{3}$. The note is blunt: *do not condone absence of limits* — without lim on each line the maximum is **M1A1A0M1A1**.

Task 5 — twice, and stop

May 2024 TZ2 P1 Q8, 6 marks

Substituting gives $\frac{1-1}{0-0} = \frac{0}{0}$, so the rule applies.

First round. $\frac{d}{dx}(\sec^4 x - \cos^2 x) = 4\sec^4 x \tan x + 2\sin x \cos x$ and $\frac{d}{dx}(x^4 - x^2) = 4x^3 - 2x$, so

$$\lim_{x \rightarrow 0} \frac{4\sec^4 x \tan x + 2\sin x \cos x}{4x^3 - 2x}.$$

At $x = 0$ this is $\frac{0}{0}$ again — a second licence.

Second round. Differentiating $4\sec^4 x \tan x$ by the product rule gives $16\sec^4 x \tan^2 x + 4\sec^6 x$, and $2\sin x \cos x = \sin 2x$ gives $2\cos 2x = 2\cos^2 x - 2\sin^2 x$:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{16\sec^4 x \tan^2 x + 4\sec^6 x - 2\sin^2 x + 2\cos^2 x}{12x^2 - 2} &= \frac{0 + 4 - 0 + 2}{-2} \\ &= \frac{6}{-2} \\ &= -3. \end{aligned}$$

At $x = 0$ the denominator is -2 , not 0: the form is no longer indeterminate and the work stops here. The markscheme is explicit — **M1** for the second use of the

rule providing their expression is in indeterminate form as $x \rightarrow 0$ and providing there is no third attempt at using l'Hôpital's rule.

Its other note is the one that quietly costs marks: to award full marks limit notation $\lim_{x \rightarrow 0}$ must be seen at least once in the working. If no limit notation is seen but otherwise all correct, do not award the final A1.

Where the six go: **M1** for attempting the rule, **A1A1** for the two first derivatives, **M1** for the second application, **A1A1** for the second derivatives — and the answer -3 falls out of them.

Task 6 — three rounds, or one line

May 2022 TZ1 P1 Q12(d), 5 marks ·
Q12(e), 3 marks

(a) With $g(x) = e^x \cos x$: $g(0) = 1$, $g'(0) = 1$, and the relation $g'' = 2(g' - g)$ from part (c) of the original question gives $g''(0) = 0$, $g'''(0) = -2$, $g^{(4)}(0) = -4$. So

$$\begin{aligned}
& g(x) && = 1 \\
& + x \\
& + 0 \cdot \frac{x^2}{2} \\
& - \frac{2x^3}{6} \\
& - \frac{4x^4}{24} \\
& = 1 + x - \frac{x^3}{3} - \frac{x^4}{6}.
\end{aligned}$$

(b) Now the limit is one subtraction:

$$\begin{aligned}
\frac{e^x \cos x - 1 - x}{x^3} &= \frac{-\frac{x^3}{3} - \frac{x^4}{6} + \dots}{x^3} \\
&= \\
&\quad -\frac{1}{3} \\
&\quad -\frac{x}{6} \\
&\quad + \dots \longrightarrow \\
&\quad -\frac{1}{3}.
\end{aligned}$$

Three marks. The l'Hôpital route reaches the same place after three rounds — the markscheme prints it as METHOD 2 and it runs to half a page.

Task 7 — pull the power out, keep the parameter

November 2021 P1 Q11(c), 4 marks

May 2025 TZ1 P1 Q12(e), 5 marks

(a) $x^9 = (x^3)^3$, so the whole expression is a cube:

$$\begin{aligned}\frac{(x^2 e^x - x^2)^3}{x^9} &= \left(\frac{x^2(e^x - 1)}{x^3} \right)^3 \\ &= \left(\frac{e^x - 1}{x} \right)^3.\end{aligned}$$

And $\frac{e^x - 1}{x} = 1 + \frac{x}{2} + \dots \rightarrow 1$, so the limit is $1^3 = 1$.

(b) From $\cos x = 1 - \frac{x^2}{2} + \dots$, raising to the n th power and keeping only what survives division by x^2 :

$$\cos^n x = \left(1 - \frac{x^2}{2} + \dots \right)^n = 1 - \frac{nx^2}{2} + \dots$$

(the binomial's next term carries x^4). Equivalently, differentiate twice: $f'_n(x) = -n \cos^{n-1} x \sin x$ and $f''_n(x) = -n \cos^n x + n(n-1) \cos^{n-2} x \sin^2 x$, so $f_n(0) = 1$, $f'_n(0) = 0$, $f''_n(0) = -n$.

(c)

$$\frac{f_n(x) - 1}{x^2} = \frac{-\frac{nx^2}{2} + \dots}{x^2} \longrightarrow -\frac{n}{2}.$$

The markscheme's note is worth having: *do not award FT marks for an expression that does not involve n* . An answer without n in it is not an answer to this question.

Task 8 — the other indeterminate form

May 2023 TZ1 P3 Q1(c)(i), 2 marks · May 2025 TZ2 P2 Q11(c), 5 marks

(a) Both parts run to infinity, so the form is $\frac{\infty}{\infty}$:

$$\lim_{b \rightarrow \infty} \frac{e^b - b - 1}{e^b} = \lim_{b \rightarrow \infty} \frac{e^b - 1}{e^b} = \lim_{b \rightarrow \infty} \frac{e^b}{e^b} = 1.$$

One mark for the correct quotient, one for the 1.

(b) A product of the form $\infty \cdot 0$ is not a quotient. Rewrite first — this is the mark the markscheme insists on seeing:

$$\begin{aligned}
(3x + 1)e^{-3x} &= \frac{3x + 1}{e^{3x}} \\
&\Rightarrow \lim_{x \rightarrow \infty} \frac{3}{3e^{3x}} \\
&= \lim_{x \rightarrow \infty} \frac{1}{e^{3x}} \\
&= 0.
\end{aligned}$$

(c) Take the limit through the given integral:

$$\begin{aligned}
\lim_{a \rightarrow \infty} \int_0^a f(t) dt &= \frac{k}{9} \left(1 - \lim_{a \rightarrow \infty} (3a + 1)e^{-3a} \right) \\
&= \frac{k}{9} (1 - 0) \\
&= \frac{k}{9}.
\end{aligned}$$

A probability density function integrates to 1 over its whole range, so $\frac{k}{9} = 1$ and $k = 9$.

Task 9 — solve for the constant

May 2022 TZ2 P2 Q7, 8 marks

November 2023 TZ1 P1 Q9(b), 6 marks

(a) $\lim_{x \rightarrow 0} x^2 = 0$. A finite limit therefore requires the numerator to go to 0 as well

— otherwise the quotient is unbounded.
So

$$\begin{aligned}\lim_{x \rightarrow 0} (\arctan(\cos x) - k) &= 0 \\ &\implies \arctan 1 - k \\ &= 0 \\ &\implies k \\ &= \frac{\pi}{4}.\end{aligned}$$

Note the direction: you deduce k from the requirement, you do not substitute k and check. **M1A0** for the second.

(b) With $k = \frac{\pi}{4}$ the form is $\frac{0}{0}$. First round:

$$\lim_{x \rightarrow 0} \frac{-\sin x}{1 + \cos^2 x} = \lim_{x \rightarrow 0} \frac{-\sin x}{2x(1 + \cos^2 x)},$$

still $\frac{0}{0}$. Second round, or simply $\frac{\sin x}{x} \rightarrow 1$ and $1 + \cos^2 x \rightarrow 2$:

$$\longrightarrow \frac{-1}{2 \cdot 2} = -\frac{1}{4}.$$

(c)

$$\frac{\sin^2 kx}{x^2} = \left(\frac{\sin kx}{x} \right)^2 = k^2 \left(\frac{\sin kx}{kx} \right)^2 \longrightarrow k^2.$$

So $k^2 = 16$, and $k \in \mathbb{R}^+$ leaves $k = 4$. Saying why -4 is rejected is part of the answer.

Task 10 — let the parameter go

May 2025 TZ3 P3 Q1(d), 4 marks
November 2022 P2 Q7, 6 marks

(a) Only m^n moves, and $-1 < m < 1$ makes $m^n \rightarrow 0$. Hence $m^n x \rightarrow 0$ and $\frac{1 - m^n}{1 - m} \rightarrow \frac{1}{1 - m}$, so

$$L: y = \frac{c}{1 - m}.$$

(M1) for using $m^n \rightarrow 0$, **(A1)** for the x term vanishing, **A1** for the constant term, **A1** for the line.

(b) $\mathbf{u} \cdot \mathbf{v} = \cos \frac{1}{n} + \sin \frac{1}{n}$, $|\mathbf{u}| = \sqrt{2}$, $|\mathbf{v}| = 1$, so

$$\cos \theta = \frac{\cos \frac{1}{n} + \sin \frac{1}{n}}{\sqrt{2}}.$$

(c) $\frac{1}{n} \rightarrow 0$, so $\cos \theta \rightarrow \frac{1+0}{\sqrt{2}} = \frac{1}{\sqrt{2}}$ and

$$\theta \longrightarrow \frac{\pi}{4}.$$

45° is accepted; 0.785 is not, because the question said *exact*.

Task 11 — a whole Paper 3 part

November 2022 P3 Q1(g), 6 marks

(a) At $x = 1$ the numerator is $n \cdot 1 - (n + 1) \cdot 1 + 1 = 0$ and the denominator is 0, so

$$\lim_{x \rightarrow 1} f_1(x) = \frac{n - (n + 1) + 1}{0} = \frac{0}{0}.$$

One mark, **R1**, and the note says it is refused if $\lim_{x \rightarrow 1}$ is not written down.

(b) First round:

$$\lim_{x \rightarrow 1} \frac{n(n + 2)x^{n+1} - (n + 1)^2x^n + 1}{2(x - 1)}.$$

At $x = 1$ the numerator is $n(n + 2) - (n + 1)^2 + 1 = n^2 + 2n - n^2 - 2n - 1 + 1 = 0$, so it is $\frac{0}{0}$ again. Second round:

$$\begin{aligned} & \lim_{x \rightarrow 1} \frac{n(n + 1)(n + 2)x^n - n(n + 1)^2x^{n-1}}{2} \\ &= \frac{n(n + 1)[(n + 2) - (n + 1)]}{2} = \frac{n(n + 1)}{2}. \end{aligned}$$

Which is $1 + 2 + \cdots + n$ — and it had to be, because $f_1(1) = 1 + 2 \cdot 1 + \cdots + n \cdot 1$.

Task 12 — two limits taken in something else

November 2023 P3 Q2(f), 2 marks

November 2025 TZ3 P3 Q1(g), 3 marks

(a) As $\alpha \rightarrow \frac{\pi}{2}$, $\tan \alpha$ is unbounded, so divide top and bottom by it and let $\frac{1}{\tan \alpha} \rightarrow 0$:

$$g = \frac{\frac{f}{\tan \alpha} + 1}{\frac{1}{\tan \alpha} - f} \longrightarrow \frac{0 + 1}{0 - f} = -\frac{1}{f}.$$

Two marks: **M1** for the division, **R1** for $\frac{1}{\tan \alpha} \rightarrow 0$, and the **R1** depends on the **M1**.

(b) $S_n(0) = 0$ and the denominator is 0, so the form is $\frac{0}{0}$. One round of the rule:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{S_n(x)}{x} &= \lim_{x \rightarrow 0} \frac{S'_n(x)}{1} \\ &= \lim_{x \rightarrow 0} \cos(S_{n-1}(x)) \cdots \cos(S_1(x)) \cos x. \end{aligned}$$

Every $S_k(0) = 0$, so every factor is $\cos 0 = 1$, and the product of n ones is 1. The proof by induction in the previous part is what makes this one line long.

On the timer

May 2025 TZ3 P1 Q9, 6 marks

Substituting gives $\frac{0.0}{1-1} = \frac{0}{0}$. First round:

$$\lim_{x \rightarrow 0} \frac{\sin x + x \cos x}{\sin x},$$

still $\frac{0}{0}$. Second round:

$$\lim_{x \rightarrow 0} \frac{2 \cos x - x \sin x}{\cos x} = \frac{2 - 0}{1} = 2.$$

The one-line route: $\sin x \approx x$ and $1 - \cos x \approx \frac{x^2}{2}$, so

$$\frac{x \sin x}{1 - \cos x} \approx \frac{x \cdot x}{x^2/2} = 2.$$

Or, without any series at all, multiply above and below by $1 + \cos x$:

$$\begin{aligned} \frac{x \sin x (1 + \cos x)}{1 - \cos^2 x} &= \frac{x \sin x (1 + \cos x)}{\sin^2 x} \\ &= \frac{x}{\sin x} (1 + \cos x) \rightarrow 1 \cdot 2 \\ &= 2. \end{aligned}$$

Three routes, one answer, and the third is the one to have on a no-calculator paper.

Key to the recognition drill

#	code	why
1	substi tute	$x^2 - 9 = (x - 3)(x + 3)$; cancel the shared factor and put 3 in
2	infini ty	same degree top and bottom: ratio of leading coefficients, $\frac{5}{2}$
3	context	the number is 800; the mark is for saying it is the ceiling the population never quite reaches
4	lhopit al	$\frac{0}{0}$, nothing to cancel, one round gives $\frac{2}{5}$
5	again	denominator x^3 and no series offered: expect three rounds, ending at $\frac{1}{3}$
6	series	the series is handed to you; use it rather than differentiating three times
7	finite	the denominator dies, so the numerator must: $a = 2$
8	parame ter	the limit is in n ; $m^n \rightarrow 0$ and a, b sit still, so the answer is b
9	substi tute	$\frac{\pi}{4}$ is in the domain and nothing vanishes; substitute and stop
10	lhopit al	$\infty \cdot 0$: rewrite as $\frac{\ln(1 + 1/x)}{1/x}$ first, then the rule
11	infini ty	degree three over degree three, so 3 — and $-\infty$ changes nothing

#	code	why
12	symbol ic	$\frac{0}{0}$ with n in the exponent; one round of the rule gives n

Numbers 5 and 6 are the pair to look at twice. Both have a high power in the denominator; the difference is that 6 hands you the series and 5 does not. When a series is given, the question is telling you which technique it wants.

Where the marks went, across the topic

technique	blocks	marks
substitute, and cancel	1	part of 3
highest power	2	part of 6
l'Hôpital, one round	3	8
l'Hôpital, more than one	3	18
Maclaurin instead	3	9
choose the constant	3	10
limit in a parameter	3	9
the rule with a letter in it	3	9
what it means	2	part of 5

Rows three to five are 35 of the 73 marks and they are one instruction repeated: **substitute, name the form, differentiate, substitute again.** The topic is small and it is nearly all one habit. The surprise is row seven — nine marks for limits that have nothing to do with x — and it is where a student who has only practised $\frac{0}{0}$ loses the most.