

全国大学生数学竞赛非数学类模拟七

清疏竞赛考研数学

2023 年 10 月 9 日

摘要

$$\mathbb{N} = \{1, 2, \dots\}, \mathbb{N}_0 = \{0, 1, 2, \dots\}.$$

模拟试题应当规定时间独立完成并给予反馈.

1 填空题

填空题 1.1 设 $f(x)$ 连续且 $f(0) \neq 0$, 则 $\lim_{x \rightarrow 0} \frac{\int_0^x (x-t)f(t)dt}{x \int_0^x f(x-t)dt} = \underline{\frac{1}{2}}$

填空题 1.2 设函数 $y = y(x)$ 由 $\int_0^x x \sin t^2 dt = \int_0^{x+y} e^{-u^2} du$ 确定, 则 $y'(0) = \underline{-1}$

$x \int_0^x \sin t^2 dt - \int_0^{x+y} e^{-u^2} du = 0$
 $y'(0) = - \frac{\int_0^x \sin t^2 dt + x \sin x^2}{-e^{-(x+y)^2}} \Big|_{x=0} = -1$

填空题 1.3 曲面 $r(u, v) = (\sin u, \cos u \sin v, \sin v)$ 在点 $\left(\frac{1}{2}, \frac{\sqrt{3}}{4}, \frac{1}{2}\right)$ 的切平面方程为 $\underline{-\frac{5}{8}x - \frac{3}{4}y + \frac{35}{8}z = 0}$. $r(u, v) = (f(u, v), g(u, v), h(u, v))$, 则切平面方程为 $\left(\frac{\partial g}{\partial u}, \frac{\partial f}{\partial u}, \frac{\partial f}{\partial v}\right) \cdot \left(\frac{\partial g}{\partial v}, \frac{\partial f}{\partial v}, \frac{\partial h}{\partial v}\right)$

填空题 1.4 假如 z, w 是 x, y 的连续可微函数且满足 $xw^3 + yz^2 + z^3 = -1, zw^3 - xz^3 + y^2w = 1$, 在 $(x, y, z, w) = (1, -1, -1, 1)$. 计算 $\frac{\partial z}{\partial x} = \underline{-\frac{5}{4}}$.

填空题 1.5 设 $f(x) = \int_1^x \frac{\ln(1+t)}{t} dt$, 则 $\int_0^1 \frac{f(x)}{\sqrt{x}} dx = \underline{-2\pi - 4\ln 2 + 8}$

1.1 $\lim_{x \rightarrow 0} \frac{\int_0^x (x-t)f(t)dt}{x \int_0^x f(t)dt} = \lim_{x \rightarrow 0} \frac{[x \int_0^x f(t)dt - \int_0^x t f(t)dt]'}{\int_0^x f(t)dt + x f(x)}$
 $= \lim_{x \rightarrow 0} \frac{\int_0^x f(t)dt}{\int_0^x f(t)dt + x f(x)} = \lim_{x \rightarrow 0} \frac{1}{1 + \frac{x f(x)}{\int_0^x f(t)dt}} = A$
又 $\lim_{x \rightarrow 0} \frac{x}{\int_0^x f(t)dt} = \lim_{x \rightarrow 0} \frac{1}{f(x)} = \frac{1}{f(0)}$, 故 $A = \frac{1}{2}$

$$1.3. \frac{\partial(g)h}{\partial(u,v)} = \begin{vmatrix} \frac{\partial g}{\partial u} & \frac{\partial g}{\partial v} \\ \frac{\partial h}{\partial u} & \frac{\partial h}{\partial v} \end{vmatrix} = \begin{vmatrix} -\sin u \sin v & \cos u \cos v \\ 0 & \cos v \end{vmatrix} = -\sin u \sin v \cos v$$

$$\frac{\partial(f)h}{\partial(u,v)} = \begin{vmatrix} \cos u & 0 \\ 0 & \cos v \end{vmatrix} = \cos u \cdot \cos v$$

$$\frac{\partial(f)g}{\partial(u,v)} = \begin{vmatrix} \cos u & 0 \\ -\sin u \sin v & \cos u \cos v \end{vmatrix} = \cos^2 u \cos v$$


$$\text{代入 } (u,v) = \left(\frac{\pi}{6}, \frac{\pi}{6}\right), \text{ 故 } \left(-\frac{\sqrt{3}}{8}, \frac{3}{4}, \frac{3\sqrt{3}}{8}\right)$$

$$1.4 \quad \begin{cases} xw^3 + yz^2 + z^3 = -1 \\ zw^3 - xz^3 + y^2w = 1 \end{cases} \Rightarrow \begin{cases} w^3 + 3xw^2 \frac{\partial w}{\partial x} + 2yz \frac{\partial z}{\partial x} + 3z^2 \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial x} w^3 + 3zw^2 \frac{\partial w}{\partial x} - z^3 - 3xz^2 \frac{\partial z}{\partial x} + y^2 \frac{\partial w}{\partial x} = 0 \end{cases}$$

$$\text{即 } \begin{cases} 1 + 3 \frac{\partial w}{\partial x} + 2 \frac{\partial z}{\partial x} + 3 \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial x} - 3 \frac{\partial w}{\partial x} + 1 - 3 \frac{\partial z}{\partial x} + \frac{\partial w}{\partial x} = 0 \end{cases} \Rightarrow \begin{cases} 5a + 3b = -1 \\ -2a - 2b = -1 \end{cases}$$

$$\begin{cases} 10a + 6b = -2 \\ -6a - 6b = -3 \end{cases} \Rightarrow 4a = -5 \Rightarrow a = -\frac{5}{4}$$

$$1.5 \quad \int_0^1 \frac{f(x)}{\sqrt{x}} dx = \int_0^1 \int_1^x \frac{\ln(1+t)}{t\sqrt{x}} dt dx = -\int_0^1 \int_x^1 \frac{\ln(1+t)}{t\sqrt{x}} dt dx$$



$$= -\int_0^1 dt \int_0^t \frac{\ln(1+t)}{t\sqrt{x}} dx$$

$$= -\int_0^1 2 \frac{\ln(1+t)}{t} \sqrt{t} dt$$

$$= -2 \int_0^1 \frac{\ln(1+t)}{\sqrt{t}} dt$$

$$\stackrel{t=u^2}{=} -4 \int_0^1 \frac{\ln(1+u^2)}{u} u du$$

$$= -4 \int_0^1 \ln(1+u^2) du$$

$$= -4 [u \ln(1+u^2)]_0^1 - \int_0^1 u d(\ln(1+u^2))$$


$$= \text{可求(原函数)}$$

$$= -2 - 4 \ln 2 + 8$$

3 解答题

解答题 3.1 计算 $\iiint_D 2xdV$ 这里 D 由 $5(z^2 + y^2) = x, x = 5$ 围成的有界区域.

解: 不妨考虑 $D: z = 5(x^2 + y^2)$ 和 $z = 5$ 围成 计算 $\iiint_D 2z dV$
 $z = 5r^2 = 5 \Rightarrow r = 1$



$$\iiint_D 2x dV = \int_0^{2\pi} d\theta \int_0^1 r dr \int_{5r^2}^5 2z dz$$

$$= 2\pi \int_0^1 [5^2 - (5r^2)^2] \cdot r dr$$

$$= \pi \int_0^1 [25 - 25x^2] dx$$

$$= \frac{50}{3}\pi$$

解答题 3.2 设 $\varphi, \psi \in C^1(\mathbb{R})$, $\varphi(0) = -2, \psi(0) = 1$ 且对任意一光滑闭曲线 L , 都有

$$\int_L [2x\varphi(y) + \psi(y)] dx + [x^2\varphi(y) + 2xy^2 - 2x\varphi(y)] dy = 0.$$

(1): 求 φ, ψ .

(2): 对简单曲线 $L: (0,0) \rightarrow (\pi, \frac{\pi}{2})$, 计算上述积分.

解 (1): $2x\varphi(y) + 2y^2 - 2\varphi(y) = 2x\varphi'(y) + 2\psi'(y)$

故 $2\varphi(y) = 2\varphi'(y), \quad y^2 - \varphi(y) = \psi'(y)$

故 $\varphi(y) = -2e^y, \quad \psi'(y) = y^2 + 2e^y$

$\psi(y) = \frac{y^3}{3} + 2e^y - 1$

(2): $\int_L (4xe^y + \frac{2}{3}y^3 + 4e^y - 2) dx + [-2x^2e^y + 2xy^2 + 4xe^y] dy$

$\frac{d}{dy} [-2x^2e^y + (\frac{2}{3}y^3 + 4e^y - 2)x + C(y)]$

$= -2x^2e^y + (2y^2 + 4e^y)x + C'(y)$

$= -2x^2e^y + 2xy^2 + 4xe^y$

故 $C(y) = 0$, 故 $\int_L = [-2x^2e^y + (\frac{2}{3}y^3 + 4e^y - 2)x]_{(0,0)}^{(\pi, \frac{\pi}{2})}$
 $= -2\pi^2e^{\frac{\pi}{2}} + \pi(\frac{\pi^2}{12} + 4e^{\frac{\pi}{2}} - 2)$

解答题 3.3 设 $f, g \in C^1[a, b] \cap D^2(a, b)$ 且满足 $f''(x), g''(x) \neq 0, \forall x \in (a, b)$, 证明存在 $c \in (a, b)$, 使得

$$\frac{f(b) - f(a) - (b-a)f'(a)}{g(b) - g(a) - (b-a)g'(a)} = \frac{f''(c)}{g''(c)}.$$

证明: 设 $F(x) = f(x) - f(a) - (x-a)f'(a)$, $F'(x) = f'(x) - f'(a)$

$G(x) = g(x) - g(a) - (x-a)g'(a)$, $G'(x) = g'(x) - g'(a)$

由 Cauchy 中值定理, $\frac{F(b) - F(a)}{G(b) - G(a)} = \frac{F'(c)}{G'(c)} = \frac{f'(c) - f'(a)}{g'(c) - g'(a)} = \frac{f''(c)}{g''(c)}$, $c \in (a, b)$.

因此我们完成了证明.

解答题 3.4 设 $(a_n)_{n=1}^{\infty}, (b_n)_{n=1}^{\infty}$ 是两个正数列, 证明下述结果等价.

(a): 存在正实数列 $(c_n)_{n=1}^{\infty}$, 使得 $\sum_{n=1}^{\infty} \frac{a_n}{c_n}, \sum_{n=1}^{\infty} \frac{c_n}{b_n}$ 都收敛.

(b): $\sum_{n=1}^{\infty} \sqrt{\frac{a_n}{b_n}}$ 收敛.

证明: (a) \Rightarrow (b): 设正实数列 (c_n) 使 $\sum_{n=1}^{\infty} \frac{a_n}{c_n}, \sum_{n=1}^{\infty} \frac{c_n}{b_n}$ 收敛.
 则 $\sum_{n=1}^{\infty} \sqrt{\frac{a_n}{b_n}} \leq \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{a_n}{c_n} + \frac{c_n}{b_n} \right) < +\infty$, 故 $\sum_{n=1}^{\infty} \sqrt{\frac{a_n}{b_n}}$ 收敛.

(b) \Rightarrow (a): 若 $\sum_{n=1}^{\infty} \sqrt{\frac{a_n}{b_n}}$ 收敛, 取 $c_n = \sqrt{a_n b_n}$,

则 $\sum_{n=1}^{\infty} \frac{a_n}{c_n} = \sum_{n=1}^{\infty} \sqrt{\frac{a_n}{b_n}} = \sum_{n=1}^{\infty} \frac{c_n}{b_n}$. 证毕!

解答题 3.5 让 $a_0 = 1, a_1 = \frac{1}{2}, a_{n+1} = \frac{na_n^2}{1+(n+1)a_n}, n \geq 1$, 证明 $\sum_{k=0}^{\infty} \frac{a_{k+1}}{a_k}$ 收敛并求值.

证明:
$$\frac{(1+(n+1)a_n)a_{n+1}}{a_n} = na_n = \frac{a_{n+1} + (n+1)a_{n+1}a_n}{a_n}$$

故 $\frac{a_{n+1}}{a_n} = na_n - (n+1)a_{n+1}, n \geq 1$ 且 na_n 递减有下界.

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{a_{k+1}}{a_k} &= \frac{a_1}{a_0} + \sum_{k=1}^{\infty} (ka_k - (k+1)a_{k+1}) \\ &= \frac{1}{2} + 1 \cdot a_1 - \lim_{n \rightarrow \infty} na_n \\ &= 1 - \lim_{n \rightarrow \infty} na_n \text{ 收敛.} \end{aligned}$$

故 $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 0$, 故 $\exists N \geq 1, \forall n \geq N$, 有 $a_{n+1} \leq \frac{1}{2}a_n$. 则

$$a_{n+1} \leq \frac{1}{2}a_n \leq \left(\frac{1}{2}\right)^2 a_{n+1} \leq \dots \leq \left(\frac{1}{2}\right)^{n-N+1} a_N.$$

故 $\lim_{n \rightarrow \infty} na_n = 0$. 故 $\sum_{k=0}^{\infty} \frac{a_{k+1}}{a_k} = 1$.

解答题 3.6 设 $f(x) = e^{\frac{x^2}{2}} \int_x^\infty e^{-\frac{y^2}{2}} dy$, 证明

(1): f 是 $[0, +\infty)$ 的递减函数且 $\lim_{x \rightarrow +\infty} f(x) = 0$.

(2): 试计算 $\lim_{x \rightarrow +\infty} x f(x)$.

证明:

$$(1): f'(x) = x \cdot e^{\frac{x^2}{2}} \int_x^\infty e^{-\frac{y^2}{2}} dy - e^{\frac{x^2}{2}} \cdot e^{-\frac{x^2}{2}} \\ = x e^{\frac{x^2}{2}} \int_x^\infty e^{-\frac{y^2}{2}} dy - 1$$

$$\text{证 } \int_x^\infty e^{-\frac{y^2}{2}} dy \leq \frac{1}{x} e^{-\frac{x^2}{2}},$$

$$\text{证 } f(x) = \int_x^\infty e^{-\frac{y^2}{2}} dy - \frac{1}{x} e^{-\frac{x^2}{2}} \quad f'(x) = -e^{-\frac{x^2}{2}} + \frac{e^{-\frac{x^2}{2}}}{x^2} - \frac{1}{x} \cdot e^{-\frac{x^2}{2}} \cdot (-x) \\ = \frac{e^{-\frac{x^2}{2}}}{x^2} > 0.$$

故 $f(x) \leq f(+\infty) = 0$, 故 $f(x) \leq 0$, 故 $f(x)$ 递减.

(2): 我们直接求 $\lim_{x \rightarrow +\infty} x f(x)$, 从而 $\lim_{x \rightarrow +\infty} f(x) = 0$.

$$\lim_{x \rightarrow +\infty} x f(x) = \lim_{x \rightarrow +\infty} \frac{\int_x^\infty e^{-\frac{y^2}{2}} dy}{\frac{1}{x} e^{-\frac{x^2}{2}}} = \lim_{x \rightarrow +\infty} \frac{-e^{-\frac{x^2}{2}}}{-\frac{1}{x^2} \cdot e^{-\frac{x^2}{2}} - x \cdot \frac{1}{x} e^{-\frac{x^2}{2}}} \\ = \lim_{x \rightarrow +\infty} \frac{-1}{-\frac{1}{x^2} - 1} = 1.$$