

## 第十届清疏竞赛班非数学类 25

$B_r = \{x \in \mathbb{R}^3 : \|x\| \leq r\}$ , 设  $f \in C^2(\mathbb{R}^3 - B_1)$  满足  $\Delta f = 0$ ,

若  $F(x) = \frac{1}{4\pi \|x\|^2} \iint_{\partial B_{\|x\|}} f(y) dS(y)$ , 证明  $\Delta F = 0$ .

知识点:

证明(多元微分学训练题)在球坐标下拉普拉斯算子有表达式:

$$\Delta u = \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin \varphi} \partial_\varphi (\sin \varphi \partial_\varphi) + \frac{1}{r^2 \sin^2 \varphi} \partial_\theta^2,$$

$$\text{这里 } \partial_r = \frac{\partial}{\partial r}, \partial_\varphi = \frac{\partial}{\partial \varphi}, \partial_\theta^2 = \frac{\partial^2}{\partial_\theta \partial_\theta}$$

$x = r \sin \varphi \cos \theta, y = r \sin \varphi \sin \theta, z = r \cos \varphi, \varphi \in [0, \pi], \theta \in [0, 2\pi], r \geq 0$   
并推导极坐标下  $\Delta$  的表达式.

证明:

$$\begin{aligned} \Delta F(x) &= \Delta \left( \frac{1}{4\pi \|x\|^2} \iint_{\partial B_{\|x\|}} f(y) dS(y) \right) \stackrel{y=\|x\|y}{=} \Delta \left( \frac{1}{4\pi} \iint_{\partial B_1} f(\|x\| \cdot y) dS(y) \right) \\ &= \frac{1}{4\pi} \iint_{\partial B_1} \Delta_x f(\|x\| \cdot y) dS(y) = \frac{1}{4\pi} \iint_{\partial B_1} \frac{1}{r^2} \partial_r (r^2 \partial_r f(r y)) dS(y) \\ &= \frac{1}{4\pi} \int_0^{2\pi} d\theta \int_0^\pi \frac{\sin \varphi}{r^2} \partial_r (r^2 \partial_r f(r \sin \varphi \cos \theta, r \sin \varphi \sin \theta, r \cos \varphi)) d\varphi \\ &= - \int_0^{2\pi} d\theta \int_0^\pi \sin \varphi \left[ \frac{1}{r^2 \sin \varphi} \partial_\varphi (\sin \varphi \partial_\varphi f) + \frac{1}{r^2 \sin^2 \varphi} \partial_\theta^2 f \right] d\varphi \\ &= - \int_0^{2\pi} d\theta \int_0^\pi \left[ \frac{1}{r^2} \partial_\varphi (\sin \varphi \partial_\varphi f) \right] d\varphi - \int_0^{2\pi} d\theta \int_0^\pi \frac{1}{r^2 \sin \varphi} \partial_\theta^2 f d\varphi \\ &= - \frac{1}{r^2} \int_0^{2\pi} \left( \sin \varphi \partial_\varphi f \Big|_0^\pi \right) d\theta - \int_0^\pi d\varphi \int_0^{2\pi} \frac{1}{r^2 \sin \varphi} \partial_\theta^2 f d\theta \\ &= - \int_0^\pi \frac{\partial_\theta f(r \sin \varphi \cos \theta, r \sin \varphi \sin \theta, r \cos \varphi) \Big|_0^{2\pi}}{r^2 \sin \varphi} d\varphi = 0, \end{aligned}$$

从而我们证明了  $\Delta F = 0$ .

$$0 < \rho < 1, \text{ 计算 } \frac{1}{\sqrt{2\pi(1-\rho^2)}} \iint_{\mathbb{R}^2} \max\{x, y\} e^{-\frac{x^2-2\rho xy+y^2}{2(1-\rho^2)}} dx dy$$

积累:

$$\max\{x, y\} = \frac{x+y+|x-y|}{2}, \min\{x, y\} = \frac{x+y-|x-y|}{2}$$

$$\text{做 } x+y=u, x-y=v, \text{ 即 } \begin{cases} x = \frac{u+v}{2} \\ y = \frac{u-v}{2} \end{cases}, \text{ 那么:}$$

$$x^2 - 2\rho xy + y^2 = (\rho+1)v^2 + (1-\rho)u^2, |J| = \left\| \begin{array}{cc} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{array} \right\| = \frac{1}{2}$$

$$\frac{1}{\sqrt{2\pi(1-\rho^2)}} \iint_{\mathbb{R}^2} \frac{x+y+|x-y|}{2} e^{-\frac{x^2-2\rho xy+y^2}{2(1-\rho^2)}} dx dy$$

$$= \frac{1}{4\sqrt{2\pi(1-\rho^2)}} \iint_{\mathbb{R}^2} (u+|v|) e^{-\frac{(\rho+1)v^2+(1-\rho)u^2}{2(1-\rho^2)}} du dv$$

$$= \frac{1}{4\sqrt{2\pi(1-\rho^2)}} \iint_{\mathbb{R}^2} |v| e^{-\frac{(\rho+1)v^2+(1-\rho)u^2}{2(1-\rho^2)}} du dv$$

$$= \frac{1}{4\sqrt{2\pi(1-\rho^2)}} \int_{-\infty}^{\infty} e^{-\frac{u^2}{2(1-\rho)}} du \int_{-\infty}^{\infty} |v| e^{-\frac{v^2}{2(1-\rho)}} dv$$

$$= \frac{\sqrt{1-\rho}}{2}.$$

$f \in C(\mathbb{R})$  满足  $f(x) > 0$  且对所有  $t \in \mathbb{R}$ , 都有

$\int_{-\infty}^{\infty} e^{-|t-x|} f(x) dx \leq 1$ , 证明: 对任何  $b > a$ , 都有:

$$\int_a^b f(x) dx \leq \frac{b-a}{2} + 1.$$

证明:

注意到  $\int_a^b \int_{-\infty}^{\infty} e^{-|t-x|} f(x) dx dt \leq b-a$

另外一方面, 注意到  $\int_a^b \int_{-\infty}^{\infty} e^{-|t-x|} f(x) dx dt$

$$= \int_{-\infty}^{\infty} f(x) dx \int_a^b e^{-|t-x|} dt \geq \int_a^b f(x) dx \int_a^b e^{-|t-x|} dt$$

$$= \int_a^b f(x) dx \left[ \int_a^x e^{t-x} dt + \int_x^b e^{x-t} dt \right]$$

$$= \int_a^b f(x) \left[ 2 - e^{a-x} - e^{x-b} \right] dx$$

$$\text{因此 } \int_a^b f(x) dx \leq \frac{b-a}{2} + \frac{1}{2} \int_a^b f(x) \left[ e^{a-x} + e^{x-b} \right] dx$$

$$\int_a^b e^{a-x} f(x) dx \leq \int_{-\infty}^{\infty} e^{-|a-x|} f(x) dx \leq 1, \text{ 同理 } \int_a^b e^{x-b} f(x) dx \leq 1,$$

$$\text{因此我们证明了 } \int_a^b f(x) dx \leq \frac{b-a}{2} + 1.$$

计算曲线积分

$$\oint_{\Gamma} (y^2 + z^2)dx + (z^2 + x^2)dy + (x^2 + y^2)dz$$

$$\text{其中 } \Gamma: \begin{cases} x^2 + y^2 + z^2 = 2Rx \\ x^2 + y^2 = 2rx \end{cases}, 0 < r < R, z \geq 0.$$

方向与 $z$ 轴正向符合右手螺旋法则.

证明:

由 $Stokes$ 定理, 我们知道

$$\begin{aligned} & \oint_{\Gamma} (y^2 + z^2)dx + (z^2 + x^2)dy + (x^2 + y^2)dz \\ &= \iint_S \begin{vmatrix} dydz & dzdx & dxdy \\ \partial_x & \partial_y & \partial_z \\ y^2 + z^2 & z^2 + x^2 & x^2 + y^2 \end{vmatrix} \\ &= \iint_S (2y - 2z)dydz - (2x - 2z)dzdx + (2x - 2y)dxdy \end{aligned}$$

$S$ 只要是任何 $\Gamma$ 围成的光滑曲面都是对的.

可取 $S$ :  $x^2 + y^2 + z^2 = 2Rx$ , 此时 $(x - R)^2 + y^2 + z^2 = R^2$

球面外法向量为:  $\left(\frac{x - R}{R}, \frac{y}{R}, \frac{z}{R}\right)$ , 因此

$$\begin{aligned} & \iint_S (2y - 2z)dydz - (2x - 2z)dzdx + (2x - 2y)dxdy \\ &= \iint_S (2y - 2z) \frac{x - R}{R} dS - (2x - 2z) \frac{y}{R} dS + (2x - 2y) \frac{z}{R} dS \\ &= \frac{2}{R} \iint_S (y - z)(x - R) - (x - z)y + (x - y)z dS \\ &= \frac{2}{R} \iint_S -z(x - R) + xz dS \\ &= 2 \iint_S z dS \\ &= 2 \iint_{x^2 + y^2 \leq 2rx} z \frac{R}{z} dxdy \\ &= 2R \iint_{x^2 + y^2 \leq 2rx} 1 dxdy \\ &= 2\pi r^2 R. \end{aligned}$$

$z = f(x, y)$  是区域  $D = \{(x, y) \in \mathbb{R}^2 : 0 \leq x, y \leq 1\}$  上的可微函数.

若  $f(0, 0) = 0$ , 且  $dz|_{(0,0)} = 3dx + 2dy$ , 求极限

$$\lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} f(t, u) du dt}{1 - \sqrt[4]{1 - x^4}}$$

注:

本题可以改为  $f \in C(D)$  且在  $D$  处一点可微, 那么二元函数的中值定理就不在适用.

证明:  $\frac{\partial f}{\partial x}(0, 0) = 3, \frac{\partial f}{\partial y}(0, 0) = 2$

$$\lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} f(t, u) du dt}{1 - \sqrt[4]{1 - x^4}} = \lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} f(t, u) du dt}{1 - \left(1 - \frac{1}{4}x^4 + o(x^4)\right)} = 4 \lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} f(t, u) du dt}{x^4}$$

注意到是原点局部的积分, 因此对  $t, u > 0$ , 有渐进写法:

$$f(t, u) = f(0, 0) + 3t + 2u + o(\sqrt{t^2 + u^2}),$$

$$\frac{t + u}{\sqrt{2}} \leq \sqrt{t^2 + u^2} \leq t + u, o(\sqrt{t^2 + u^2}) = o(t + u), \text{ 则}$$

$$4 \lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} (3t + 2u) du dt + o\left(\int_0^{x^2} \int_x^{\sqrt{t}} (t + u) du dt\right)}{x^4} = 4 \lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} (3t + 2u) du dt}{x^4}$$

严格写法1:

$\forall \varepsilon > 0$ , 存在  $\delta > 0$ , 当  $t, u \in (0, \delta^2)$ , 有

$$3t + 2u - \varepsilon(t + u) \leq f(t, u) \leq 3t + 2u + \varepsilon(t + u).$$

当  $x \in (0, \delta)$

$$4 \lim_{x \rightarrow 0^+} \frac{-\int_0^{x^2} \int_{\sqrt{t}}^x f(t, u) du dt}{x^4} \leq 4 \lim_{x \rightarrow 0^+} \frac{-\int_0^{x^2} \int_{\sqrt{t}}^x 3t + 2u - \varepsilon(t + u) du dt}{x^4}$$

$$= \frac{1}{5} \lim_{x \rightarrow 0^+} \frac{x^4(2\varepsilon x + 5\varepsilon - 6x - 10)}{x^4} = \frac{1}{5}(5\varepsilon - 10)$$

$$4 \lim_{x \rightarrow 0^+} \frac{-\int_0^{x^2} \int_{\sqrt{t}}^x f(t, u) du dt}{x^4} \geq -\frac{1}{5}(5\varepsilon + 10),$$

$$\text{因此由 } \varepsilon \text{ 任意性我们知 } \lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} f(t, u) du dt}{1 - \sqrt[4]{1 - x^4}} = -2.$$

方法2: 不洞察到本质, 直接怎么说怎么列.

含参积分求导公式当  $a(x), b(x), f(x, s)$  连续可微, 我们有

$$F(x) = \int_{a(x)}^{b(x)} f(x, s) ds,$$

$$F'(x) = b'(x)f(x, b(x)) - a'(x)f(x, a(x)) + \int_{a(x)}^{b(x)} \partial_x f(x, s) ds$$

$$\lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_x^{\sqrt{t}} f(t, u) du dt}{1 - \sqrt[4]{1 - x^4}}$$

$$= -4 \lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} \int_{\sqrt{t}}^x f(t, u) du dt}{x^4}$$

含参积分求导公式

$$= - \lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} f(t, x) dt}{x^3}$$

此时我们可以对  $f(t, x)$  运用可微定义的 *peano* 余项形式类似方法1证明.

或者  $\lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} f(t, x) dt}{x^3} = \lim_{x \rightarrow 0^+} \frac{x^2 f(\xi(x), x)}{x^3} = \lim_{x \rightarrow 0^+} \frac{f(\xi(x), x)}{x}$

$$f(\xi(x), x) = 3\xi(x) + 2x + o\left(\sqrt{x^2 + \xi^2(x)}\right), 0 < \xi(x) < x^2$$

因此可以知道  $-\lim_{x \rightarrow 0^+} \frac{\int_0^{x^2} f(t, x) dt}{x^3} = -2.$

设  $f(x, y)$  在  $D = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq r^2\}$  上具有二阶连续偏导数, 且

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = x^2 + y^2, \text{ 求 } \lim_{r \rightarrow 0^+} \frac{\iint_D \left( x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} \right) dx dy}{(\tan r - \sin r)^2}.$$

证明:

适用下述公式的二重积分版本.

$$\iiint_{\Omega} \nabla(x^2 + y^2) \cdot \nabla f dV = - \iiint_{\Omega} (x^2 + y^2) \Delta f dV + \iint_S (x^2 + y^2) \frac{\partial f}{\partial n} dS$$

因此

$$\begin{aligned} & \lim_{r \rightarrow 0^+} \frac{\iint_D \left( x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} \right) dx dy}{(\tan r - \sin r)^2} \\ &= 2 \lim_{r \rightarrow 0^+} \frac{- \iint (x^2 + y^2)^2 dx dy + \oint (x^2 + y^2) \left( f_x \frac{x}{r} + f_y \frac{y}{r} \right) ds}{r^6} \\ &= 2 \lim_{r \rightarrow 0^+} \frac{-2\pi \int_0^r \rho^5 d\rho}{r^6} + 2 \lim_{r \rightarrow 0^+} \frac{\oint \left( f_x \frac{x}{r} + f_y \frac{y}{r} \right) ds}{r^4} \\ &= -\frac{2\pi}{3} + 2 \lim_{r \rightarrow 0^+} \frac{\iint \Delta f dx dy}{r^4} \\ &= -\frac{2\pi}{3} + 2 \lim_{r \rightarrow 0^+} \frac{\iint (x^2 + y^2) dx dy}{r^4} \\ &= -\frac{2\pi}{3} + 4\pi \lim_{r \rightarrow 0^+} \frac{\int_0^r \rho^3 d\rho}{r^4} \\ &= \frac{\pi}{3} \end{aligned}$$

