



# Decoding m<sup>6</sup>A mRNA methylation by reader proteins in cancer

Bing Han <sup>a</sup>, Saisai Wei <sup>a</sup>, Fengying Li <sup>a</sup>, Jun Zhang <sup>a</sup>, Zhongxiang Li <sup>b</sup>, Xiangwei Gao <sup>a,\*</sup>

<sup>a</sup> Sir Run-Run Shaw Hospital, School of Public Health, Zhejiang University School of Medicine, Hangzhou, 310058, China

<sup>b</sup> Medical Research Institute, Baoan Women's and Children's Hospital, Jinan University, Shenzhen, 518133, China

## ARTICLE INFO

### Keywords:

m<sup>6</sup>A modification  
YTH domain protein  
mRNA metabolism  
Cancer  
m<sup>6</sup>A editor

## ABSTRACT

N<sup>6</sup>-methyladenosine (m<sup>6</sup>A), the most prevalent internal modification in eukaryotic mRNAs, regulates gene expression at the post-transcriptional level. The reader proteins of m<sup>6</sup>A, mainly YTH domain-containing proteins, specifically recognize m<sup>6</sup>A-modified mRNAs and regulate their metabolism. Recent studies have highlighted essential roles of m<sup>6</sup>A readers in the initiation and development of human cancers. In this review, we summarize recent findings about the biological functions of YTH domain proteins in cancers, the underlying mechanisms, and clinical implications. Gene expression reprogramming by dysregulated m<sup>6</sup>A reader proteins offers potential targets for cancer treatment, while targeted m<sup>6</sup>A editors and readers provide tools to manipulate m<sup>6</sup>A metabolism in cancers.

## 1. Introduction

Chemical modifications of RNA are critical for gene expression at the post-transcriptional level, which subsequently induce changes in biological outputs. In particular, the N<sup>6</sup>-methyladenosine (m<sup>6</sup>A) modification on RNAs, methylation at the sixth position nitrogen atom of adenosine (A), is an essential regulator because of its high abundance: it has been found in 0.1–0.4 % of messenger RNA (mRNA) adenosine residues in diverse eukaryotic cells [1–3]. Despite the early discovery in various cellular mRNAs in the 1970s, the distribution of m<sup>6</sup>A was recently clarified with the development of detecting methodologies (m<sup>6</sup>A-seq and its derivatives) [4,5]. These methods allow analyzing m<sup>6</sup>A distribution at the transcriptomic level, opening up the era of “epitranscriptomics”. Mapping of m<sup>6</sup>A in different organisms revealed a highly conserved feature of this modification. A consensus motif of m<sup>6</sup>A modification, RRACH ([G/A/U][G/A]m<sup>6</sup>AC[U/A/C]), has been identified by the m<sup>6</sup>A-seq data [5,6]. As for the exact positions, m<sup>6</sup>A is enriched around stop codons and in the 3' untranslated regions (3' UTRs) [4,5]. In addition to methodological breakthrough, the characterization of its enzyme system further brought this modification into the spotlight [7]. It is now known that the m<sup>6</sup>A modification is dynamically regulated by methyltransferases (“writers”) and demethylases (“erasers”) [8,9]. The writers, including core methyltransferase components (METTL3 and METTL14) and their cofactors (WTAP, RBM15, HAKAI, VIRMA, and ZC3H13), promote the deposition of m<sup>6</sup>A on mRNAs [10]. The reversibility of m<sup>6</sup>A relies on erasers, including

FTO and ALKBH5, leading to the removal of m<sup>6</sup>A modifications for a balanced equilibrium [7,9].

## 2. “Reader” proteins regulate the fate of m<sup>6</sup>A-modified mRNAs

m<sup>6</sup>A methylation alters Watson-Crick base pairing strength, RNA secondary structure, or protein-RNA interactions, which in turn affects gene expression by modulating almost all aspects of RNA metabolism, including processing, localization, translation, and decay [11–13]. Recognition by proteins is most widely studied mechanism, and the m<sup>6</sup>A “reader” proteins directly regulate RNA fate. Combined approaches involving RNA affinity chromatography and mass spectrometry have been used to identify several m<sup>6</sup>A readers, including YT521-B homology domain-containing proteins (YTHDF1-3 and YTHDC1-2) [5], insulin-like growth factor 2 (IGF2) mRNA-binding proteins (IGF2BP1-3) [14], FMR1 [15], and heterogeneous nuclear ribonucleoprotein (HNRNP) protein family members [16,17]. These reader proteins decode m<sup>6</sup>A and decide the metabolic fate of modified mRNAs, indicating that they are crucial in the regulation of RNA metabolism and post-transcriptional gene expression.

m<sup>6</sup>A readers process RNA at least through 3 mechanisms: selectively binding, weakening the cognate binding proteins, altering the secondary structure of RNA and RNA-protein interactions [8]. Highly conserved YTH domain proteins selectively bind to m<sup>6</sup>A modifications with high affinity and a large binding interface, further altering the secondary structure [18]. Interestingly, these m<sup>6</sup>A readers have distinguishing

\* Corresponding author.

E-mail address: [xiangweigao@zju.edu.cn](mailto:xiangweigao@zju.edu.cn) (X. Gao).

<https://doi.org/10.1016/j.canlet.2021.07.047>

Received 13 May 2021; Received in revised form 25 July 2021; Accepted 28 July 2021

Available online 30 July 2021


0304-3835/© 2021 Elsevier B.V. All rights reserved.

features of RNA processing. YTHDF1 has recently been extensively studied as a cytoplasmic m<sup>6</sup>A reader. YTHDF1 binds to m<sup>6</sup>A at exact G [G > A](m<sup>6</sup>A)C sites and promotes translation initiation by interacting with initiation factors and ribosomes [19]. As a homologous gene of YTHDF1, YTHDF2 was identified by RNA immunoprecipitation (RIP) and photoactivatable ribonucleotide crosslinking and immunoprecipitation (PAR-CLIP) experiments to have a conserved G(m<sup>6</sup>A)C[U > A] motif, and this protein plays an important role in RNA decay by recruiting the mRNAs to processing bodies (P-bodies) [20]. With similar sequence identity to YTHDF2, YTHDF3 facilitates m<sup>6</sup>A-containing mRNA translation and decay together with YTHDF1 and YTHDF2 [21, 22]. YTHDC2, another cytoplasmic YTH family member, regulates mRNA stability by interacting with the 5'-3' exoribonuclease XRN1 and enhances the translation efficiency of its targets in spermatogenesis and oogenesis [23,24]. However, IGF2BPs represent a different facet of the m<sup>6</sup>A reading process, mainly promoting mRNA stability and translation by recognizing the consensus GG(m<sup>6</sup>A)C sequence [14]. As for nuclear-specific m<sup>6</sup>A readers, YTHDC1 preferentially recognizes the GG (m<sup>6</sup>A)C sequence and mainly functions in mRNA splicing and nuclear export [25–27]. Heterogeneous nuclear ribonucleoprotein G and C

(HNRNPG/HNRNPC) are other two nuclear readers that bind m<sup>6</sup>A to regulate alternative splicing [16,28,29]. Additionally, another hnRNP family protein, HNRNPA2B1, binds to m<sup>6</sup>A in the pri-miRNA to facilitate miRNA processing by recruiting a microprocessor [17] (Fig. 1).


### 3. Manipulating YTH domain protein-regulated m<sup>6</sup>A metabolism

Recently developed m<sup>6</sup>A editors offer a way to manipulate m<sup>6</sup>A specifically in individual transcripts [30–32]. Based on CRISPR/Cas system, these editors compose a methyltransferase or a demethylase fused with a catalytically inactive Cas proteins (dCas9 or dCas13). With the help of guide RNAs (gRNAs), site-specific m<sup>6</sup>A incorporation or demethylation in specific transcripts could be achieved (Fig. 2). Studies using these editors demonstrated that single site methylation in different mRNA regions may play different roles, either promoting mRNA translation or degradation. In addition to these m<sup>6</sup>A editors, targeted m<sup>6</sup>A readers have also been developed by fusing m<sup>6</sup>A reader proteins, YTHDF1 or YTHDF2, to dCas13b [33]. The fusion proteins can target the reader to specific RNA of interest using gRNA (Fig. 2). Tethering targeted mRNA with the fusion proteins can trigger enhanced protein



**Fig. 1.** The functions of m<sup>6</sup>A reader proteins in mRNA metabolism.

The m<sup>6</sup>A reader proteins recognize m<sup>6</sup>A and determine target mRNA fate. The nuclear m<sup>6</sup>A reader proteins YTHDC1, HNRNPG, and HNRNPA2B1 regulate mRNA splicing, nuclear export, and pre-miRNA processing, while the cytosolic m<sup>6</sup>A reader proteins YTHDF1-3, YTHDC2, and IGF2BP1-3 regulate mRNA translation and decay.



**Fig. 2.** Schematic of targeted  $m^6A$  editors and readers.

Targeted  $m^6A$  editors or readers have been developed by fusing  $m^6A$  writers, erasers, or readers to dCas13b/dCas9. The fusion proteins can target the mRNAs of interest with the help of gRNA and regulate their metabolism.

production or RNA degradation. The development of dCas-based targeted  $m^6A$  editors or readers reveal the effect of individual  $m^6A$  modifications and dissect their functional roles.

#### 4. The roles of YTH domain proteins in cancer initiation and development

Abnormal epigenetic regulation contributes to cancer initiation and development, during which  $m^6A$  has gained increasing attention in cancer biology [34–36].  $m^6A$  modification affects many cancer processes through writers, erasers and readers, such as cancer initiation, progression, metastasis, therapeutic response and cancer relapse [35]. For  $m^6A$  readers, accumulated evidence has shown that YTH domain proteins have essential roles in various complicated cancer pathogenesis. Because of the various RNA fate outcomes mediated by these readers, the  $m^6A$  machinery can play either oncogenic or tumor-suppressor roles in different cancer contexts. YTHDF1/3 and YTHDC2 are usually critical in promoting oncogene translation and thus act as positive tumor markers. However, YTHDF2 is reported to have dual roles in cancers that depend on the degradation of its target mRNAs. We briefly summarized the recent studies on the functions of YTH domain proteins in various cancers (Table 1). The dysregulation of  $m^6A$  readers in human cancers highlights the potential of using these proteins as new biomarkers and therapeutic targets in clinical practice.

#### 5. Aberrant expression of YTH domain proteins in cancers

Emerging data have suggested that aberrant expression of YTH proteins in many types of cancers (Table 1) could be strongly associated with cancer progression and treatment outcomes. Many studies have shown an increase of YTHDF1 expression in cancers, indicating a positive correlation with cancer pathogenesis (Table 1). However, how YTHDF1 expression programs cells to adapt to long-term cancer environmental changes remains largely unknown. DNA copy number alterations [38] or transcriptional regulation [37] are both suspected to account for the overexpression of YTHDF1 in certain cancers. For example, in colorectal cancer (CRC), the oncogenic transcription factor c-Myc was shown to be associated with the 5' region of the transcription start site of the YTHDF1 gene, indicating that c-Myc promotes YTHDF1 expression in CRC cells. Posttranscriptional regulation is also a key part of controlling YTHDF1 expression. In the intestine, the activated Wnt/APC signaling pathway promotes YTHDF1 translation by interacting with its 5' UTR [39]. MiR-376c was reported to directly target the 3'UTR of YTHDF1 in non-small-cell lung cancer (NSCLC) cells [40]. Moreover, YTHDF1 could be regulated by USF1 and c-Myc at both the mRNA and protein levels in hepatocellular carcinoma (HCC) [49]. Despite the fact that YTHDF1 is dysregulated in cancers or at least can be a risk factor for cancer initiation and prognosis, future research needs to

identify the upstream regulators of YTHDF1.

YTHDF2 has been found to participate in many cancers and might be associated with cancer progression. Interestingly, YTHDF2 expression is markedly inconsistent: it is upregulated in some types of cancers and downregulated in others. For example, YTHDF2 is highly expressed in bladder cancer, glioblastoma, acute myeloid leukemia (AML), lung cancer, ovarian cancer and prostate cancer but expressed at low levels in gastric cancer. Furthermore, YTHDF2 can act as either an oncogene or a tumor suppressor in HCC. Similar to the situation for other  $m^6A$  readers, evidence of how YTHDF2 responds to the cancer microenvironment is still limited. Hypoxia has been identified as a key driver of low YTHDF2 expression in HCC, and induction of HIFs, such as HIF-1 $\alpha$  and HIF-2 $\alpha$ , has been proven to mediate and enhance YTHDF2 function [78,79]. Recently, miRNAs have also been revealed as regulators of YTHDF2 expression. Luciferase reporter assays and statistical analysis confirmed that YTHDF2 is a direct target gene of miR-145 in ovarian cancer [86] and miR-495/miR-493-3p in prostate cancer [88,89].

#### 6. YTH proteins in colorectal cancer

Colorectal cancer (CRC) is the third leading cause of cancer-related death in both men and women worldwide. However, many cancer cases and deaths could be prevented with appropriate screening and surveillance, which indicates the high clinical need for precise tumor biomarkers and therapeutic targets.  $m^6A$  related genes are found abundantly expressed in colon cancer, including reader protein YTHDC1 and HNRNPC [94]. Recently, most related studies have indicated YTHDF1 as a potential oncogene in CRC. Given that aberrant Wnt/ $\beta$ -catenin signaling is the driving force for intestinal tumorigenesis, some studies have focused on how YTHDF1 interacts with this pathway. The studies revealed that YTHDF1 is upregulated in CRC tissues versus normal tissues [39–41]. Deletion of the YTHDF1 gene in mouse intestinal stem cells (ISCs) induces tumor shrinkage and prolongs survival by blocking the Wnt signaling pathway [39]. In addition to playing a role in ISCs, YTHDF1 is also involved in promoting cancer stem-cell like activity by enhancing Wnt activity, which further regulates tumorigenicity in human CRC [40]. According to The Cancer Genome Atlas (TCGA) database, the expression of YTHDF1 in CRC also shows a positive correlation with an important Wnt target gene, c-Myc. Mechanistically, c-Myc activates the transcriptional activity of YTHDF1, promoting cancer cell proliferation and increasing chemosensitivity [41]. However, univariate and multivariate Cox regression analyses have clarified that YTHDF1 is a negative prognostic factor for colon cancer [42]. Studies revealed lower expression in the high-risk cancer group, indicating good value for predicting prognosis. Further studies implied that YTHDF1 identifies the  $m^6A$ -modified *ANKLE1* transcript, a tumor suppressor in CRC [106], indicating that YTHDF1 may be required for ANKLE1-regulated CRC inhibition and prevention. Similar to YTHDF1, YTHDC2 contributes to colon cancer metastasis by promoting *HIF-1 $\alpha$*  translation, suggesting that YTHDC2 is a potential biomarker for diagnosis and treatment response evaluation for colon cancer patients [107].

In CRC, YTHDF3 was identified as a novel target of YAP signaling, by which YTHDF3 selectively binds to  $m^6A$ -modified *GAS5* to promote its decay and triggers CRC proliferation and metastasis [92]. Unlike that of other YTH domain proteins, the expression of YTHDF2 was unaffected in CRC tissues compared to nontumor tissues. Nonetheless, the binding of YTHDF2 to *SOX4* mRNA can be increased to induce the degradation of *SOX4*, indicating another view of the regulation and impact of  $m^6A$  on CRC [73]. Another study shows that high expression of METTL3 promotes  $m^6A$ -modified *SOX2* mRNAs, which is subsequently recognized by IGF2BP2 and leads to inhibition of *SOX2* mRNA degradation [105]. Although these observations provide the significant signatures of YTH domain proteins in CRC progression, more details and the related mechanisms need to be uncovered.

**Table 1**  
Functional characterization of YTH domain-containing proteins in cancers.

m <sup>6</sup> A readers	Cancer	Expression	Targets and Mechanism	Role	Reference
YTHDF1	Breast cancer	Upregulated	Not mentioned.	Positive prognostic signature	[37]
		Upregulated	Positive associated with <i>CDK1</i> , <i>MKI67</i> , <i>VEGFA</i> .	Positive prognostic signature	[38]
	Colorectal cancer	Upregulated	Enhances <i>TCF7L2</i> translation.	Oncogene	[39]
		Upregulated	Increases Wnt/ $\beta$ -catenin activity.	Oncogene	[40]
		Upregulated	Not mentioned.	Oncogene	[41]
		Downregulated	Not mentioned.	Negative prognostic signature	[42]
	Gastric cancer	Unknown	Facilitates <i>ANKLE1</i> expression.	Tumor suppressor	[43]
		Unaffected	Not mentioned.	Tumor suppressive signature	[44]
		Upregulated	Not mentioned.	Positive prognostic signature	[45]
	Hepatocellular carcinoma	Upregulated	Not mentioned.	Oncogene	[46]
		Upregulated	Not mentioned.	Oncogene	[47]
		Upregulated	Promotes translation of <i>Snail</i> .	Oncogene	[48]
		Upregulated	Promotes <i>FZD5</i> mRNA translation.	Oncogene	[49]
		Unknown	Increases the stability of <i>FOXO3</i> mRNA.	Oncogene	[50]
	Lung cancer	Upregulated	Promotes YAP mRNA translation.	Oncogene	[51,52]
		Upregulated	Promotes <i>PRPF6</i> mRNA expression.	Oncogene	[53]
		Upregulated	Targets <i>CDKs</i> and <i>Keap1-Nrf2-AKR1C1</i> axis	Dual effect	[54]
		Upregulated		Positive prognostic signature	[55]
		Upregulated in tumors but downregulated in higher pathological stages	Not mentioned.	Dual effect	[56]
		Upregulated	Disrupts the Wnt/b-catenin pathway.	Positive therapeutic target	[57]
	Melanoma	Unknown	Promotes <i>HINT2</i> translation.	Oncogene	[58]
		Upregulated	Interacts with <i>CDK1/2</i> .	Positive therapeutic target	[59]
	Ovarian cancer	Upregulated	Augments <i>EIF3C</i> translation.	Oncogene	[60]
		Unaffected	Promotes <i>TRIM29</i> translation.	Oncogene	[61]
	Cervical cancer	Unknown	Induces <i>PKD4</i> expression.	Positive therapeutic target	[62]
		Upregulated	Enhances <i>HK2</i> stability.	Positive therapeutic target	[63]
	Squamous cell carcinoma	Upregulated	Not mentioned.	Positive prognostic signature	[64]
		Upregulated	Promotes c-Myc stability.	Oncogene	[65]
	Merkel Cell Carcinoma	Upregulated	Enhances <i>TFRC</i> expression.	Positive therapeutic target	[66]
		Upregulated	Activates <i>eIF3</i> translation.	Oncogene	[67]
	Abdominal aortic aneurysm	Upregulated	Not mentioned.	Unknown	[68]
	Glioma	Upregulated	Not mentioned.	Positive prognostic signature	[69]
		Upregulated	Not mentioned.	Positive prognostic signature	[70]
YTHDF2	Bladder cancer	Upregulated	Promotes <i>SETD7</i> , <i>KLF4</i> degradation.	Oncogene	[71]
	Cervical cancer	Unknown	Mediates <i>GAS5</i> RNA degradation.	Potential therapeutic target	[72]
	Colorectal cancer	Unaffected	Promotes <i>SOX4</i> mRNA degradation.	Potential therapeutic target	[73]
	Gastric cancer	Downregulated	Promotes <i>FOXC2</i> degradation.	Negative prognostic signature	[74]
	Glioblastoma	Upregulated	Stabilizes <i>MYC</i> , <i>VEGFA</i> transcripts.	Potential therapeutic target	[75]
		Upregulated	Positively correlated with PD-1, TIM-3 and CTLA-4.	Positive prognostic signature	[76]
		Upregulated	Facilitates <i>LXRA</i> , <i>HIVEP2</i> mRNA decay.	Potential prognostic signature	[77]
		Downregulated	Processes <i>IL11</i> , <i>SERPINE2</i> mRNA decay.	Potential therapeutic target	[78]
		Downregulated	Destabilizing <i>EGFR</i> mRNA.	Tumor suppressor	[79]
		Unknown	Mediates <i>SOC2</i> mRNA degradation.	Potential therapeutic target	[80]
		Upregulated	Not mentioned.	Positive prognostic signature	[81]
		Upregulated	Increases <i>OCT4</i> translation.	Potential therapeutic target	[82]
	Acute myeloid leukemia	Upregulated	Destabilizes <i>TNFRSF1B</i> mRNA.	Potential therapeutic target	[83]
	Lung cancer	Unknown	Reduces <i>PFKF</i> , <i>LDHB</i> expression.	Negative prognostic signature	[84]
		Upregulated	Facilitates <i>6PGD</i> mRNA translation.	Potential prognostic signature	[85]
	Ovarian cancer	Upregulated	Inversely correlated with miR-145.	Positive therapeutic target	[86]
	Pancreatic cancer	Unknown	Promotes <i>PER1</i> mRNA degradation.	Potential diagnostic and therapeutic target	[87]
		Upregulated	Facilitates <i>MOB3B</i> mRNA degradation.	Potential therapeutic target	[88]
		Upregulated	Not mentioned.	Positive therapeutic target	[89]
		Upregulated	Induces <i>LHPP</i> , <i>NKX3-1</i> mRNA degradation.	Potential diagnostic and therapeutic target	[90]
YTHDF3	Breast cancer	Upregulated	Positive associated with <i>CDK1</i> , <i>MKI67</i> , <i>VEGFA</i> .	Positive prognostic signature	[38]
		Upregulated	Induces <i>ST6GALNAC5</i> , <i>GJA1</i> translation.	Potential prognostic signature	[91]
	Colorectal cancer	Upregulated	Facilitates <i>GAS5</i> degradation.	Potential prognostic signature	[92]
	Hepatocellular carcinoma	Unknown	Enhancing <i>Zeb1</i> mRNA stability.	Potential therapeutic target	[93]
YTHDC1	Colorectal cancer	Upregulated	Not mentioned.	Positive therapeutic target	[94]
	Glioblastoma	Unaffected	Promotes <i>SRSF</i> decay.	Potential therapeutic target	[95]

(continued on next page)



Table 1 (continued)

m <sup>6</sup> A readers	Cancer	Expression	Targets and Mechanism	Role	Reference
YTHDC2	Hepatocellular carcinoma	Upregulated	Not mentioned.	Positive prognostic target	[96]
	Prostate cancer	Unknown	Mediates <i>Cd44v5-Luc</i> splicing.	Positive therapeutic target	[97]
	Colon cancer	Upregulated	Promotes translation of <i>HIF-1α</i> and <i>Twist1</i> .	Potential prognostic signature	[98]
IGF2BPs	Lung adenocarcinoma	Downregulated	Promotes <i>SLC7A11</i> mRNA decay.	Negative prognostic signature	[99]
	Breast cancer	Downregulated	Not mentioned.	Negative prognostic signature	[100]
		Unknown	Promotes <i>CERS6</i> mRNA stability.	Potential therapeutic signature	[101]
	Gastric cancer	Upregulated	Promotes <i>HDGF</i> mRNA stability.	Potential prognostic biomarker	[102]
	Lung cancer	Upregulated	Not mentioned.	Positive prognostic signature	[103]
	Colorectal cancer	Unknown	Enhances <i>VANG1</i> mRNA stability.	Potential therapeutic target	[104]
		Upregulated	Prevents <i>SOX2</i> mRNA degradation.	Potential biomarker	[105]

## 7. YTH proteins in liver cancer

Hepatocellular carcinoma (HCC) is the most common type of primary liver cancer [108]. The high prevalence of HCC has long been attributed to chronic viral hepatitis, while other risk factors, particularly genetic or epigenetic susceptibility, are gaining importance. Studies have shown that HCC is highly associated with abnormal m<sup>6</sup>A deposition [80,109]. Interestingly, m<sup>6</sup>A modification in these studies showed quite different effects on HCC progression: METTL3 promoted HCC via m<sup>6</sup>A modification, but METTL14 was an adverse prognostic factor in HCC patients. Thus, the final outcome of m<sup>6</sup>A modification in HCC could be determined by assessing m<sup>6</sup>A-containing mRNAs and their readers.

Recently, a correlation analysis using the TCGA database showed that YTHDF1 is significantly overexpressed in HCC and positively associated with pathology stage, implying that YTHDF1 is a potential new therapeutic and prognostic marker in HCC [46]. Another similar study revealed the independent predictive value of both YTHDF1 and METTL3 for HCC patient overall survival [47]. In addition, in vitro and in vivo evidence indicated that YTHDF1 induces the progression of migration, invasion and epithelial-mesenchymal transition (EMT) in HCC cells by regulating *Snail* mRNA translation [48]. In addition, Wnt receptor-*FZD5* is another YTHDF1 target at the translational level in HCC. YTHDF1 promotes HCC cell proliferation and metastasis by accelerating the translational output of *FZD5* mRNA in an m<sup>6</sup>A-dependent manner [49]. For HCC therapy, resistance to sorafenib (a first-line treatment for advanced HCC) remains a problem in developing individual therapeutic strategies. YTHDF1 was demonstrated to have an important role in modifying the hypoxic tumor environment by identifying *FOXO3* mRNA. This study validated that METTL3 mediates and that YTHDF1 recognizes *FOXO3* methylation, leading to inhibition of the autophagy signaling pathway to maintain sorafenib sensitivity in HCC [110]. Therefore, YTHDF1 may act as an oncogene and has the potential to serve as a positive molecular biomarker for evaluating the prognosis and treatment response of HCC patients. As a synergy factor of YTHDF1 in protein synthesis, YTHDF3 enhances the stability and lifetime of m<sup>6</sup>A-methylated *Zeb1* mRNA. Upregulated *Zeb1* mediates circ K-IAA1429 expression, leading to a robust driving force for HCC migration, invasion and EMT [93].

However, the performance of YTHDF2 in HCC is more complicated, as shown above. YTHDF2 can work as both a molecular ‘rheostat’ and an ‘accelerator’, which depends on the mRNAs it processes. When it binds to *interleukin 11 (IL11)/serpin family E member 2 (SERPINE2)* or *EGFR*, YTHDF2 may act as a tumor suppressor [78,79]. In contrast, when YTHDF2 recognizes *OCT4* mRNA, it promotes HCC development and cancer metastasis [82]. In addition, there are several other studies showing that YTHDF2 has the potential to participate in HCC-related phenotypes, such as proliferation and migration [80] and immune cell infiltration [81], but further exploration is needed to determine the specific regulatory mechanisms.

## 8. YTH proteins in lung cancer

Lung cancer is a molecularly heterogeneous disease [111]. Understanding the molecular biology of lung cancer is helpful for finding promising strategies for early prognosis and treatment. Lung cancer is classified into two types: NSCLC and small-cell lung cancer (SCLC). NSCLC is the most common type, of which lung adenocarcinoma (LUAD) is the most common subtype. Recently, METTL3 was identified as a potential therapeutic target for patients with lung cancer as it promotes the translation of oncogenes [112,113]. The functions of m<sup>6</sup>A readers in lung cancer remain largely unknown; the possible functions are discussed below.


The YAP signaling pathway plays a critical role in lung cancer, including in tumorigenesis, aggressiveness, metastasis, and resistance to drug treatment [114]. As a central component of the YAP signaling pathway, YTHDF1 is recruited to m<sup>6</sup>A-modified YAP mRNA, inducing NSCLC drug resistance and metastasis by promoting YAP translation [51,52]. Another m<sup>6</sup>A target, *PRPF6*, screened by LUAD TCGA dataset analysis, was shown to be positively expressed with YTHDF1. In this study, the authors found that YTHDF1-dependent *PRPF6* m<sup>6</sup>A methylation is a key regulator of ATTM-induced anticancer effects. The mechanism mediated by the m<sup>6</sup>A-*PRPF6*-YTHDF1 axis indicates an opportunity for overcoming the side effects caused by ATTM therapy by scavenging H2S [53]. Consistently, according to different datasets from the TCGA and Gene Expression Omnibus (GEO), YTHDF1 is overexpressed in LUAD patients versus normal controls [115]. In addition, YTHDF1, together with 5 other m<sup>6</sup>A-related genes, was screened to build a risk scoring signature that was strongly related to pathological stage, sex and overall survival, indicating that YTHDF1 has good predictive value in LAUD [56]. Additionally, m<sup>6</sup>A modification was found to participate in lung cancer cell metabolism via its reader YTHDF2 [85]. In this study, upregulated YTHDF2 bound to 6-phosphogluconate dehydrogenase (*6PGD*) and promoted *6PGD* mRNA translation, indicating that YTHDF2 is a lung cancer promoter. IGF2BPs are also detected abnormally upregulated in LAUD tissue [103]. After irradiation, m<sup>6</sup>A level of *VANG1* is upregulated, as well as the *VANG1* mRNA stability regulated by IGF2BP2/3, suggesting a regulatory role of IGF2BPs in LUAD radio resistance [104].

Resistance to hypoxia-induced apoptosis, leading to distinctive hypoxia adaption in de novo lung adenocarcinomas (ADCs), has been an inevitable barrier leading to poor clinical outcomes of NSCLC treatment. The search for more specific hallmarks and treatments in both ADC and hypoxia-adapted NSCLC has become necessary and urgent. In the two different progressions of NSCLC, YTHDF1 behaves in completely distinct manners and provides us with new insights into cancer biology [54]. In ADC, YTHDF1 deficiency impedes cancer progression by inhibiting NSCLC cell proliferation and xenograft tumor formation through regulating the translation activity of cyclin proteins. However, lower expression of YTHDF1 is correlated with a worse clinical outcome

because of the resistance to cisplatin treatment in cancerous cells. Depletion of YTHDF1 in hypoxic solid tumors makes cancerous cells more resistant to cisplatin-dependent chemotherapy by regulating the *Keap1-Nrf2-AKR1C1* axis. In contrast, YTHDC2 is frequently suppressed in LUAD [99,100]. Mouse model studies have shown that YTHDC2 preferentially binds to *SLC7A11* mRNA and decreases tumorigenesis by suppressing cystine uptake and blocking the downstream antioxidant program [99].

## 9. YTH proteins in other cancers

The mRNAs recognized by YTH domain-containing proteins largely determine their roles in the progression of specific cancers. In oral squamous cell carcinoma, *c-Myc* is a direct YTHDF1 target gene and is responsible for the acceleration of tumor growth [65]. The negative p53 regulators *YY1* and *MDM2* are targets stimulated by YTHDF2 to promote human keratinocyte transformation in arsenic carcinogenesis [116]. In ocular melanoma, YTHDF1 inhibits the progression of both uveal and



**Fig. 3.** Deregulation of m<sup>6</sup>A reader proteins in human cancers.

A comprehensive overview of m<sup>6</sup>A readers in human carcinogenesis, including proliferation, migration, invasion, metastasis, and chemoresistance regulation. These readers specifically identify m<sup>6</sup>A-modified mRNAs and subsequently regulate critical signaling pathways in the pathogenesis of cancers. Red box line indicates upregulated expression and oncogenic functions. Blue line indicates downregulated expression and tumor suppressive functions. Black line indicates contradicted roles within different studies. Dashed gray line indicates unaffected or unknown roles.

conjunctival melanoma by promoting the translation of the tumor suppressor *HINT2* [58]. Notably, in addition to being recognized by YTHDF3 in CRC, the tumor suppressor *GAS* is also identified by YTHDF2 in cervical cancer and acts as a critical molecule for cervical cancer progression [72]. YTHDF2 also binds to *FOXO2*, *TNFRSF1B*, *PER1*, and *MOB3B* in gastric cancer, AML, pancreatic cancer and prostate cancer, respectively, providing similar insights into the underlying mechanisms of carcinogenesis induced by RNA degradation [74,83,87,88]. In glioblastoma (GBM), despite different target transcripts, YTHDF1/2 and YTHDC1 display similar function on GBM growth and progression, indicating positive prognostic and therapeutic prospects of these readers [69,70,75–77,95].

Studies have also indicated that more than one individual transcript participates in m<sup>6</sup>A reader-mediated effects in cancers. In ovarian cancer and Merkel cell carcinoma, YTHDF1 augments m<sup>6</sup>A-modified *EIF3C* translation and concomitantly increases the global translational output, thus facilitating tumorigenesis and cancer metastasis [60,67]. Additionally, YTHDF1 has been verified to bind to m<sup>6</sup>A-modified *PDK4* or *HK2* and promote cancer cell metabolic processes, such as glycolysis or the Warburg effect in cervical cancer, which suggests that YTHDF1 could be a promising therapeutic target [62,63]. Similarly, a group of transcripts encoding lysosomal proteases in dendritic cells are methylated and then recognized by YTHDF1. Deletion of YTHDF1 markedly inhibits these cathepsins, thus enhancing the antitumor response of antigen-specific CD8<sup>+</sup> T cells [117]. In bladder cancer, *SETD7* and *KLF4* were both identified as direct targets of YTHDF2 through transcriptome sequencing and confirmed with methylated RNA immunoprecipitation (MeRIP)-RT-qPCR [71].

In addition, other studies have indicated a positive correlation between YTHDF1 expression and cancer development, but there is a lack of mechanistic evidence in abdominal aortic aneurysm [118] and head and neck squamous cell carcinoma [119]. Likewise, low expression of YTHDC2 is significantly associated with worse overall survival in lung cancer patients. However, the potential RNAs that interact with YTHDC2 still need to be explored [100]. Further research is required to decipher the specific targets of m<sup>6</sup>A readers in these cancers, which may contribute to the understanding of their clinical significance and potential applications.

## 10. Conclusions and future perspectives

In this review, we summarized nearly all the recent findings on YTH domain-containing proteins in cancers. Dysregulation of m<sup>6</sup>A readers in cancer tissues results in abnormal mRNA metabolism of oncogenes and tumor suppressors, leading to the acceleration or delay of cancer development (Fig. 3). However, our understanding of how YTH proteins contribute to these processes remains incomplete. It is essential to identify the intrinsic and extrinsic signals that regulate the recruitment of YTH proteins to m<sup>6</sup>A-modified mRNAs and to identify these transcripts in a specific biological environment. In addition, the mechanisms that control the expression and activities of YTH proteins remain largely unknown. Therefore, future elucidation of the accurate roles of YTH proteins in regulating gene expression in cancers with new tools and methods is needed.

The crucial role of m<sup>6</sup>A readers in cancer initiation and progression provides potential targets for the treatment of cancers. Indeed, inhibition of YTHDF1 showed therapeutic potential alone or in combination with anti-PD-L1 immunotherapy in CRC [39,117]. However, small-molecule agents targeting YTH proteins, by blocking their interaction with m<sup>6</sup>A-modified mRNAs or other regulatory proteins, remain to be identified or developed. These small molecules might serve as a potential way to inhibit dysregulated mRNA metabolism by YTH proteins in cancer. On the other hand, targeted m<sup>6</sup>A editors and readers based on CRISPR/dCas system have been recently developed [30–33]. Taking advantage of these tools, manipulating the oncogenic or tumor suppressive mRNAs of interest with specific editors or readers could be

achieved to modulate cancer cell characteristics. Of note, Li et al. have used dCas13b-ALKBH5 to demethylate m<sup>6</sup>A modifications on oncogene transcripts such as *EGFR* and *MYC*, resulting in altered cancer cell proliferation in vitro [31]. However, whether these tools could be applied to cancer treatment remains to be explored.

## Author contributions

All the authors summarized and discussed. B.H. and X.G. wrote the manuscript.

## Declaration of competing interest

The authors declare that they have no conflict of interest.

## Acknowledgements

This work was supported by grants from National Natural Science Foundation of China (82073110 and 81672847 to X.G.), Zhejiang Youth Talent Support Program (ZJWR0308085 to X.G.), Zhejiang Provincial Natural Science Foundation of China (LQ19C050004 to S.W.), and Guangdong Provincial Science and Technology Project (2017A020214008).

## References

- [1] R. Desrosiers, K. Friderici, F. Rottman, Identification of methylated nucleosides in messenger RNA from Novikoff hepatoma cells, *Proc. Natl. Acad. Sci. U.S.A.* 71 (1974) 3971–3975.
- [2] C.M. Wei, A. Gershowitz, B. Moss, Methylated nucleotides block 5' terminus of HeLa cell messenger RNA, *Cell* 4 (1975) 379–386.
- [3] J.M. Adams, S. Cory, Modified nucleosides and bizarre 5'-termini in mouse myeloma mRNA, *Nature* 255 (1975) 28–33.
- [4] K.D. Meyer, Y. Saletore, P. Zumbo, O. Elemento, C.E. Mason, S.R. Jaffrey, Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons, *Cell* 149 (2012) 1635–1646.
- [5] D. Dominissini, S. Moshitch-Moshkovitz, S. Schwartz, M. Salmon-Divon, L. Ungar, S. Osenberg, K. Cesarkas, J. Jacob-Hirsch, N. Amariglio, M. Kupiec, R. Sorek, G. Rechavi, Topology of the human and mouse m<sup>6</sup>A RNA methylomes revealed by m<sup>6</sup>A-seq, *Nature* 485 (2012) 201–206.
- [6] B. Linder, A.V. Grozhik, A.O. Olarerin-George, C. Meydan, C.E. Mason, S. R. Jaffrey, Single-nucleotide-resolution mapping of m<sup>6</sup>A and m<sup>6</sup>Am throughout the transcriptome, *Nat. Methods* 12 (2015) 767–772.
- [7] G. Jia, Y. Fu, X. Zhao, Q. Dai, G. Zheng, Y. Yang, C. Yi, T. Lindahl, T. Pan, Y. G. Yang, C. He, N<sup>6</sup>-methyladenosine in nuclear RNA is a major substrate of the obesity-associated FTO, *Nat. Chem. Biol.* 7 (2011) 885–887.
- [8] Y. Fu, D. Dominissini, G. Rechavi, C. He, Gene expression regulation mediated through reversible m<sup>6</sup>A RNA methylation, *Nat. Rev. Genet.* 15 (2014) 293–306.
- [9] G. Zheng, J.A. Dahl, Y. Niu, P. Fedorcsak, C.M. Huang, C.J. Li, C.B. Vagbo, Y. Shi, W.L. Wang, S.H. Song, Z. Lu, R.P. Bosmans, Q. Dai, Y.J. Hao, X. Yang, W.M. Zhao, W.M. Tong, X.J. Wang, F. Bogdan, K. Furu, Y. Fu, G. Jia, X. Zhao, J. Liu, H. E. Krokan, A. Klungland, Y.G. Yang, C. He, ALKBH5 is a mammalian RNA demethylase that impacts RNA metabolism and mouse fertility, *Mol. Cell* 49 (2013) 18–29.
- [10] T. Lence, C. Paolantoni, L. Wörpenberg, J.Y. Roignant, Mechanistic insights into m<sup>6</sup>A RNA enzymes, *Biochim. Biophys. Acta Gene Regul. Mech.* 1862 (2019) 222–229.
- [11] I.A. Roundtree, M.E. Evans, T. Pan, C. He, Dynamic RNA modifications in gene expression regulation, *Cell* 169 (2017) 1187–1200.
- [12] B.S. Zhao, I.A. Roundtree, C. He, Post-transcriptional gene regulation by mRNA modifications, *Nat. Rev. Mol. Cell Biol.* 18 (2017) 31–42.
- [13] C. Roost, S.R. Lynch, P.J. Batista, K. Qu, H.Y. Chang, E.T. Kool, Structure and thermodynamics of N<sup>6</sup>-methyladenosine in RNA: a spring-loaded base modification, *J. Am. Chem. Soc.* 137 (2015) 2107–2115.
- [14] H. Huang, H. Weng, W. Sun, X. Qin, H. Shi, H. Wu, B.S. Zhao, A. Mesquita, C. Liu, C.L. Yuan, Y.C. Hu, S. Hüttelmaier, J.R. Skibbe, R. Su, X. Deng, L. Dong, M. Sun, C. Li, S. Nachtergaele, Y. Wang, C. Hu, K. Ferchen, K.D. Greis, X. Jiang, M. Wei, L. Qu, J.L. Guan, C. He, J. Yang, J. Chen, Recognition of RNA N<sup>6</sup>-methyladenosine by IGF2BP proteins enhances mRNA stability and translation, *Nat. Cell Biol.* 20 (2018) 285–295.
- [15] R.R. Edupuganti, S. Geiger, R.G.H. Lindeboom, H. Shi, P.J. Hsu, Z. Lu, S.Y. Wang, M.P.A. Baltissen, P. Jansen, M. Rossa, M. Muller, H.G. Stunnenberg, C. He, T. Carell, M. Vermeulen, N<sup>6</sup>-methyladenosine (m<sup>6</sup>A) recruits and repels proteins to regulate mRNA homeostasis, *Nat. Struct. Mol. Biol.* 24 (2017) 870–878.
- [16] N. Liu, K.I. Zhou, M. Parisien, Q. Dai, L. Diatchenko, T. Pan, N<sup>6</sup>-methyladenosine alters RNA structure to regulate binding of a low-complexity protein, *Nucleic Acids Res.* 45 (2017) 6051–6063.



- [17] C.R. Alarcon, H. Goodarzi, H. Lee, X. Liu, S. Tavazoie, S.F. Tavazoie, HNRNPA2B1 is a mediator of m(6)a-dependent nuclear RNA processing events, *Cell* 162 (2015) 1299–1308.
- [18] D. Theler, C. Dominguez, M. Blatter, J. Boudet, F.H. Allain, Solution structure of the YTH domain in complex with N6-methyladenosine RNA: a reader of methylated RNA, *Nucleic Acids Res.* 42 (2014) 13911–13919.
- [19] X. Wang, B.S. Zhao, I.A. Roundtree, Z. Lu, D. Han, H. Ma, X. Weng, K. Chen, H. Shi, C. He, N(6)-methyladenosine modulates messenger RNA translation efficiency, *Cell* 161 (2015) 1388–1399.
- [20] X. Wang, Z. Lu, A. Gomez, G.C. Hon, Y. Yue, D. Han, Y. Fu, M. Parisien, Q. Dai, G. Jia, B. Ren, T. Pan, C. He, N6-methyladenosine-dependent regulation of messenger RNA stability, *Nature* 505 (2014) 117–120.
- [21] H. Shi, X. Wang, Z. Lu, B.S. Zhao, H. Ma, P.J. Hsu, C. Liu, C. He, YTHDF3 facilitates translation and decay of N(6)-methyladenosine-modified RNA, *Cell Res.* 27 (2017) 315–328.
- [22] A. Li, Y.S. Chen, X.L. Ping, X. Yang, W. Xiao, Y. Yang, H.Y. Sun, Q. Zhu, P. Baidya, X. Wang, D.P. Bhattarai, Y.L. Zhao, B.F. Sun, Y.G. Yang, Cytoplasmic m(6)A reader YTHDF3 promotes mRNA translation, *Cell Res.* 27 (2017) 444–447.
- [23] P.J. Hsu, Y. Zhu, H. Ma, Y. Guo, X. Shi, Y. Liu, M. Qi, Z. Lu, H. Shi, J. Wang, Y. Cheng, G. Luo, Q. Dai, M. Liu, X. Guo, J. Sha, B. Shen, C. He, Ythdc2 is an N(6)-methyladenosine binding protein that regulates mammalian spermatogenesis, *Cell Res.* 27 (2017) 1115–1127.
- [24] J. Kretschmer, H. Rao, P. Hackert, K.E. Sloan, C. Höbartner, M.T. Bohnsack, The m6A reader protein YTHDC2 interacts with the small ribosomal subunit and the 5'-3' exoribonuclease XRN1, *RNA* 24 (2018) 1339–1350.
- [25] C. Xu, X. Wang, K. Liu, I.A. Roundtree, W. Tempel, Y. Li, Z. Lu, C. He, J. Min, Structural basis for selective binding of m6A RNA by the YTHDC1 YTH domain, *Nat. Chem. Biol.* 10 (2014) 927–929.
- [26] W. Xiao, S. Adhikari, U. Dahal, Y.S. Chen, Y.J. Hao, B.F. Sun, H.Y. Sun, A. Li, X. L. Ping, W.Y. Lai, X. Wang, H.L. Ma, C.M. Huang, Y. Yang, N. Huang, G.B. Jiang, H.L. Wang, Q. Zhou, X.J. Wang, Y.L. Zhao, Y.G. Yang, Nuclear m(6)A reader YTHDC1 regulates mRNA splicing, *Mol. Cell.* 61 (2016) 507–519.
- [27] I.A. Roundtree, G.Z. Luo, Z. Zhang, X. Wang, T. Zhou, Y. Cui, J. Sha, X. Huang, L. Guerrero, P. Xie, E. He, B. Shen, C. He, YTHDC1 mediates nuclear export of N(6)-methyladenosine methylated mRNAs, *Elife* 6 (2017) e31311.
- [28] N. Liu, Q. Dai, G. Zheng, C. He, M. Parisien, T. Pan, N(6)-methyladenosine-dependent RNA structural switches regulate RNA-protein interactions, *Nature* 518 (2015) 560–564.
- [29] K.I. Zhou, H. Shi, R. Lyu, A.C. Wylder, Z. Matuszek, J.N. Pan, C. He, M. Parisien, T. Pan, Regulation of Co-transcriptional pre-mRNA splicing by m(6)A through the low-complexity protein hnRNP, *Mol. Cell.* 76 (2019) 70–81.
- [30] X.M. Liu, J. Zhou, Y. Mao, Q. Ji, S.B. Qian, Programmable RNA N(6)-methyladenosine editing by CRISPR-Cas9 conjugates, *Nat. Chem. Biol.* 15 (2019) 865–871.
- [31] J. Li, Z. Chen, F. Chen, G. Xie, Y. Ling, Y. Peng, Y. Lin, N. Luo, C.M. Chiang, H. Wang, Targeted mRNA demethylation using an engineered dCas13b-ALKBH5 fusion protein, *Nucleic Acids Res.* 48 (2020) 5684–5694.
- [32] C. Wilson, P.J. Chen, Z. Miao, D.R. Liu, Programmable m(6)A modification of cellular RNAs with a Cas13-directed methyltransferase, *Nat. Biotechnol.* 38 (2020) 1431–1440.
- [33] S. Rauch, C. He, B.C. Dickinson, Targeted m(6)A reader proteins to study epitranscriptomic regulation of single RNAs, *J. Am. Chem. Soc.* 140 (2018) 11974–11981.
- [34] X. Deng, R. Su, H. Weng, H. Huang, Z. Li, J. Chen, RNA N(6)-methyladenosine modification in cancers: current status and perspectives, *Cell Res.* 28 (2018) 507–517.
- [35] H. Huang, H. Weng, J. Chen, m(6)A modification in coding and non-coding RNAs: roles and therapeutic implications in cancer, *Canc. Cell* 37 (2020) 270–288.
- [36] W. Zhao, X. Qi, L. Liu, S. Ma, J. Liu, J. Wu, Epigenetic regulation of m(6)A modifications in human cancer, *Mol. Ther. Nucleic Acids* 19 (2020) 405–412.
- [37] L. Liu, X. Liu, Z. Dong, J. Li, Y. Yu, X. Chen, F. Ren, G. Cui, R. Sun, N6-methyladenosine-related genomic targets are altered in breast cancer tissue and associated with poor survival, *J. Canc.* 10 (2019) 5447–5459.
- [38] R. Anita, A. Paramasivam, J.V. Priyadarsini, S. Chitra, The m6A readers YTHDF1 and YTHDF3 aberrations associated with metastasis and predict poor prognosis in breast cancer patients, *Am. J. Canc. Res.* 10 (2020) 2546–2554.
- [39] B. Han, S. Yan, S. Wei, J. Xiang, K. Liu, Z. Chen, R. Bai, J. Sheng, Z. Xu, X. Gao, YTHDF1-mediated translation amplifies Wnt-driven intestinal stemness, *EMBO Rep.* 21 (2020) e49229.
- [40] Y. Bai, C. Yang, R. Wu, L. Huang, S. Song, W. Li, P. Yan, C. Lin, D. Li, Y. Zhang, YTHDF1 regulates tumorigenicity and cancer stem cell-like activity in human colorectal carcinoma, *Front. Oncol.* 9 (2019) 332.
- [41] Y. Nishizawa, M. Konno, A. Asai, J. Koseki, K. Kawamoto, N. Miyoshi, H. Takahashi, N. Nishida, N. Haraguchi, D. Sakai, T. Kudo, T. Hata, C. Matsuda, T. Mizushima, T. Satoh, Y. Doki, M. Mori, H. Ishii, Oncogene c-Myc promotes epitranscriptome m6A reader YTHDF1 expression in colorectal cancer, *Oncotarget* 9 (2017) 7476–7486.
- [42] T. Liu, C. Li, L. Jin, C. Li, L. Wang, The prognostic value of m6A RNA methylation regulators in colon adenocarcinoma, *Med. Sci. Mon. Int. Med. J. Exp. Clin. Res.* 25 (2019) 9435–9445.
- [43] J. Tian, P. Ying, J. Ke, Y. Zhu, Y. Yang, Y. Gong, D. Zou, X. Peng, N. Yang, X. Wang, S. Mei, Y. Zhang, C. Wang, R. Zhong, J. Chang, X. Miao, ANKLE1 N(6)-Methyladenosine-related variant is associated with colorectal cancer risk by maintaining the genomic stability, *Int. J. Canc.* 146 (2020) 3281–3293.
- [44] C. Zhang, M. Zhang, S. Ge, W. Huang, X. Lin, J. Gao, J. Gong, L. Shen, Reduced m6A modification predicts malignant phenotypes and augmented Wnt/PI3K-Akt signaling in gastric cancer, *Canc. Med.* 8 (2019) 4766–4781.
- [45] T. Liu, S. Yang, Y.P. Cheng, X.L. Kong, D.D. Du, X. Wang, Y.F. Bai, L.H. Yin, Y. P. Pu, G.Y. Liang, The N6-methyladenosine (m6A) methylation gene YTHDF1 reveals a potential diagnostic role for gastric cancer, *Canc. Manag. Res.* 12 (2020) 11953–11964.
- [46] X. Zhao, Y. Chen, Q. Mao, X. Jiang, W. Jiang, J. Chen, W. Xu, L. Zhong, X. Sun, Overexpression of YTHDF1 is associated with poor prognosis in patients with hepatocellular carcinoma, *Canc. Biomarkers* 21 (2018) 859–868.
- [47] Y. Zhou, Z. Yin, B. Hou, M. Yu, R. Chen, H. Jin, Z. Jian, Expression profiles and prognostic significance of RNA N6-methyladenosine-related genes in patients with hepatocellular carcinoma: evidence from independent datasets, *Canc. Manag. Res.* 11 (2019) 3921–3931.
- [48] X. Lin, G. Chai, Y. Wu, J. Li, F. Chen, J. Liu, G. Luo, J. Tauler, J. Du, S. Lin, C. He, H. Wang, RNA m(6)A methylation regulates the epithelial mesenchymal transition of cancer cells and translation of Snail, *Nat. Commun.* 10 (2019) 2065.
- [49] X. Liu, J. Qin, T. Gao, C. Li, B. He, B. Pan, X. Xu, X. Chen, K. Zeng, M. Xu, C. Zhu, Y. Pan, H. Sun, L. Sun, T. Xu, S. Wang, YTHDF1 facilitates the progression of hepatocellular carcinoma by promoting FZD5 mRNA translation in an m6A-dependent manner, *Mol. Ther. Nucleic Acids* 22 (2020) 750–765.
- [50] Z. Lin, Y. Niu, A. Wan, D. Chen, H. Liang, X. Chen, L. Sun, S. Zhan, L. Chen, C. Cheng, X. Zhang, X. Bu, W. He, G. Wan, RNA m(6)A methylation regulates sorafenib resistance in liver cancer through FOXO3-mediated autophagy, *EMBO J.* 39 (12) (2020) e103181.
- [51] D. Jin, J. Guo, Y. Wu, J. Du, L. Yang, X. Wang, W. Di, B. Hu, J. An, L. Kong, L. Pan, G. Su, m(6)A mRNA methylation initiated by METTL3 directly promotes YAP translation and increases YAP activity by regulating the MALAT1-miR-1914-3p-YAP axis to induce NSCLC drug resistance and metastasis, *J. Hematol. Oncol.* 12 (2019) 135.
- [52] D. Jin, J. Guo, Y. Wu, L. Yang, X. Wang, J. Du, J. Dai, W. Chen, K. Gong, S. Miao, X. Li, H. Sun, m(6)A demethylase ALKBH5 inhibits tumor growth and metastasis by reducing YTHDFs-mediated YAP expression and inhibiting miR-107/LATS2-mediated YAP activity in NSCLC, *Mol. Cancer* 19 (2020) 40.
- [53] X. Li, N. Li, L. Huang, S. Xu, X. Zheng, A. Hamsath, M. Zhang, L. Dai, H. Zhang, J. J. Wong, M. Xian, C.T. Yang, J. Liu, Is hydrogen sulfide a concern during treatment of lung adenocarcinoma with ammonium tetrathiomolybdate? *Front. Oncol.* 10 (2020) 234.
- [54] Y. Shi, S. Fan, M. Wu, Z. Zuo, X. Li, L. Jiang, Q. Shen, P. Xu, L. Zeng, Y. Zhou, Y. Huang, Z. Yang, J. Zhou, J. Gao, H. Zhou, S. Xu, H. Ji, P. Shi, D.D. Wu, C. Yang, Y. Chen, YTHDF1 links hypoxia adaptation and non-small cell lung cancer progression, *Nat. Commun.* 10 (2019) 4892.
- [55] Y. Zhang, X. Liu, L. Liu, J. Li, Q. Hu, R. Sun, Expression, Prognostic Significance, Of m6A-related genes in lung adenocarcinoma, *Med. Sci. Mon. Int. Med. J. Exp. Clin. Res.* 26 (2020) e919644.
- [56] J. Zhu, M. Wang, D. Hu, Deciphering N(6)-methyladenosine-related genes signature to predict survival in lung adenocarcinoma, *BioMed Res. Int.* 2020 (2020) 2514230.
- [57] J. Zhou, D. Xiao, T. Qiu, J. Li, Z. Liu, Loading MicroRNA-376c in extracellular vesicles inhibits properties of non-small cell lung cancer cells by targeting YTHDF1, *Technol. Canc. Res. Treat.* 19 (2020), 1533033820977525.
- [58] R. Jia, P. Chai, S. Wang, B. Sun, Y. Xu, Y. Yang, S. Ge, R. Jia, Y.G. Yang, X. Fan, m(6)A modification suppresses ocular melanoma through modulating HINT2 mRNA translation, *Mol. Canc.* 18 (2019) 161.
- [59] T. Li, M. Gu, A. Deng, C. Qian, Increased expression of YTHDF1 and HNRNPA2B1 as potent biomarkers for melanoma: a systematic analysis, *Canc. Cell Int.* 20 (2020) 239.
- [60] T. Liu, Q. Wei, J. Jin, Q. Luo, Y. Liu, Y. Yang, C. Cheng, L. Li, J. Pi, Y. Si, H. Xiao, L. Li, S. Rao, F. Wang, J. Yu, J. Yu, D. Zou, P. Yi, The m6A reader YTHDF1 promotes ovarian cancer progression via augmenting EIF3C translation, *Nucleic Acids Res.* 48 (2020) 3816–3831.
- [61] L. Hao, J.M. Wang, B.Q. Liu, J. Yan, C. Li, J.Y. Jiang, F.Y. Zhao, H.Y. Qiao, H. Q. Wang, m6A-YTHDF1-mediated TRIM29 upregulation facilitates the stem cell-like phenotype of cisplatin-resistant ovarian cancer cells, *Biochim. Biophys. Acta Mol. Cell Res.* 1868 (2021) 118878.
- [62] Z. Li, Y. Peng, J. Li, Z. Chen, F. Chen, J. Tu, S. Lin, H. Wang, N6-methyladenosine regulates glycolysis of cancer cells through PDK4, *Nat. Commun.* 11 (2020) 2578.
- [63] Q. Wang, X. Guo, L. Li, Z. Gao, X. Su, M. Ji, J. Liu, N6-methyladenosine METTL3 promotes cervical cancer tumorigenesis and Warburg effect through YTHDF1/HK2 modification, *Cell Death Dis.* 11 (2020) 911.
- [64] L.C. Xinyuan Zhao, Development and validation of a m6A RNA methylation regulators-based signature for predicting the prognosis of head and neck squamous cell carcinoma, *Am. J. Canc. Res.* 9 (2019) 14.
- [65] W. Zhao, Y. Cui, L. Liu, X. Ma, X. Qi, Y. Wang, Z. Liu, S. Ma, J. Liu, J. Wu, METTL3 facilitates oral squamous cell carcinoma tumorigenesis by enhancing c-myc stability via YTHDF1-mediated m(6)A modification, *Mol. Ther. Nucleic Acids* 20 (2020) 1–12.
- [66] J. Ye, Z. Wang, X. Chen, X. Jiang, Z. Dong, S. Hu, W. Li, Y. Liu, B. Liao, W. Han, J. Shen, M. Xiao, YTHDF1-enhanced iron metabolism depends on TFRC m<sup>6</sup>A methylation, *Theranostics* 10 (2020) 12072–12089.
- [67] E. Orouji, W.K. Peitsch, A. Orouji, R. Houben, J. Utikal, Oncogenic role of an epigenetic reader of m6A RNA modification: YTHDF1 in Merkel cell carcinoma, *Cancers* 12 (2020) 202.
- [68] Y. He, J. Xing, S. Wang, S. Xin, Y. Han, J. Zhang, Increased m6A methylation level is associated with the progression of human abdominal aortic aneurysm, *Ann. Transl. Med.* 7 (2019) 797.



- [69] C. Xu, B. Yuan, T. He, B. Ding, S. Li, Prognostic values of YTHDF1 regulated negatively by mir-3436 in Glioma, *J. Cell Mol. Med.* 24 (2020) 7538–7549.
- [70] A.A. Yarmishyn, Y.P. Yang, K.H. Lu, Y.C. Chen, Y. Chien, S.J. Chou, P.H. Tsai, H. I. Ma, C.S. Chien, M.T. Chen, M.L. Wang, Musashi-1 promotes cancer stem cell properties of glioblastoma cells via upregulation of YTHDF1, *Canc. Cell Int.* 20 (2020) 597.
- [71] H. Xie, J. Li, Y. Ying, H. Yan, K. Jin, X. Ma, L. He, X. Xu, B. Liu, X. Wang, X. Zheng, L. Xie, METTL3/YTHDF2 m(6) A axis promotes tumorigenesis by degrading SETD7 and KLF4 mRNAs in bladder cancer, *J. Cell Mol. Med.* 24 (2020) 4092–4104.
- [72] X. Wang, J. Zhang, Y. Wang, Long noncoding RNA GAS5-AS1 suppresses growth and metastasis of cervical cancer by increasing GAS5 stability, *Am. J. Transl. Res.* 11 (2019) 4909–4921.
- [73] X. Chen, M. Xu, X. Xu, K. Zeng, X. Liu, B. Pan, C. Li, L. Sun, J. Qin, T. Xu, B. He, Y. Pan, H. Sun, S. Wang, METTL14-mediated N6-methyladenosine modification of SOX4 mRNA inhibits tumor metastasis in colorectal cancer, *Mol. Canc.* 19 (2020) 106.
- [74] X. Shen, K. Zhao, L. Xu, G. Cheng, J. Zhu, L. Gan, Y. Wu, Z. Zhuang, YTHDF2 inhibits gastric cancer cell growth by regulating FOXC2 signaling pathway, *Front. Genet.* 11 (2020) 592042.
- [75] D. Dixit, B. Prager, R. Gimple, H. Poh, Y. Wang, Q. Wu, Z. Qiu, R. Kidwell, L. Kim, Q. Xie, K. Vitting-Seerup, S. Bhargava, Z. Dong, L. Jiang, Z. Zhu, P. Hamerlik, S. Jaffrey, J. Zhao, X. Wang, J. Rich, The RNA m6A reader YTHDF2 maintains oncogene expression and is a targetable dependency in glioblastoma stem cells, *Canc. Discov.* 11 (2020) 480–499.
- [76] X. Lin, Z. Wang, G. Yang, G. Wen, H. Zhang, YTHDF2 correlates with tumor immune infiltrates in lower-grade glioma, *Aging* 12 (2020) 18476–18500.
- [77] R. Fang, X. Chen, S. Zhang, H. Shi, Y. Ye, H. Shi, Z. Zou, P. Li, Q. Guo, L. Ma, C. He, S. Huang, EGFR/SRC/ERK-stabilized YTHDF2 promotes cholesterol dysregulation and invasive growth of glioblastoma, *Nat. Commun.* 12 (2021) 177.
- [78] J. Hou, H. Zhang, J. Liu, Z. Zhao, J. Wang, Z. Lu, B. Hu, J. Zhou, Z. Zhao, M. Feng, H. Zhang, B. Shen, X. Huang, B. Sun, M.J. Smyth, C. He, Q. Xia, YTHDF2 reduction fuels inflammation and vascular abnormalization in hepatocellular carcinoma, *Mol. Canc.* 18 (2019) 163.
- [79] L. Zhong, D. Liao, M. Zhang, C. Zeng, X. Li, R. Zhang, H. Ma, T. Kang, YTHDF2 suppresses cell proliferation and growth via destabilizing the EGFR mRNA in hepatocellular carcinoma, *Canc. Lett.* 442 (2019) 252–261.
- [80] M. Chen, L. Wei, C.T. Law, F.H. Tsang, J. Shen, C.L. Cheng, L.H. Tsang, D.W. Ho, D.K. Chiu, J.M. Lee, C.C. Wong, I.O. Ng, C.M. Wong, RNA N6-methyladenosine methyltransferase-like 3 promotes liver cancer progression through YTHDF2-dependent posttranscriptional silencing of SOCS2, *Hepatology* 67 (2018) 2254–2270.
- [81] X.Y. Shao, J. Dong, H. Zhang, Y.S. Wu, L. Zheng, Systematic analyses of the role of the reader protein of N (6)-methyladenosine RNA methylation, YTH domain family 2, in liver hepatocellular carcinoma, *Front. Mol. Biosci.* 7 (2020) 577460.
- [82] C. Zhang, S. Huang, H. Zhuang, S. Ruan, Z. Zhou, K. Huang, F. Ji, Z. Ma, B. Hou, X. He, YTHDF2 promotes the liver cancer stem cell phenotype and cancer metastasis by regulating OCT4 expression via m6A RNA methylation, *Oncogene* 39 (2020) 4507–4518.
- [83] J. Paris, M. Morgan, J. Campos, G.J. Spencer, A. Shmakova, I. Ivanova, C. Mapperley, H. Lawson, D.A. Wotherspoon, C. Sepulveda, M. Vukovic, L. Allen, A. Sarapu, A. Tavasani, A.V. Guitart, A. Villacres, C. Much, J. Choe, A. Azar, L.N. van de Lagemaat, D. Vernimmen, A. Nehme, F. Mazurier, T.C.P. Somerville, R.I. Gregory, D. O'Carroll, K.R. Kranc, Targeting the RNA m(6)A reader YTHDF2 selectively compromises cancer stem cells in acute myeloid leukemia, *Cell Stem Cell* 25 (2019) 137–148.
- [84] Y. Qing, L. Dong, L. Gao, C. Li, Y. Li, L. Han, E. Prince, B. Tan, X. Deng, C. Wetzel, C. Shen, M. Gao, Z. Chen, W. Li, B. Zhang, D. Braas, J. Ten Hoeve, G.J. Sanchez, H. Chen, L.N. Chan, C.W. Chen, D. Ann, L. Jiang, M. Muschen, G. Marcucci, D. R. Plas, Z. Li, R. Su, J. Chen, R-2-hydroxyglutarate attenuates aerobic glycolysis in leukemia by targeting the FTO/m(6)A/PFKP/LDHB axis, *Mol. Cell.* 81 (2021) 922–939.
- [85] H. Sheng, Z. Li, S. Su, W. Sun, X. Zhang, L. Li, J. Li, S. Liu, B. Lu, S. Zhang, C. Shan, YTH domain family 2 promotes lung cancer cell growth by facilitating 6-phosphogluconate dehydrogenase mRNA translation, *Carcinogenesis* 41 (2020) 541–550.
- [86] J. Li, L. Wu, M. Pei, Y. Zhang, YTHDF2, a protein repressed by miR-145, regulates proliferation, apoptosis, and migration in ovarian cancer cells, *J. Ovarian Res.* 13 (2020) 111.
- [87] X. Guo, K. Li, W. Jiang, Y. Hu, W. Xiao, Y. Huang, Y. Feng, Q. Pan, R. Wan, RNA demethylase ALKBH5 prevents pancreatic cancer progression by posttranscriptional activation of PER1 in an m6A-YTHDF2-dependent manner, *Mol. Canc.* 19 (2020) 91.
- [88] C. Du, C. Lv, Y. Feng, S. Yu, Activation of the KDM5A/miRNA-495/YTHDF2/m6A-MOB3B axis facilitates prostate cancer progression, *J. Exp. Clin. Canc. Res.* 39 (2020) 223.
- [89] J. Li, S. Meng, M. Xu, S. Wang, L. He, X. Xu, X. Wang, L. Xie, Downregulation of N 6-methyladenosine binding YTHDF2 protein mediated by miR-493-3p suppresses prostate cancer by elevating N 6-methyladenosine levels, *Oncotarget* 9 (2017) 3752–3764.
- [90] J. Li, H. Xie, Y. Ying, H. Chen, H. Yan, L. He, M. Xu, X. Xu, Z. Liang, B. Liu, X. Wang, X. Zheng, L. Xie, YTHDF2 mediates the mRNA degradation of the tumor suppressors to induce AKT phosphorylation in N6-methyladenosine-dependent way in prostate cancer, *Mol. Canc.* 19 (2020) 152.
- [91] G. Chang, L. Shi, Y. Ye, H. Shi, L. Zeng, S. Tiwary, J. Huse, L. Huo, L. Ma, Y. Ma, S. Zhang, J. Zhu, V. Xie, P. Li, L. Han, C. He, S. Huang, YTHDF3 induces the translation of mA-enriched gene transcripts to promote breast cancer brain metastasis, *Canc. Cell* 38 (2020) 857–871.
- [92] W. Ni, S. Yao, Y. Zhou, Y. Liu, P. Huang, A. Zhou, J. Liu, L. Che, J. Li, Long noncoding RNA GAS5 inhibits progression of colorectal cancer by interacting with and triggering YAP phosphorylation and degradation and is negatively regulated by the m(6)A reader YTHDF3, *Mol. Canc.* 18 (2019) 143.
- [93] M. Wang, Y. Yang, J. Yang, J. Yang, S. Han, circ KIAA1429 accelerates hepatocellular carcinoma advancement through the mechanism of m(6)A-YTHDF3-Zeb1, *Life Sci.* 257 (2020) 118082.
- [94] X. Liu, L. Liu, Z. Dong, J. Li, Y. Yu, X. Chen, F. Ren, G. Cui, R. Sun, Expression patterns and prognostic value of m(6)A-related genes in colorectal cancer, *Am. J. Transl. Res.* 11 (2019) 3972–3991.
- [95] F. Li, Y. Yi, Y. Miao, W. Long, T. Long, S. Chen, W. Cheng, C. Zou, Y. Zheng, X. Wu, J. Ding, K. Zhu, D. Chen, Q. Xu, J. Wang, Q. Liu, F. Zhi, J. Ren, Q. Cao, W. Zhao, N(6)-Methyladenosine modulates nonsense-mediated mRNA decay in human glioblastoma, *Canc. Res.* 79 (2019) 5785–5798.
- [96] J. Liu, G. Sun, S. Pan, M. Qin, R. Ouyang, Z. Li, J. Huang, The Cancer Genome Atlas (TCGA) based m(6)A methylation-related genes predict prognosis in hepatocellular carcinoma, *Bioengineered* 11 (2020) 759–768.
- [97] H.J. Luxton, B.S. Simpson, I.G. Mills, N.R. Brindle, Z. Ahmed, V. Stavrindes, S. Heavey, S. Stamm, H.C. Whitaker, The oncogene metadherin interacts with the known splicing proteins YTHDC1, Sam68 and T-STAR and plays a novel role in alternative mRNA splicing, *Cancers* 11 (2019) 1233.
- [98] Y. Li, J.N. Zheng, E.H. Wang, C.J. Gong, K.F. Lan, X. Ding, The m6A reader protein YTHDC2 is a potential biomarker and associated with immune infiltration in head and neck squamous cell carcinoma, *PeerJ* 8 (2020) e10385.
- [99] L. Ma, T. Chen, X. Zhang, Y. Miao, X. Tian, K. Yu, X. Xu, Y. Niu, S. Guo, C. Zhang, S. Qiu, Y. Qiao, W. Wang, L. Du, Y. Yu, J. Wang, The m(6)A reader YTHDC2 inhibits lung adenocarcinoma tumorigenesis by suppressing SLC7A11-dependent antioxidant function, *Redox Biol* 38 (2021) 101801.
- [100] S. Sun, Q. Han, M. Liang, Q. Zhang, J. Zhang, J. Cao, Downregulation of m(6) A reader YTHDC2 promotes tumor progression and predicts poor prognosis in non-small cell lung cancer, *Thorac. Canc.* 11 (2020) 3269–3279.
- [101] G. Bao, J. Huang, W. Pan, X. Li, T. Zhou, Long noncoding RNA CERS6-AS1 functions as a malignancy promoter in breast cancer by binding to IGF2BP3 to enhance the stability of CERS6 mRNA, *Canc. Med.* 9 (2020) 278–289.
- [102] Q. Wang, C. Chen, Q. Ding, Y. Zhao, Z. Wang, J. Chen, Z. Jiang, Y. Zhang, G. Xu, J. Zhang, J. Zhou, B. Sun, X. Zou, S. Wang, METTL3-mediated m(6)A modification of HDGF mRNA promotes gastric cancer progression and has prognostic significance, *Gut* 69 (2020) 1193–1205.
- [103] W. Guo, Q. Huai, H. Wan, L. Guo, P. Song, S. Gao, J. He, Prognostic impact of IGF2BP3 expression in patients with surgically resected lung adenocarcinoma, *DNA Cell Biol.* 40 (2021) 316–331.
- [104] C.C. Hao, C.Y. Xu, X.Y. Zhao, J.N. Luo, G. Wang, L.H. Zhao, X. Ge, X.F. Ge, Up-regulation of VANG1 by IGF2BPs and miR-29b-3p attenuates the detrimental effect of irradiation on lung adenocarcinoma, *J. Exp. Clin. Canc. Res.* 39 (2020) 256.
- [105] T. Li, P.S. Hu, Z. Zuo, J.F. Lin, X. Li, Q.N. Wu, Z.H. Chen, Z.L. Zeng, F. Wang, J. Zheng, D. Chen, B. Li, T.B. Kang, D. Xie, D. Lin, H.Q. Ju, R.H. Xu, METTL3 facilitates tumor progression via an m(6)A-IGF2BP2-dependent mechanism in colorectal carcinoma, *Mol. Canc.* 18 (2019) 112.
- [106] J. Tian, P. Ying, J. Ke, Y. Zhu, Y. Yang, Y. Gong, D. Zou, X. Peng, N. Yang, X. Wang, S. Mei, Y. Zhang, C. Wang, R. Zhong, J. Chang, X. Miao, ANKLE1 N-Methyladenosine-related variant is associated with colorectal cancer risk by maintaining the genomic stability, *Int. J. Canc.* 146 (2020) 3281–3293.
- [107] A. Tanabe, K. Tanikawa, M. Tsunetomi, K. Takai, H. Ikeda, J. Konno, T. Torigoe, H. Maeda, G. Kutomi, K. Okita, M. Mori, H. Sahara, RNA helicase YTHDC2 promotes cancer metastasis via the enhancement of the efficiency by which HIF-1α mRNA is translated, *Canc. Lett.* 376 (2016) 34–42.
- [108] F. Bray, J. Ferlay, I. Soerjomataram, R.L. Siegel, L.A. Torre, A. Jemal, Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries, *CA A Cancer J. Clin.* 68 (2018) 394–424.
- [109] J.Z. Ma, F. Yang, C.C. Zhou, F. Liu, J.H. Yuan, F. Wang, T.T. Wang, Q.G. Xu, W. P. Zhou, S.H. Sun, METTL14 suppresses the metastatic potential of hepatocellular carcinoma by modulating N(6)-methyladenosine-dependent primary MicroRNA processing, *Hepatology* 65 (2017) 529–543.
- [110] Z. Lin, Y. Niu, A. Wan, D. Chen, H. Liang, X. Chen, L. Sun, S. Zhan, L. Chen, C. Cheng, X. Zhang, X. Bu, W. He, G. Wan, RNA m(6) A methylation regulates sorafenib resistance in liver cancer through FOXO3-mediated autophagy, *EMBO J.* 39 (2020) e103181.
- [111] R.S. Herbst, D. Morgensztern, C. Boshoff, The biology and management of non-small cell lung cancer, *Nature* 553 (2018) 446–454.
- [112] J. Choe, S. Lin, W. Zhang, Q. Liu, L. Wang, J. Ramirez-Moya, P. Du, W. Kim, S. Tang, P. Sliz, P. Santisteban, R.E. George, W.G. Richards, K.K. Wong, N. Locker, F.J. Slack, R.I. Gregory, mRNA circularization by METTL3-eIF3h enhances translation and promotes oncogenesis, *Nature* 561 (2018) 556–560.
- [113] S. Lin, J. Choe, P. Du, R. Triboulet, R.I. Gregory, The m(6)A methyltransferase METTL3 promotes translation in human cancer cells, *Mol. Cell.* 62 (2016) 335–345.
- [114] F. Lo Sardo, S. Strano, G. Blandino, YAP and TAZ in lung cancer: oncogenic role and clinical targeting, *Cancers* 10 (2018) 137.
- [115] Y. Zhang, X. Liu, L. Liu, J. Li, Q. Hu, R. Sun, Expression and Prognostic Significance of m6A-Related Genes in Lung Adenocarcinoma, *Med. Sci. Mon.* 26 (2020) e919644.

- [116] T. Zhao, D. Sun, M. Zhao, Y. Lai, Y. Liu, Z. Zhang, N(6)-methyladenosine mediates arsenite-induced human keratinocyte transformation by suppressing p53 activation, *Environ. Pollut.* 259 (2020) 113908.
- [117] D. Han, J. Liu, C. Chen, L. Dong, Y. Liu, R. Chang, X. Huang, Y. Liu, J. Wang, U. Dougherty, M.B. Bissonnette, B. Shen, R.R. Weichselbaum, M.M. Xu, C. He, Anti-tumour immunity controlled through mRNA m(6)A methylation and YTHDF1 in dendritic cells, *Nature* 566 (2019) 270–274.
- [118] Y. He, J. Xing, S. Wang, S. Xin, Y. Han, J. Zhang, Increased m6A methylation level is associated with the progression of human abdominal aortic aneurysm, *Ann. Transl. Med.* 7 (2019) 797.
- [119] X. Zhao, L. Cui, Development and validation of a mA RNA methylation regulators-based signature for predicting the prognosis of head and neck squamous cell carcinoma, *Am. J. Canc. Res.* 9 (2019) 2156–2169.