

In many applications “higher order” terms are neglected in order to find simpler or manageable formulas.

The base for this and for an estimate of the resulting error is the following theorem:

Theorem (Taylor theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be an $n + 1$ -times continuously differentiable function.

Then for any two points x and x_0 in (a, b) there exists a number $\Theta_{x, x_0} \in (0, 1)$ such that

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \underbrace{\frac{f^{(n+1)}(x_0 + \Theta_{x, x_0} \cdot (x - x_0))}{(n+1)!} (x - x_0)^{n+1}}_{\approx |x - x_0|^n}$$

Notation:

$$* (x - x_0)^0 = 1$$

$$* 0! = 1$$

$$* f^{(0)} = f$$

Definition (Taylor Polynomial, Lagrange Form of the Remainder)

We call

$$T_n(f, x, x_0) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$$


the **Taylor polynomial of n-th order** (not of n-th degree!) around the center x_0 .

The term

$$\frac{f^{(n+1)}(x_0 + \Theta_{x,x_0} \cdot (x - x_0))}{(n+1)!} (x - x_0)^{n+1}$$

is called the **Lagrange form of the remainder**.

There are other forms of the ~~R~~ Remainder: Peano form,
Cauchy form, ...
↑



Examples for the Taylor Theorem

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

$$f(x) = \sqrt{1+x} = (1+x)^{1/2}$$

Taylor polynomial of order 3 around $x_0 = 0$

$$f'(x) = \frac{1}{2} (1+x)^{-1/2}$$

$$f''(x) = -\frac{1}{4} (1+x)^{-3/2}$$

$$f'''(x) = \frac{3}{8} (1+x)^{-5/2}$$


$$f^{(4)}(x) = -\frac{15}{16} (1+x)^{-7/2}$$

$$\text{Thus } T_3(f, x, 0) = f(0) + \frac{f'(0)}{1!} (x-0) + \frac{f''(0)}{2!} (x-0)^2 + \frac{f^{(3)}(0)}{3!} (x-0)^3$$

$$= 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3$$

$$\text{and } R_3(f, x, 0) = \frac{-15}{16 \cdot \frac{24}{8}} \frac{x^4}{(1 + \underbrace{\partial_{x, x_0} x}_{\in (0,1)})^{7/2}} = \frac{-5x^4}{128 (1 + \partial_{x, x_0} x)^{7/2}}$$

$$\text{--- } 4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$$



Definition (Taylor Series)

Let $f : [a, b] \rightarrow \mathbb{R}$ be arbitrarily many times differentiable.

Then for $x \in [a, b]$ the power series

$$T(f, x, x_0) = \sum_{k=0}^{\infty} \frac{f^{(k)}(\overset{x_0}{\cancel{0}})}{k!} (x - x_0)^k.$$

is called the **Taylor series** of the function f with the center x_0 .

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series


Fourier Series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Theorem (Convergence Criterion for Taylor Series)


The Taylor series $T(f, x, x_0)$ converges for a given $x \in [a, b]$ to $f(x)$

iff

$$\lim_{n \rightarrow \infty} R_n(f, x, x_0) = 0.$$

We are interested in an interval $[a, b]$ such that the remainder converges to 0 and this should happen “fast”.

Warning: There exist functions whose Taylor series represent the function in a single point only!



An important special case: $x_0 = 0$.


Let f be as in the Taylor theorem and $0 \in (a, b)$. For any point $x \in (a, b)$ there exists a number $\Theta_x \in (0, 1)$ such that

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(0)}{k!} x^k + \frac{f^{(n+1)}(\Theta_x x)}{(n+1)!} x^{n+1}.$$

The corresponding series is called **Maclaurin series** of the function f :

$$M(f, x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k.$$

Its center is $x_0 = 0$.



A Taylor Series for the Logarithm

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series


Fourier Series

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



$$\bullet \ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}, \quad x \in \left[-\frac{1}{2}, 1\right], \quad x_0 = 0$$

By induction: $f^{(k)}(x) = (-1)^{k-1} \frac{(k-1)!}{(1+x)^k}, \quad k \in \mathbb{N}$

$M(f, x)$

$$T(f, x, 0) = \underbrace{\ln(1)}_0 + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(n-1)!}{(1+0)^n} \frac{1}{n} (x-0)^n$$

$$= \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

$$R_n(f, x, 0) = \frac{f^{(n+1)}(\theta_x x)}{(n+1)!} x^{n+1} = \frac{(-1)^n n!}{(n+1)!} \frac{x^{n+1}}{(1+\theta_x x)^{n+1}} = \frac{(-1)^n}{n+1} \left(\frac{x}{1+\theta_x x} \right)^{n+1}$$

for $0 \leq x \leq 1$: $\left| \frac{x}{1+\theta_x x} \right| \leq 1$ clearly also holds

for $-\frac{1}{2} \leq x < 0$: $|x| = -x \leq \frac{1}{2} \leq 1+x < 1+\theta_x x$

$$\bullet \ln(2) = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots$$

$$= \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n}$$

alternating harmonic series

\Rightarrow the series converges