

## Problem (General Interpolation Task)


*For a given function  $g : \mathbb{R} \rightarrow \mathbb{R}$  with given points*

$$(x_j, g(x_j)), \quad j = 1, \dots, n,$$

*we search for a function  $p$ , s.t. the graph of  $p$  goes through these given points.*

*$p$  is chosen within a class of reasonable (and manageable) functions, e.g.*

- *the class of real polynomials,*
- *the class of trigonometric polynomials in case of  $T$ -periodic functions.*



# Trigonometric Approximation Revisited

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform  
Convergence

Power Series

Taylor Series

Fourier Series


Discrete Fourier Transform

Differentiation in  
Higher Dimensions

Integration in  
Higher Dimensions

Further Topics in  
Calculus

Summary -  
Outlook and  
Review



In the later case we consider a function  $g : [0, 2\pi] \rightarrow \mathbb{R}$ , periodically extended with  $T = 2\pi$ .

Let

$$x_0 = 0, x_1 = \frac{2\pi}{n}, \dots, x_{n-1} = (n-1)\frac{2\pi}{n}, x_n = 2\pi.$$

We search for a trigonometric polynomial

$$p_n(x) := \sum_{k=0}^{n-1} c_k \exp(ikx)$$

such that

$$p_n(x_j) = g(x_j) \quad \text{for } j = 0, \dots, n-1.$$

Remark: Due to the periodicity, we have  $g(x_0) = g(x_n)$ .

By inserting:

$$\sum_{k=0}^{n-1} \exp(ikx_j) c_k = g(x_j) \quad \text{for } j = 0, \dots, n-1.$$

This is a linear equation system (LES) with coefficient matrix

$$f_{jk} = \exp\left(2\pi i k \frac{j}{n}\right) = \zeta^{jk} \quad \text{with } \zeta := \exp\left(2\pi i \frac{1}{n}\right).$$

This motivates the definition of the so-called **Fourier matrix** :

$$F_n = (f_{ij})_{i,j=1,\dots,n}$$

Properties of the Fourier matrix:


- $F_n$  is symmetric
- $\frac{1}{\sqrt{n}} F_n$  is unitary, i.e.  $\left(\frac{1}{\sqrt{n}} F_n\right)^{-1} = \overline{\left(\frac{1}{\sqrt{n}} F_n\right)}^\top$  (details later in LA)

Thus, we may rewrite the LES

$$F_n \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} = \begin{pmatrix} g(x_1) \\ \vdots \\ g(x_n) \end{pmatrix}$$

that uniquely determines the trigonometric polynomial  $p_n$  by

$$\begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} = \frac{1}{n} \overline{F_n}^\top \begin{pmatrix} g(x_1) \\ \vdots \\ g(x_n) \end{pmatrix}$$



The mapping

$$\mathbb{C}^n \rightarrow \mathbb{C}^n, v \mapsto F_n v$$


is called **discrete Fourier transform (DFT)**, i.e.

$$(F_n v)_j = \sum_{k=0}^{n-1} v_k \exp\left(2\pi i \frac{jk}{n}\right).$$

This may be interpreted as a discretization of the (continuous) Fourier transform (at the end of lecture).

The inverse DFT solves the trigonometric interpolation problem.

By an algorithm, called **fast Fourier transform (FFT)** the DFT may be computed with only "few" operations!



# Example: Fourier Matrix

Let  $v^T = (1, -1, 0, 1)$ .  $= \begin{pmatrix} 1 \\ -1 \\ 0 \\ 1 \end{pmatrix}$

$\zeta_{(4)} = e^{i\frac{2\pi}{4}} = i$

Compute

$$F_4 v = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & \zeta^1 & \zeta^2 & \zeta^3 \\ 1 & \zeta^2 & \zeta^4 & \zeta^6 \\ 1 & \zeta^3 & \zeta^6 & \zeta^9 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \\ 0 \\ 1 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & i & i^2 = -1 & -i \\ 1 & -1 & 1 & -1 \\ 1 & -i & -1 & i \end{pmatrix} \begin{pmatrix} 1 \\ -1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 - 1 + 0 + 1 \\ 1 - i + 0 - i \\ 1 + 1 + 0 - 1 \\ 1 + i + 0 + i \end{pmatrix} = \begin{pmatrix} 1 \\ 1 - 2i \\ 1 \\ 1 + 2i \end{pmatrix}$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series

Fourier Series


Discrete Fourier Transform

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review






An image with decreasing quality from the left to the right.  
(Source: [Wikipedia](#))

Typical standards: `jpeg`, `mpeg`

An important step (among others) in these data compression methods is the discrete Fourier cosine transform that maps into the frequency domain.

Theoretically, this step is invertible, i.e. without loss of information possible.



At the end of a lecture (if time permits) we will have a short look at

- (Continuous) Fourier transform
- Laplace transform
- Short-time Fourier transform
- Wavelet

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform  
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in  
Higher Dimensions

Integration in  
Higher Dimensions

Further Topics in  
Calculus

Summary -  
Outlook and  
Review

