

Introduction

Power series

Differentiation in
Higher Dimensions


Integration in
Higher Dimensions

**Further Topics in
Calculus**

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



- 1 Introduction
- 2 Power series
- 3 Differentiation in Higher Dimensions
- 4 Integration in Higher Dimensions
- 5 Further Topics in Calculus**
 - Vector Analysis
 - Integral Transforms
- 6 Summary - Outlook and Review

Let $D \subseteq \mathbb{R}^n$.

A function $\mathbf{F} : D \rightarrow \mathbb{R}^n$ with $n \geq 2$ is also called a **vector field**.

Example:


Let $f : D \rightarrow \mathbb{R}$, then ∇f is a vector field.

A vector field F is called **conservative** or **gradient field**, if there exists a function ϕ s.t.:

$$\mathbf{F} = \nabla \phi.$$

↑ with a " - " in physics

Then we call ϕ a (scalar) **potential (function)**.



Path independence:


Let D be a simply connected domain.

$$\int_{\mathbf{a}}^{\mathbf{b}} \mathbf{F} \cdot d\mathbf{s} = \phi(\mathbf{b}) - \phi(\mathbf{a})$$

for a sufficiently smooth curve from \mathbf{a} to \mathbf{b} .

Note that the integral does not depend on the curve, but only on the start and end points, \mathbf{a} and \mathbf{b} .

How can we decide whether a vector field is conservative?



Consider a simply connected domain D .

The following statements are equivalent:


- $\int_a^b \mathbf{F} \cdot d\mathbf{s} = \phi(\mathbf{b}) - \phi(\mathbf{a})$ for a curve from \mathbf{a} to \mathbf{b} .
- The curve integral over \mathbf{F} along any closed curve is zero.

- $\nabla \times \mathbf{F} = 0$ $\stackrel{n=3}{<=}$ $\frac{\partial}{\partial x_i} F_j = \frac{\partial}{\partial x_j} F_i, 1 \leq i, j \leq n$

integrability condition

We suppose that the involved curves are sufficiently smooth.

Note that ϕ is unique up to an additive constant.




Examples: Potential

Consider $\vec{v}: \mathbb{R}^2 \rightarrow \mathbb{R}^2, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ 1 \end{pmatrix}$

$\vec{s}_1: [0, 1] \rightarrow \mathbb{R}^2, t \mapsto \begin{pmatrix} t \\ \sqrt{1-t^2} \end{pmatrix}$


$\vec{s}_2: [0, \frac{\pi}{2}] \rightarrow \mathbb{R}^2, t \mapsto \begin{pmatrix} \cos t \\ \sin t \end{pmatrix}$



\vec{v} is conservative, i.e. there exists a potential ϕ

$$\frac{\partial}{\partial x} v_2 = \frac{\partial}{\partial x} 1 = 0 = \frac{\partial}{\partial y} v_1 = \frac{\partial}{\partial y} x \quad \checkmark$$

$$-\int_{\vec{s}_1} \vec{v} \cdot d\vec{s} \stackrel{\text{cons.}}{=} \int_{\vec{s}_2} \vec{v} \cdot d\vec{s} = \int_0^{\pi/2} \vec{v}(\vec{r}(t)) \cdot \vec{r}'(t) dt = \dots = \frac{1}{2}$$



The fundamental theorem of Analysis 1

$$F(x) := \int_a^x F'(\tilde{x}) d\tilde{x}$$

or

$$\int_a^b F'(x) dx = F(b) - F(a)$$

cannot be easily generalized to several dimensions.

Introduction

Power series

Differentiation in
Higher Dimensions


Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



Some generalizations are:

- Gradient theorem (“1d”)

$$\int_{\gamma} \nabla f(\mathbf{r}) \cdot d\mathbf{r} = f(\mathbf{r}(\mathbf{b})) - f(\mathbf{r}(\mathbf{a}))$$

(γ a curve from \mathbf{a} to \mathbf{b} in \mathbb{R}^n with parametrization \mathbf{r})

- Stokes theorem (2d)

$$\iint_M (\nabla \times \mathbf{F}) \cdot d\mathbf{A} = \int_{\partial M} \mathbf{F} \cdot d\boldsymbol{\gamma}$$


(M a 2d submanifold of \mathbb{R}^3 , bounded by a curve γ representing ∂M)

- Gauss divergence theorem in 3d

\leftarrow outer normal
 $= \vec{n} \cdot d\mathbf{a}$

$$\iiint_K \underbrace{\nabla \cdot \mathbf{F}}_{\text{sinks/sources}} dV = \iint_{\partial K} \mathbf{F} \cdot d\mathbf{A}$$

($K \subseteq \mathbb{R}^3$ compact. May be generalized to \mathbb{R}^n .)



Recall: gradient

$$\operatorname{grad} f(\mathbf{x}) = \nabla f(\mathbf{x}) = \left(\frac{\partial}{\partial x_1} f(\mathbf{x}), \dots, \frac{\partial}{\partial x_n} f(\mathbf{x}) \right)^\top$$

Divergence (sources or sinks)

$$\operatorname{div} \mathbf{F}(\mathbf{x}) = \nabla \cdot \mathbf{F}(\mathbf{x}) = \sum_{i=1}^n \frac{\partial}{\partial x_i} F_i(\mathbf{x})$$

Curl or rot(ation), in \mathbb{R}^3 (\mathbb{R}^2 also possible):

$$\operatorname{curl} \mathbf{F}(\mathbf{x}) = \nabla \times \mathbf{F}(\mathbf{x}) = \begin{pmatrix} \frac{\partial}{\partial x_2} F_3(\mathbf{x}) - \frac{\partial}{\partial x_3} F_2(\mathbf{x}) \\ \frac{\partial}{\partial x_3} F_1(\mathbf{x}) - \frac{\partial}{\partial x_1} F_3(\mathbf{x}) \\ \frac{\partial}{\partial x_1} F_2(\mathbf{x}) - \frac{\partial}{\partial x_2} F_1(\mathbf{x}) \end{pmatrix}$$

Laplace operator

$$\operatorname{div} \operatorname{grad} f(\mathbf{x}) = \nabla \cdot \nabla f(\mathbf{x}) = \Delta f(\mathbf{x}) = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} f(\mathbf{x})$$

Introduction

Power series

Differentiation in
Higher Dimensions


Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review




Integration by parts

$$\iiint_V g \nabla \cdot \mathbf{f} dV = \iint_{\partial V} g \mathbf{f} \cdot d\mathbf{A} - \iiint_V \nabla g \cdot \mathbf{f} dV$$

(May be generalized to \mathbb{R}^n .)

$$\nabla \cdot (g \vec{f}) = \nabla g \cdot \vec{f} + g (\nabla \cdot \vec{f})$$


- Introduction
- Power series
- Differentiation in Higher Dimensions
- Integration in Higher Dimensions
- Further Topics in Calculus
- Vector Analysis
- Integral Transforms
- Summary - Outlook and Review



Fourier and Laplace transforms are useful tools for solving (partial) differential equations

Applications of the Fourier transform

- Information technology/electrical engineering:
Low-pass filter
- Spectroscopy (NMR, FTIR, ...)
- Acoustics
- Quantum mechanics
- ...
- Generalizes Fourier series from periodic to (some) non-periodic functions



Suppose

$$f(x) = \sum_{k=-\infty}^{\infty} c_k \exp(ik(\Delta\omega)x), \quad \Delta\omega = \frac{2\pi}{T}$$

with complex Fourier coefficients

$$\begin{aligned} c_k &= \frac{1}{T} \int_0^T f(x) \exp(-ik\cancel{\omega}x) dx \\ &= \frac{1}{T} \int_{-T/2}^{T/2} f(x) \overline{\exp(i\cancel{\omega}x)} dx, \quad k \in \mathbb{Z} \end{aligned}$$


Handwritten notes: $\Delta\omega$ above the first ω , $\Delta\omega$ below the second ω .

Introduce $\omega_k := k\Delta\omega$ and insert c_k into the series:

$$f(x) = \sum_{k=-\infty}^{\infty} \left(\frac{1}{T} \int_{-T/2}^{T/2} f(x) \overline{\exp(i\cancel{\omega}x)} \right) \exp(i\omega_k x) = \dots$$

Handwritten notes: k and dx below the integral, with an arrow pointing from k to the ω_k term.


- Introduction
- Power series
- Differentiation in Higher Dimensions
- Integration in Higher Dimensions
- Further Topics in Calculus
- Vector Analysis
- Integral Transforms
- Summary - Outlook and Review



$$\begin{aligned}
 f(x) &= \sum_{k=-\infty}^{\infty} \left(\frac{1}{T} \int_{-T/2}^{T/2} f(\xi) \overline{\exp(i\omega_k \xi)} \right) \exp(i\omega_k x) \\
 &= \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \frac{2\pi}{T} \left(\int_{-T/2}^{T/2} f(\xi) \overline{\exp(i\omega_k \xi)} d\xi \right) \exp(i\omega_k x) \\
 &= \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \left(\int_{-T/2}^{T/2} f(\xi) \overline{\exp(i\omega_k \xi)} d\xi \right) \exp(i\omega_k x) \Delta\omega
 \end{aligned}$$

$\Delta\omega \rightarrow 0, \quad T \rightarrow \infty$

The series may be interpreted as a limit of Riemann sums that approximate an integral on $[-\infty, \infty]$ with refinement $\Delta\omega$ and grid points $\omega_k, k \in \mathbb{Z}$.



For $T \rightarrow \infty$ (and $\Delta\omega \rightarrow 0$) we may expect convergence to an improper integral:

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) \exp(i\omega x) d\omega$$

with

$$F(\omega) = \int_{-\infty}^{\infty} f(x) \overline{\exp(i\omega x)} dx$$


Definition:

$$F(\omega) := \mathcal{F}\{f(x)\} := \int_{-\infty}^{\infty} f(x) \exp(-i\omega x) dx$$


is called **Fourier transform** (or spectral function) of $f : \mathbb{R} \rightarrow \mathbb{C}$, if the integral exists for all $\omega \in \mathbb{R}$.

$$\mathcal{F}^{-1}\{F(\omega)\} := f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) \exp(i\omega x) d\omega$$

is called **inverse Fourier transform** of $F : \mathbb{R} \rightarrow \mathbb{C}$, if the integral exists for all $x \in \mathbb{R}$.



- Linearity $\mathcal{F}\{c_1 f_1(x) + c_2 f_2(x)\} = c_1 \mathcal{F}\{f_1(x)\} + c_2 \mathcal{F}\{f_2(x)\}$
 $= c_1 F_1(\omega) + c_2 F_2(\omega)$
 $c_1, c_2 \in \mathbb{R}$
- Displacement
- Similarity
- Derivatives $\mathcal{F}\{f'(t)\} = i\omega \mathcal{F}\{f(t)\} = i\omega F(\omega)$
- Modulation



Laplace transform

$$F(s) := \mathcal{L}\{f(t)\} := \int_0^\infty f(t) \exp(-st) dt,$$


is called **Laplace transform** of $f : \mathbb{R}^+ \rightarrow \mathbb{R}$,
if the integral exists for **some** $s \in \mathbb{R}$.

The inverse Laplace transform is hard to calculate.
Usually, reference tables are used.

The Laplace transform has similar properties as the
Fourier transform,
in particular

$$\mathcal{L}\{y'(t)\} = s\mathcal{L}\{y(t)\} - y(0) = sY(s) - y(0)$$

transforming differential equations (incl. initial values) into
algebraic equations



- 1 Introduction
- 2 Power series
- 3 Differentiation in Higher Dimensions
- 4 Integration in Higher Dimensions
- 5 Further Topics in Calculus
- 6 **Summary - Outlook and Review**
 - Insight and outlook
 - References

Introduction

Power series

Differentiation in
Higher Dimensions


Integration in
Higher Dimensions

Further Topics in
Calculus

**Summary -
Outlook and
Review**

Insight and outlook

References



Outlook and insight: What we did in Analysis 2

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions


Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References

- **Differential calculus in higher dimensions (mainly 2d, 3d)**
- **Integral calculus in higher dimensions (mainly 2d, 3d)**
- Sequences of functions
- **Power series,**
Taylor series/Taylor expansions
Fourier series
- ...



What we did in Analysis 2 - More Details

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions


Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References


- Sequences of functions, uniform convergence
- Power series, especially Taylor series, Fourier series, discrete Fourier transform
- Limits and continuity (for any d), curves (application: ECC)
- Differential calculus (for any d), especially optimization, linear regression
- Integrals along curves, on sectors; solids of revolution
- Integral calculus (for any d)
- Vector analysis (brief insight)
- Fourier transform (very brief insight)



What we have not discussed here

- Total differential, implicit functions, implicit differentiation
- Optimization with constraints, Lagrange multipliers
- Curvature
- **Differential equations**
- Integrals over curved surfaces, ...
- Laplace transform
- Fundamental theorem of Algebra to be proved by analytical methods ;)
- ...

... and all topics of this lecture may be discussed with all proofs and more details and extensions



References: recommended literature

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References



James Stewart: *Calculus*. Brooks/Cole, 6th edition, 2009.



J. Hass, C. Heil, M.D. Weir: *Thomas' Calculus: Early transcendentals*. Pearsons, 14th edition, 1999.

In German only:



Forster, O.: *Analysis 1*. Springer-Spektrum, 11. Auflage, 2013.



Forster, O.: *Analysis 2*. Springer-Spektrum, 11. Auflage, 2017.




Forster, O.: *Analysis 3*. Springer-Spektrum, 8. Auflage, 2017.



Bronstein, Semendjajew, Musiol, Mühlig: *Taschenbuch der Mathematik*, Verlag Harri Deutsch, Frankfurt a. M./Thun, 4., überarb. u. erw. Aufl. der Neubearb., 1999.

Formula collections are also available in English.



Further sources (for sake of completeness)

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions


Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References

-  Kimmerle, S.-J., Dvorsky, K., Ließ, H.-D., Avenhaus, R.: Time-to-Failure under Varying Thermal Stresses, Preprint, 2022.
-  Gerdts, M.: *Mathematik II*, Vorlesungsskript, Bachelorstudiengänge Luft- und Raumfahrttechnik, Elektrotechnik und Bau- und Umweltingenieurwesen, Universität der Bundeswehr München, Herbsttrimester 2014.
-  Fetzer, A., Fränkel, H.: *Mathematik 1*, Springer, Berlin/Heidelberg, 6. Aufl., 2000.
-  Jäger W.: *Analysis I-III*, Vorlesungsskripte, Diplom-Studiengänge Mathematik und Physik, Universität Heidelberg, Wintersemester 1998/99 – Wintersemester 1999/2000.
-  Kimmerle, W., Stroppel, M.: *Analysis für Ingenieure, Mathematiker und Physiker*, edition Delkhofen, 4. Aufl., 1., korr. Nachdruck, 2015.
-  Wellisch, U.: *Mathematik 2 Maschinenbau*, lecture notes, TH Rosenheim, 2021.
-  Meyberg, K., Vachenaue, P.: *Höhere Mathematik 1*, Springer, 6. korr. Aufl., 1. korr. Nachdruck, 2003.
-  Papula, L.: *Mathematik für Ingenieure und Naturwissenschaftler, Band 1*, Springer Vieweg, Wiesbaden, 14. Auflage, 2014.
-  Papula, L.: *Mathematische Formelsammlung*, Springer Vieweg, Wiesbaden, 11. Auflage, 2014.
-  Tapken, G., Weiß, M.: *Mathematik für Informatiker*, Vorlesungsskript, BSc.-Studiengänge “Informatik” und “Wirtschaftsinformatik”, OTH Regensburg, 2020.
-  Schneeberger, S.: *Analysis 1 & 2*, Vorlesungsskripte, BSc.-Studiengang WMA, TH Rosenheim, 2020/21.
-  Stingl, P.: *Mathematik für Fachhochschulen, Technik und Informatik*, Hanser, München, 8., aktual. Auflage, 2009.



General mathematical software:

- SCILAB, free software, `scilab.org`)
- MATLAB, campus licence
- Wolfram Mathematica
- ...

Computer algebra, i.e. “symbolic calculators”:

- MAPLE
- MATLAB, symbolic toolbox, campus licence ???
- Wolfram Alpha
- ...

For optimization, finite element methods, machine learning,
... specialized software packages exist.

Programming languages as Python, Java, C, ... are always
useful.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References

