

Technische
Hochschule
Rosenheim



Analysis 2

Priv.-Doz. Dr. Sven-Joachim Kimmerle

Summer term 2022
Bachelor Applied Artificial Intelligence (AAI)

- 1 Introduction
 - Lecturer
 - Motivation
 - Administrative and organisational matters
- 2 Power series
- 3 Differentiation in Higher Dimensions
- 4 Integration in Higher Dimensions
- 5 Further Topics in Calculus
- 6 Summary - Outlook and Review

Introduction

Lecturer
Motivation
Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



- 2000: “Vordiplom” in Mathematics & “Vordiplom” in Physics (U Heidelberg)
- 2002: Maîtrise in Mathematics (U Paris 7, France)
- 2004: Diploma in Mathematics (U Heidelberg)
- 2004-2009: Research center MATHEON, Berlin
- 2009: PhD in Mathematics (HU Berlin)
- 2010: Toyota/U Ottawa, Ottawa, Canada
- 2011-2018: Postdoc & deputy professor, UniBw München, Neubiberg
- 2019: “Habilitation” in Mathematics (UniBw München, Neubiberg)
- Since 2018: Physical Software Solutions GmbH, Münsing & Ottobrunn
- Since 2021: Lecturer (part-time), TH Rosenheim

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



What do you expect from the lecture?

Analysis 2

S.-J. Kimmerle

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Teaching mathematical basics (continued):

- Review of mathematics from Analysis 1 (& “school”)
- Completion of power series; Taylor series
- Fourier series
- Differentiation in higher dimensions
- Integration in higher dimensions
- Further topics: vector calculus, integral transformations (shortly)
- Mathematical thinking, techniques & working

In parallel this semester: Linear Algebra

- Systems of equations, matrices & vectors, eigenvalues, vector spaces, . . .

Introduction

Lecturer

Motivation

Administrative and organisational matters

Power series

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



Administrative & organisational matters 1

Analysis 2

S.-J. Kimmerle

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

4 SWS: ~ 2.67 hours lecture with ~ 1.33 hours exercise

- Lecture (2-3 hours per week)

Thursday, 09:45-11:15 (every week) in A3.14

Thursday, 11:45-13:15 (begin of the semester) in
A3.14 with integrated exercises

In presence (at least for the moment ...).

Please wear masks at all times!

- 2 exercise groups:

We start at the middle of the semester with 2
exercise groups instead of the 2nd lecture block.

Thursday, 11:45-13:15 in A3.14

Thursday, 13:45-15:15 in A3.14

In presence

Please register **later** for a group in the LC!

In case of (technical) issues, **we wait for 20 minutes!**



- Presentations, exercises and other material can be found in the Learning Campus (LC)
 - learning-campus.th-rosenheim.de
 - → Department ANG
 - Mathematics
 - Analysis 2 (AAI - B2), SoSe 2022
 - shortly: “Analysis 2 AAI, 22”
 - login: Kennelch!
- Office hours & contact
 - After each lecture/exercise group or
 - some time Wednesday afternoons or on Thursday
 - by appointment by email:
`sven-joachim.kimmerle@th-rosenheim.de`



● Examination type

- Written exam: 90 min.
- Auxiliary tools: 1 sheet (DIN A4) both sides, hand written with formulas, e.g.
- No calculators (or smartphones etc.) will be permitted.

● Homework and bonus system

- Marked homework (bonus), sometimes in groups up to 2
- To hand-in each Friday morning online, discussion next Thursday



According to the module handbook for 5 ECTS we expect a workload of about 150 hours:

- 60 hours contact (in presence or virtual):
~ 40 hours lecture, ~ 20 hours exercise
- 90 hours independent study

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Preliminaries:

- Good math skills from school or previous semesters
- Sound understanding of English
- Perseverance and endurance

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Recommended literature

Analysis 2

S.-J. Kimmerle

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

 James Stewart: *Calculus*. Brooks/Cole, 6th edition, 2009.

 J. Hass, C. Heil, M.D. Weir: *Thomas' Calculus: Early
transcendentals*. Pearsons, 14th edition, 1999.

In German only:

 Forster, O.: *Analysis 1*. Springer-Spektrum, 11. Auflage, 2013.

 Forster, O.: *Analysis 2*. Springer-Spektrum, 11. Auflage, 2017.

 Forster, O.: *Analysis 3*. Springer-Spektrum, 8. Auflage, 2017.

Further literature and material (software, e.g.) will be given during the course.



Copying ban

All materials made available in this lecture have been protected by me with a password, which has only been made available to the registered participants of this course.

Any form of distribution is prohibited!

Sven-Joachim Kimmerle

Analysis 2

S.-J. Kimmerle

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



S.-J. Kimmerle

Introduction

Lecturer

Motivation

Administrative and
organisational matters

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

• • •



1 Introduction

2 Power series

- Sequences of Functions
- Uniform Convergence
- Continuity and Uniform Convergence
- Power Series
- Taylor Series
- Fourier Series
- Discrete Fourier Transform

3 Differentiation in Higher Dimensions

4 Integration in Higher Dimensions

5 Further Topics in Calculus

6 Summary - Outlook and Review

Introduction

Power series

Sequences of Functions
Uniform Convergence
Continuity and Uniform
Convergence
Power Series
Taylor Series
Fourier Series
Discrete Fourier Transform

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



We know the concepts of functions, sequences, and series.

We are interested in series representing a function $f(x)$ at every x :

$$f(x) = \sum_{j=0}^{\infty} a_j (x - x_0)^j$$

For this purpose we need the concept of a sequence of functions at first.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



A sequence of elements of \mathbb{R} (cf. Analysis 1):

$$g : \mathbb{N} \rightarrow \mathbb{R}, n \mapsto a_n =: g(n)$$

Definition (Sequences of Functions)

Let D be a set. A mapping

$$g : D \times \mathbb{N} \rightarrow \mathbb{R}, (x, n) \mapsto f_n = f_n(x) =: g(x, n)$$

is called a **sequence** of functions $f_n : D \rightarrow \mathbb{R}, n \in \mathbb{N}$.

Other notations by writing the functions, e.g., are:

$$\{f_n\}_{n \in \mathbb{N}} = \{f_n\}_{n \geq 1} = \{f_1, f_2, f_3, \dots, f_n, \dots\}$$

The domain of definition D and the target area, here \mathbb{R} , have to be identical for all functions f_n .

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Example (Pointwise Convergence)

Consider $\{f_n\}_{n \in \mathbb{N}}$, $f_n : [0, 2] \rightarrow \mathbb{R}$ with

$$f_n(x) = \begin{cases} n^2 x, & 0 \leq x \leq \frac{1}{n}, \\ 2n - n^2 x, & \frac{1}{n} \leq x \leq \frac{2}{n}, \\ 0, & \frac{2}{n} \leq x \leq 2. \end{cases}$$

This example exhibits that we may not swap the limit and the integral (another limit process) in general!

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Definition (Pointwise Convergence)

Let D be a set. A sequence $\{f_n\}_{n \geq n_0}$ of functions $f_n : D \rightarrow \mathbb{R}$ is called **pointwise convergent** to a function $f : D \rightarrow \mathbb{R}$, if and only if

$$\lim_{n \rightarrow \infty} f_n(x) = f(x) \quad \text{for any } x \text{ in } D.$$

Equivalently,

For any $x \in D$ and $\varepsilon > 0$
there exists a $N = N(x, \varepsilon) \geq n_0$ s.t.:

$$|f_n(x) - f(x)| < \varepsilon \quad \text{for any } x \text{ in } D \text{ and all } n \geq N.$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Definition (Uniform Convergence)

Let D be a set. A sequence $\{f_n\}_{n \geq n_0}$ of functions $f_n : D \rightarrow \mathbb{R}$ is called **uniformly convergent** to a function $f : D \rightarrow \mathbb{R}$, if and only if

For any $x \in D$ and $\varepsilon > 0$ there exists a $N = N(\varepsilon) \geq n_0$ s.t.:

$$|f_n(x) - f(x)| < \varepsilon \quad \text{for any } x \text{ in } D \text{ and all } n \geq N.$$

Notice that N may depend only on ε but not on the point x .

Uniform convergence always implies **pointwise** convergence, the opposite is not true (see last example).

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Example Uniform Convergence

Let $x \in D = [0, 1)$. The sequence

$$\left\{ \frac{1}{2^{x+n}} \right\}_{n \in \mathbb{N}}$$

converges uniformly:



Theorem (Uniform convergence preserves continuity)

Let $D \subseteq \mathbb{C}$ and $f_n : D \rightarrow \mathbb{C}$, $n \in \mathbb{N}$, a sequence of continuous functions, that uniformly converge to a function $f : D \rightarrow \mathbb{C}$, then f is continuous.

The limit of a uniformly convergent sequence of continuous functions, is again continuous.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Example: Saw-tooth function

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Definition (Uniform norm or sup norm)

Let D be a set and $f : D \rightarrow \mathbb{C}$.

We set

$$\|f\|_D := \sup_{x \in D} |f(x)|.$$

$\|\cdot\|_D$ defines a norm on D .

A function f is bounded iff $\|f\| < \infty$.

When misunderstandings are excluded, we just write $\|f\|$ instead of $\|f\|_D$.

[Introduction](#)[Power series](#)[Sequences of Functions](#)[Uniform Convergence](#)[Continuity and Uniform Convergence](#)[Power Series](#)[Taylor Series](#)[Fourier Series](#)[Discrete Fourier Transform](#)[Differentiation in Higher Dimensions](#)[Integration in Higher Dimensions](#)[Further Topics in Calculus](#)[Summary - Outlook and Review](#)

By this notation we may reformulate the uniform convergence:

$\{f_n\}_{n \in \mathbb{N}}$ converges uniformly on D

$$\iff \lim_{n \rightarrow \infty} \|f_n - f\|_D = 0$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Criterion for Uniform Convergence of a Series

Analysis 2

S.-J. Kimmerle

Theorem (Weierstrass Convergence Criterion)

Let $f_n : D \rightarrow \mathbb{C}$, $n \in \mathbb{N}$.

If

$$\sum_{n=0}^{\infty} \|f_n\|_D < \infty$$

then the series

$$\sum_{n=0}^{\infty} f_n$$

converges absolutely and uniformly on D to a function $F : D \rightarrow \mathbb{C}$.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Example: Convergence of a Power Series

The series

$$\sum_{n=1}^{\infty} \frac{\cos(nx)}{n^2}$$

converges uniformly on \mathbb{R} .



Polynomials are among the functions that are easy to handle (e.g. for a machine).

Many other functions become manageable, when they are approximated by polynomials, i.e. as power series.

As for polynomials it is helpful to consider power series on \mathbb{C} from the start.

Definition (Power Series)

Let $\{a_j\}_{j \in \mathbb{N}}$ be a complex sequence and $z_0 \in \mathbb{C}$.

Then

$$f(z) = \sum_{j=0}^{\infty} a_j (z - z_0)^j = a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots$$

is called a **(complex) power series** with the **center** of the series z_0 .

[Introduction](#)[Power series](#)[Sequences of Functions](#)[Uniform Convergence](#)[Continuity and Uniform
Convergence](#)[Power Series](#)[Taylor Series](#)[Fourier Series](#)[Discrete Fourier Transform](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Properties of Power Series I

The set of points z where the power series converges form a set $M \subseteq \mathbb{C}$.

Note that $z_0 \in M$.

By this the power series defines a function $f : M \rightarrow \mathbb{C}$.

The partial sums of power series are polynomials (multiply out!).

Power series have excellent properties of convergence.



Theorem (Radius of Convergence)

Let

$$f(z) = \sum_{j=0}^{\infty} a_j (z - z_0)^j$$

be a complex power series.

Then exactly one of the following 2 cases holds:

- There exists a $\rho \in \mathbb{R}_0^+$ s.t. the series converges absolutely for all $z \in O_\rho(z_0) = \{z \in \mathbb{C} \mid |z - z_0| < \rho\}$ and diverges for all z with $|z - z_0| > \rho$.
- The series converges absolutely for all $z \in \mathbb{C}$.

$\rho \in \mathbb{R}_0^+ \cup \{+\infty\}$ is called **radius of convergence**,
 $O_\rho(z_0)$ is called **circle of convergence**.

For $|z| = \rho$ no general statement on convergence/divergence is possible.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Assume for a power series holds

$$\overline{\lim}_{j \rightarrow \infty} \sqrt[j]{|a_j|} = b \quad \text{or} \quad \lim_{j \rightarrow \infty} \left| \frac{a_{j+1}}{a_j} \right| = b,$$

where $b \in \mathbb{R}_0^+ \cup \infty$, then:

- If $b = 0$, then $\rho = +\infty$.
- If $b = +\infty$, then $\rho = 0$.
- If $0 < b$, then $\rho = \frac{1}{b}$.

The limes superior (or inferior) of the quotient is not helpful in general.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Example: Geometric Series

Geometric series:

$$\sum_{j=0}^{\infty} z^n$$

Set $a_j = 1$ for all $j \in \mathbb{N}$ (and $z_0 = 0$).

Since $b = \lim_{j \rightarrow \infty} \sqrt[j]{a_j} = 1$, we find $\rho = \frac{1}{b} = 1$.

Thus we have (as expected) convergence for $|z| < 1$ and divergence for $|z| > 1$.

What happens for $|z| = 1$? Divergence, since we do not encounter a zero sequence z^n .

Moreover $\sum_{j=0}^{\infty} z^n = \frac{1}{1-z}$ for $|z| < 1$.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Theorem (Addition & Scaling of Power Series)

Consider two power series

$f(z) = \sum_{n=0}^{\infty} a_n z^n$ with radius of convergence ρ_f and
 $g(z) = \sum_{n=0}^{\infty} b_n z^n$ with radius of convergence ρ_g .

*Then the sum/difference is given for all z with
 $|z| < \min(\rho_f, \rho_g)$ by:*

$$\sum_{n=0}^{\infty} (a_n \pm b_n) z^n = \sum_{n=0}^{\infty} a_n z^n \pm \sum_{n=0}^{\infty} b_n z^n = f(z) \pm g(z)$$

*Further, the scaling i.e. multiplication with a factor $c \in \mathbb{C}$ is
given for all z with $|z| < \rho_f$ by:*

$$\sum_{n=0}^{\infty} c a_n z^n = c \sum_{n=0}^{\infty} a_n z^n = c f(z).$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem ((Cauchy) Product of Power Series)

Consider two power series

$f(z) = \sum_{n=0}^{\infty} a_n z^n$ with radius of convergence ρ_f and
 $g(z) = \sum_{n=0}^{\infty} b_n z^n$ with radius of convergence ρ_g .

Then the product $f(z_1)g(z_2)$ is given for all z_1, z_2 with $|z_1|, |z_2| < \min(\rho_f, \rho_g)$ by:

$$f(z_1)g(z_2) = \sum_{n=0}^{\infty} \sum_{k=0}^n a_k b_{n-k} z_1^k z_2^{n-k}.$$

In particular, if $z = z_1 = z_2$:

$$f(z)g(z) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n a_k b_{n-k} \right) z^n.$$

The latter 2 theorems also hold for $f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$,
 $g(z) = \sum_{n=0}^{\infty} b_n (z - z_0)^n$ by a shift.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem (Series of polynomials are continuous)

Let

$$f(z) = \sum_{j=0}^{\infty} a_j (z - z_0)^j$$

be a complex power series with radius of convergence ρ .

Then the function

$$f : O_{\rho}(z_0) \rightarrow \mathbb{C} : z \mapsto \sum_{j=0}^{\infty} a_j (z - z_0)^j$$

is continuous.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



More Examples: Exponential Series etc.

The series
for the exponential function and
for (co)sine, resp.,
have the radius of convergence $\rho = \infty$.

The latter 3 series yield a continuous function.



Theorem (Identity Theorem for Power Series)

Let

$$f(z) = \sum_{j=0}^{\infty} a_j(z - \zeta)^j, \quad g(z) = \sum_{j=0}^{\infty} b_j(z - \zeta)^j$$

be complex power series, both with radius of convergence ρ .

Suppose there exists a series of points $z_n \in O_\rho(\zeta) \setminus \zeta$, $n \in \mathbb{N}$, s.t.

$$f(z_n) = g(z_n) \text{ for all } n \in \mathbb{N} \text{ and } \lim_{n \rightarrow \infty} z_n = \zeta,$$

then f and g are identical, i.e. $a_j = b_j$ for all $j \in \mathbb{N}$.

Remark: Power series may be extended from the real axis to the complex plane \rightsquigarrow complex analysis (“Funktionentheorie”).

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Proof of the Identity Theorem

Analysis 2

S.-J. Kimmerle

Proof:

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



We know an example where the limit of f_n and the integration do not commute.

But in case of uniform convergence we may swap the limit of the sequence and the integration:

Theorem (Commutation of integration and limit of a sequence)

Let $f_n : [a, b] \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, a sequence of continuous functions.

Let $\{f_n\}_{n \in \mathbb{N}}$ be uniformly convergent on $[a, b]$ to $f : [a, b] \rightarrow \mathbb{R}$.

Then:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \int_a^b f_n(x) dx.$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Proof: Commutation of Integration & Limit

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Examples for Commutation of Integration & Limit

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

- Check:

$$\begin{aligned}\int_a^b e^x dx &= \int_a^b \sum_{j=0}^{\infty} \frac{x^j}{j!} dx = \sum_{j=0}^{\infty} \int_a^b \frac{x^j}{j!} dx = \\ \sum_{j=0}^{\infty} \left[\frac{x^{j+1}}{(j+1)!} \right]_a^b &= \left[\sum_{j=0}^{\infty} \frac{x^{j+1}}{(j+1)!} \right]_a^b = \left[\sum_{k=1}^{\infty} \frac{x^k}{k!} \right]_a^b = [e^x]_a^b\end{aligned}$$

- Power series for the integral sine:



We ask what are the conditions that the limit of f_n and the differentiation commute.

We require the pointwise convergence of f_n and the uniform convergence of f'_n :

Theorem (Commutation of differentiation and limit of a sequence)

Let $f_n : [a, b] \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, a sequence of continuously differentiable functions, with derivatives $f'_n : [a, b] \rightarrow \mathbb{R}$.

Let $\{f_n\}_{n \in \mathbb{N}}$ be pointwise convergent on $[a, b]$ to $f : [a, b] \rightarrow \mathbb{R}$ and $\{f'_n\}_{n \in \mathbb{N}}$ be uniformly convergent on $[a, b]$.

Then f is differentiable and:

$$f'(x) = \lim_{n \rightarrow \infty} f'_n(x) \quad \text{for all } x \in [a, b].$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Proof: Commutation of Differentiation & Limit

Analysis 2

S.-J. Kimmerle

Proof.

Let $f^\star := \lim_{n \rightarrow \infty} f'_n$. According to the result from above f^\star is continuous on $[a, b]$. Thus for all $x \in [a, b]$ there holds

$$f_n(x) = f_n(a) + \int_a^x f'_n(\xi) d\xi. \quad (1)$$

According to the last theorem

$$\int_a^x f'_n(\xi) d\xi \rightarrow \int_a^x f^\star(\xi) d\xi \text{ as } n \rightarrow \infty,$$

thus taking the limit on both sides of (1) yields

$$f(x) = f(a) + \int_a^x f^\star(\xi) d\xi.$$

By differentiation $f'(x) = f^\star(x)$ for any $x \in [a, b]$. □

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

$$f_n(x) = \frac{1}{n} \sin(nx), \quad n \in \mathbb{N}$$

$$F(x) = \sum_{n=1}^{\infty} \frac{\cos(nx)}{n^2}$$



Corollary

Suppose the power series

$$f(x) = \sum_{n=0}^{\infty} c_n (x - x_0)^n$$

converges on $I := (x_0 - \rho, x_0 + \rho)$, where $\rho > 0$ is the radius of convergence.

Then $f : I \rightarrow \mathbb{R}$ is arbitrarily many times differentiable and:

$$c_n = \frac{1}{n!} f^{(n)}(x_0)$$

Proof: Apply iteratively the theorem on commutation of differentiation and limits.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

In many applications “higher order” terms are neglected in order to find simpler or manageable formulas.

The base for this and for an estimate of the resulting error is the following theorem:

Theorem (Taylor theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be an $n + 1$ -times continuously differentiable function.

Then for any two points x and x_0 in (a, b) there exists a number $\Theta_{x,x_0} \in (0, 1)$ such that

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \frac{f^{(n+1)}(x_0 + \Theta_{x,x_0} \cdot (x - x_0))}{(n+1)!} (x - x_0)^{n+1}.$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Definition (Taylor Polynomial, Lagrange Form of the Remainder)

We call

$$T_n(f, x, x_0) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$$

the **Taylor polynomial of n-th order** (not of n-th degree!) around the center x_0 .

The term

$$\frac{f^{(n+1)}(x_0 + \Theta_{x,x_0} \cdot (x - x_0))}{(n+1)!} (x - x_0)^{n+1}$$

is called the **Lagrange form of the remainder**.

There are other forms of the remainder: Peano form, Cauchy form, ...

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Examples for the Taylor Theorem

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Definition (Taylor Series)

Let $f : [a, b] \rightarrow \mathbb{R}$ be arbitrarily many times differentiable.

Then for $x \in [a, b]$ the power series

$$T(f, x, x_0) = \sum_{k=0}^{\infty} \frac{f^{(k)}(\mathbf{x}_0)}{k!} (x - x_0)^k.$$

is called the **Taylor series** of the function f with the center x_0 .

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem (Convergence Criterion for Taylor Series)

The Taylor series $T(f, x, x_0)$ converges for a given $x \in [a, b]$ to $f(x)$

iff

$$\lim_{n \rightarrow \infty} R_n(f, x, x_0) = 0.$$

We are interested in an interval $[a, b]$ such that the remainder converges to 0 and this should happen “fast”.

Warning: There exist functions whose Taylor series represent the function in a single point only!

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



An important special case: $x_0 = 0$.

Let f be as in the Taylor theorem and $0 \in (a, b)$. For any point $x \in (a, b)$ there exists a number $\Theta_x \in (0, 1)$ such that

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(0)}{k!} x^k + \frac{f^{(n+1)}(\Theta_x x)}{(n+1)!} x^{n+1}.$$

The corresponding series is called **Maclaurin series** of the function f :

$$M(f, x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k.$$

Its center is $x_0 = 0$.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

A Taylor Series for the Logarithm

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions


Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

$$\bullet \ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$$

$$\bullet \ln(2) = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots$$



Motivation:

- Periodic signals, image processing (jpeg/mpeg compression)
- Generation of signals by periodic means
- Harmonic oscillations, acoustics
- Periodic orbits
- Revolving motors

Many periodic functions may be approximated by a sum of trigonometric base functions as

$$1, \quad \cos(kx), \quad \sin(kx), \quad k \in \mathbb{N}$$

with suitable coefficients a_0, a_k, b_k .

Frequently, it may make sense to replace the finite sum by a series.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Example: Rotational Force of an Engine I

A rotational force f (due to gas pressure and mass inertia) acts on the crankshaft of a single-cylinder two-stroke engine.

At the points $t_k = k \cdot \pi/8$, $k = 0, 1, \dots, 15$, the following values of f_k were determined experimentally:

f_0	f_1	f_2	f_3	f_4	f_5
-8250	-29430	-2286	5974	-8829	-25408
f_6	f_7	f_8	f_9	f_{10}	
-22681	-28655	-8564	96560	45862	
f_{11}	f_{12}	f_{13}	f_{14}	f_{15}	
22092	-9025	-23514	-15127	12880	

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform


Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Example: Rotational Force of an Engine II

Analysis 2

S.-J. Kimmerle

The resulting trigonometric approximation of f looks as follows:



Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Definition (Periodic function)

A function $f : \mathbb{R} \rightarrow \mathbb{C}$ is called **T -periodic**, if there exists a $T > 0$ s.t.

$$f(x + T) = f(x) \quad \text{for all } x \in \mathbb{R}.$$

T is called a **period** of f .

Note that a T -periodic function f is completely defined by its values on the interval $[0, T)$. Thus it is enough to consider f only on $[0, T)$.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Let f and g be T -periodic.

- Then $\alpha f + \beta g$ is T -periodic for any $\alpha, \beta \in \mathbb{C}$.
- Then there holds for all $c \in \mathbb{R}$:

$$\int_c^{c+T} f(x) dx = \int_0^T f(x) dx.$$

- Then the function $\tilde{f} : \mathbb{R} \rightarrow \mathbb{C}$, defined by

$$\tilde{f}(x) := f\left(\frac{T}{2\pi}x\right)$$

is 2π -periodic, since:

[Introduction](#)[Power series](#)[Sequences of Functions](#)[Uniform Convergence](#)[Continuity and Uniform
Convergence](#)[Power Series](#)[Taylor Series](#)[Fourier Series](#)[Discrete Fourier Transform](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Definition (Fourier Series)

The function series

$$F(x) := \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(k\omega x) + b_k \sin(k\omega x) \quad \text{with } \omega := \frac{2\pi}{T}$$

is called **Fourier series**. The coefficients a_0, a_1, \dots and b_1, b_2, \dots are called **Fourier coefficients**. The representation of a function by its Fourier series is called **Fourier** (or harmonic) **analysis**. The finite sum is called **Fourier sum**:

$$F_n(x) := \frac{a_0}{2} + \sum_{k=1}^n a_k \cos(k\omega x) + b_k \sin(k\omega x).$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem (Fourier)

Assume a T -periodic function f may be represented as a *Fourier series*

$$F(x) := \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(k\omega x) + b_k \sin(k\omega x) \quad \text{with } \omega := \frac{2\pi}{T},$$

then the Fourier coefficients are given by

$$a_0 = \frac{2}{T} \int_0^T f(x) dx,$$

$$a_k = \frac{2}{T} \int_0^T f(x) \cos(k\omega x) dx,$$

$$b_k = \frac{2}{T} \int_0^T f(x) \sin(k\omega x) dx.$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Symmetry & Fourier Coefficients

The following symmetries may be exploited when computing Fourier coefficients, $k \in \mathbb{N}$:

- If f is an odd function, i.e. $f(x) = -f(-x)$ for all $x \in \mathbb{R}$, then

$$b_k = \frac{2}{T} \int_0^T f(x) \sin(k\omega x) dx = \frac{4}{T} \int_0^{T/2} f(x) \sin(k\omega x) dx,$$
$$a_k = \frac{2}{T} \int_0^T f(x) \cos(k\omega x) dx = 0.$$

- If f is an even function, i.e. $f(x) = f(-x)$ for all $x \in \mathbb{R}$, then

$$b_k = \frac{2}{T} \int_0^T f(x) \sin(k\omega x) dx = 0,$$
$$a_k = \frac{2}{T} \int_0^T f(x) \cos(k\omega x) dx = \frac{4}{T} \int_0^{T/2} f(x) \cos(k\omega x) dx.$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Example: Saw-tooth Function revisited

Warning: In general $f(x) \neq F(x)$ for all $x \in \mathbb{R}$!

Gibbs phenomenon: overshooting & undershooting (up to 18% for large n) at points of discontinuity



Revision: Trigonometric Functions

Analysis 2

S.-J. Kimmerle

Definition (Cosine and sine)

For $x \in \mathbb{R}$ we define:

$$\cos(x) := \operatorname{Re}(\exp(ix))$$

$$\sin(x) := \operatorname{Im}(\exp(ix))$$

We see that the Euler formula holds:

$$\exp(ix) = \cos(x) + i \sin(x), \quad x \in \mathbb{R}$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



By the complex exponential function we may rewrite

$$F(x) = \sum_{k=-\infty}^{\infty} c_k \exp(ik\omega x) \quad \text{with } \omega := \frac{2\pi}{T}$$

where the complex Fourier coefficient c_k , $k \in \mathbb{Z}$, is given by

$$c_k = \frac{1}{T} \int_0^T f(x) \exp(-ik\omega x) dx.$$

Note that the limit is to be understood symmetrically:

$$\sum_{k=-\infty}^{\infty} c_k \exp(ik\omega x) := \lim_{n \rightarrow \infty} \sum_{k=-n}^n c_k \exp(ik\omega x).$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem (Orthonormality Relations)

If $m, n \in \mathbb{N}$, then:

$$\frac{2}{T} \int_0^T \sin(m\omega x) \sin(n\omega x) dx = \delta_{m,n},$$

$$\frac{2}{T} \int_0^T \cos(m\omega x) \cos(n\omega x) dx = \delta_{m,n},$$

$$\frac{2}{T} \int_0^T \sin(m\omega x) \cos(n\omega x) dx = 0,$$

and if $m, n \in \mathbb{Z}$, then:

$$\frac{1}{T} \int_0^T \exp(im\omega x) \exp(-in\omega x) dx = \delta_{m,n}.$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Definition (Piecewise Continuously Differentiable Function)

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ with

- (i) Only at a finite number of **singularities** the function f is not continuously differentiable.
- (ii) At any singularity x_0 there exist the following one-sided limits:

$$f(x_0+) := \lim_{x \rightarrow x_0+} f(x) \qquad f(x_0-) := \lim_{x \rightarrow x_0-} f(x),$$

$$f'(x_0+) := \lim_{x \rightarrow x_0+} f'(x) \qquad f'(x_0-) := \lim_{x \rightarrow x_0-} f'(x).$$

Then f is called **piecewise continuously differentiable**.

Discontinuities are singularities, but not any singularity is a discontinuity.

[Introduction](#)[Power series](#)[Sequences of Functions](#)[Uniform Convergence](#)[Continuity and Uniform](#)[Convergence](#)[Power Series](#)[Taylor Series](#)[Fourier Series](#)[Discrete Fourier Transform](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Theorem (Convergence of Fourier Series)

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a periodic function that is piecewise continuously differentiable. Then:

- *The Fourier series F converges at any x that is not a singularity to f .*
- *At any singularity x_0 the Fourier series converges to the “mean value” of the jump*

$$\frac{1}{2} (f(x_0+) + f(x_0-)).$$

- *In any compact interval that does not contain a discontinuity, the convergence of F to f is uniform.*

Note that there exist periodic, continuous functions, whose Fourier series does not converge to f !

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Properties of Fourier Series I

Let f and g be piecewise continuous, periodic functions with Fourier series $F = \sum_{k=-\infty}^{\infty} c_k \exp(ik\omega x)$, $G = \sum_{k=-\infty}^{\infty} d_k \exp(ik\omega x)$, resp. There holds:

- For any $\alpha, \beta \in \mathbb{R}$:

$$\alpha F(x) + \beta G(x) = \sum_{k=-\infty}^{\infty} (\alpha c_k + \beta d_k) \exp(ik\omega x)$$

- $F(-x) = \sum_{k=-\infty}^{\infty} c_{-k} \exp(ik\omega x)$
- For any $\alpha \in \mathbb{R}$

$$F(\alpha x) = \sum_{k=-\infty}^{\infty} c_k \exp(ik\omega \alpha x)$$

- For any $\alpha \in \mathbb{R}$

$$F(\alpha + x) = \sum_{k=-\infty}^{\infty} (c_k \exp(ik\omega \alpha)) \exp(ik\omega x)$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Properties of Fourier Series II

Let f be a piecewise continuously differentiable, periodic function with Fourier series $F = \sum_{k=-\infty}^{\infty} c_k \exp(ik\omega x)$. There holds:

- The Fourier series F' represents f' :

$$F'(x) = \sum_{k=-\infty}^{\infty} (ik\omega c_k) \exp(ik\omega x)$$

- $F(-x) = \sum_{k=-\infty}^{\infty} c_{-k} \exp(ik\omega x)$
- Suppose $c_0 = 0$, then the Fourier series $\tilde{F} := \int F(\xi) d\xi$ represents $\tilde{f} := \int f(\xi) d\xi$:

$$\tilde{F}(x) = \frac{1}{T} \int_0^T \tilde{f}(\xi) d\xi + \sum_{k=-\infty, k \neq 0}^{\infty} \frac{c_k}{ik\omega} \exp(ik\omega x)$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Problem (General Interpolation Task)

For a given function $g : \mathbb{R} \rightarrow \mathbb{R}$ with given points

$$(x_j, g(x_j)), \quad j = 1, \dots, n,$$

we search for a function p , s.t. the graph of p goes through these given points.

p is chosen within a class of reasonable (and manageable) functions, e.g.

- *the class of real polynomials,*
- *the class of trigonometric polynomials in case of T -periodic functions.*

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Trigonometric Approximation Revisited

Analysis 2

S.-J. Kimmerle

In the later case we consider a function $g : [0, 2\pi] \rightarrow \mathbb{R}$, periodically extended with $T = 2\pi$.

Let

$$x_0 = 0, x_1 = \frac{2\pi}{n}, \dots, x_{n-1} = (n-1)\frac{2\pi}{n}, x_n = 2\pi.$$

We search for a trigonometric polynomial

$$p_n(x) := \sum_{k=0}^{n-1} c_k \exp(ikx)$$

such that

$$p_n(x_j) = g(x_j) \quad \text{for } j = 0, \dots, n-1.$$

Remark: Due to the periodicity, we have $g(x_0) = g(x_n)$.

By inserting:

$$\sum_{k=0}^{n-1} \exp(ikx_j) c_k = g(x_j) \quad \text{for } j = 0, \dots, n-1.$$

This is a linear equation system (LES) with coefficient matrix

$$f_{jk} = \exp\left(2\pi i k \frac{j}{n}\right) = \zeta^{jk} \quad \text{with } \zeta := \exp\left(2\pi i \frac{1}{n}\right).$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



This motivates the definition of the so-called **Fourier matrix** :

$$F_n = (f_{ij})_{i,j=1,\dots,n}$$

Properties of the Fourier matrix:

- F_n is symmetric
- $\frac{1}{\sqrt{n}}F_n$ is unitary, i.e. $\left(\frac{1}{\sqrt{n}}F_n\right)^{-1} = \overline{\left(\frac{1}{\sqrt{n}}F_n\right)}^T$ (details later in LA)

Thus, we may rewrite the LES

$$F_n \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} = \begin{pmatrix} g(x_1) \\ \vdots \\ g(x_n) \end{pmatrix}$$

that uniquely determines the trigonometric polynomial p_n by

$$\begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} = \frac{1}{n} \overline{F}_n^T \begin{pmatrix} g(x_1) \\ \vdots \\ g(x_n) \end{pmatrix}$$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



The mapping

$$\mathbb{C}^n \rightarrow \mathbb{C}^n, v \mapsto F_n v$$

is called **discrete Fourier transform (DFT)**, i.e.

$$(F_n v)_j = \sum_{k=0}^{n-1} v_k \exp\left(2\pi i \frac{jk}{n}\right).$$

This may be interpreted as a discretization of the (continuous) Fourier transform (at the end of lecture).

The inverse DFT solves the trigonometric interpolation problem.

By an algorithm, called **fast Fourier transform (FFT)** the DFT may be computed with only "few" operations!

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



Example: Fourier Matrix

Let $v^T = (1, -1, 0, 1)$.

Compute

$$F_4 v =$$





An image with decreasing quality from the left to the right.
(Source: [Wikipedia](#))

Typical standards: `jpeg`, `mpeg`

An important step (among others) in these data compression methods is the discrete Fourier cosine transform that maps into the frequency domain.

Theoretically, this step is invertible, i.e. without loss of information possible.

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform
Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



At the end of a lecture (if time permits) we will have a short look at

- (Continuous) Fourier transform
- Laplace transform
- Short-time Fourier transform
- Wavelet

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform

Convergence

Power Series

Taylor Series

Fourier Series

Discrete Fourier Transform

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



- 1 Introduction
- 2 Power series
- 3 **Differentiation in Higher Dimensions**
 - Limits and Continuity in \mathbb{R}^n
 - Differentiability in \mathbb{R}^n
 - Optimization in \mathbb{R}^n
 - Introduction into Linear Regression
 - Curves
- 4 Integration in Higher Dimensions
- 5 Further Topics in Calculus
- 6 Summary - Outlook and Review

Introduction

Power series

**Differentiation in
Higher Dimensions**

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



In Analysis 1 we have discussed differentiation of functions of 1 variable.

Now we consider a real-valued function in several variables

$$f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\mathbf{x} = (x_1, \dots, x_n)^\top \mapsto f(x_1, \dots, x_n) = f(\mathbf{x})$$

Introduction

Power series

**Differentiation in
Higher Dimensions**Limits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

In Analysis 1 we have discussed differentiation of functions of 1 variable.

Now we consider a real-valued function in several variables

$$f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$$
$$\mathbf{x} = (x_1, \dots, x_n)^\top \mapsto f(x_1, \dots, x_n) = f(\mathbf{x})$$

Later we are going to extend the differential calculus to vector-valued functions (of several variables)

$$f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$$
$$\mathbf{x} = (x_1, \dots, x_n)^\top \mapsto (f_1(x_1, \dots, x_n), \dots, f_m(x_1, \dots, x_n))^\top$$



Special Case: $n = 2$

A real-valued function in 2 variables

$$f : D \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$$
$$\mathbf{x} = (x_1, x_2)^T \mapsto f(x_1, x_2)$$

We may plot the function value as 3rd coordinate over the real plane \mathbb{R}^2 .

The graph of f is a subset of \mathbb{R}^3 : a "landscape" or "mountains".



Definition (Level set)

We define the **level set** of a function $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$, $(x_1, \dots, x_n)^\top \mapsto f(x_1, \dots, x_n)$ for the function value $c \in \mathbb{R}$ as the set

$$N_c := \{\mathbf{x} \in D \mid f(\mathbf{x}) = c\}.$$

The structure of N_c may be "complicated", it might also be the empty set.

For $n = 2$ the level set is called a contour line or level curve (though it may be an area, e.g.),
for $n = 3$ the level set is called an equipotential surface (though it may be a solid, e.g.).

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Partial Function: Definition

A partial function is a "cross section"-function that is obtained by freezing all but 1 variables, e.g. x_i :

$$g_i : D_i \subseteq \mathbb{R} \rightarrow \mathbb{R}$$

$$x_i \mapsto f(a_1, \dots, a_{i-1}, x_i, a_{i+1}, \dots, a_n),$$

with $(a_1, \dots, a_{i-1}, x_i, a_{i+1}, \dots, a_n) \in D$ for all $x_i \in D_i$,
 $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n$ fixed.



Partial Function: Example

$$f(x_1, x_2) = x_1^2 + x_2^2$$

Analysis 2

S.-J. Kimmerle

Introduction

Power series

**Differentiation in
Higher Dimensions**

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n


Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Definition (Partial derivative)

Let $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$, $\mathbf{x} = (x_1, \dots, x_n)^\top \mapsto f(x_1, \dots, x_n)$ and $a \in D$, D an open set.

If the derivative of the partial function

$$x_i \mapsto f(a_1, \dots, a_{i-1}, x_i, a_{i+1}, \dots, a_n)$$

exists at $x_i = a_i$,


then we call it the **partial derivative** of f w.r.t. x_i at a .

We write:

$$\frac{\partial f}{\partial x_i}(\mathbf{a}) \quad \text{or} \quad f'_{x_i}(\mathbf{a}) \quad \text{or} \quad \dots$$

We say f is **partially differentiable** in \mathbf{a} , if all $\frac{\partial f}{\partial x_i}(\mathbf{a})$ exist.

We say f is partially differentiable in $E \subseteq D$, if f is partially differentiable at any $\mathbf{a} \in E$.



Partial Derivatives: Gradient

We assume that f is partially differentiable in D .

It is convenient to arrange the (first) partial derivatives in a vector.

We define the **gradient** of $f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}$ as the column vector (function):

$$\nabla f(\mathbf{x}) := \begin{pmatrix} \frac{\partial f}{\partial x_1}(\mathbf{x}) \\ \vdots \\ \frac{\partial f}{\partial x_i}(\mathbf{x}) \\ \vdots \\ \frac{\partial f}{\partial x_n}(\mathbf{x}) \end{pmatrix}$$

The gradient is

- orthogonal to level sets and
- points into the direction of the steepest ascent.



Partial Derivatives: Examples

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Higher Partial Derivatives: Definition

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

Partial derivatives w.r.t. some x_j are again functions of x_1, x_2, \dots, x_n and, again, we may search for partial derivatives w.r.t. some x_i , yielding **second partial derivatives**. ($i, j = 1, \dots, n$)

We write:

$$\frac{\partial}{\partial x_i} \frac{\partial f}{\partial x_j}(\mathbf{a}) \quad \text{or} \quad \frac{\partial^2}{\partial x_i \partial x_j} f(\mathbf{a}) \quad \text{or} \quad f''_{x_i, x_j}(\mathbf{a}) \quad \text{or} \quad f''_{ij}(\mathbf{a}) \quad \dots$$

In 2d with $(x, y) = (x_1, x_2)$ this reads, e.g.:

$$\frac{\partial}{\partial x} \frac{\partial f}{\partial y}(\mathbf{a}) \quad \text{or} \quad \frac{\partial^2}{\partial x \partial y} f(\mathbf{a}) \quad \text{or} \quad f''_{x, y}(\mathbf{a}) \quad \text{or} \quad f''_{1, 2}(\mathbf{a}) \quad \dots$$



Higher Partial Derivatives: Hesse Matrix

We assume that f is 2x partially differentiable in D .
Again it is convenient to order the second partial
derivatives in a (quadratic) matrix.

We define the **Hesse matrix** of $f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}$ as the
quadratic matrix (function):

$$H_f(\mathbf{x}) := \begin{pmatrix} \frac{\partial^2}{\partial x_1 \partial x_1} f(\mathbf{x}) & \dots & \dots & \frac{\partial^2}{\partial x_1 \partial x_n} f(\mathbf{x}) \\ \vdots & \frac{\partial^2}{\partial x_i \partial x_j} f(\mathbf{x}) & \ddots & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \frac{\partial^2}{\partial x_n \partial x_1} f(\mathbf{x}) & \dots & \dots & \frac{\partial^2}{\partial x_n \partial x_n} f(\mathbf{x}) \end{pmatrix}$$

In 2d with $(x, y) = (x_1, x_2)$ this reads, e.g.:

$$H_f(x, y) := \begin{pmatrix} \frac{\partial^2}{\partial x \partial x} f(x, y) & \frac{\partial^2}{\partial x \partial y} f(x, y) \\ \frac{\partial^2}{\partial y \partial x} f(x, y) & \frac{\partial^2}{\partial y \partial y} f(x, y) \end{pmatrix}$$



Schwarz-Clairaut Theorem

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

According to the Schwarz-Clairaut theorem there holds:

Suppose f is defined on a disk $D \subseteq \mathbb{R}^2$ with $(x_0, y_0) \in D$.

If $\frac{\partial^2 f}{\partial x \partial y}$ and $\frac{\partial^2 f}{\partial y \partial x}$ are both continuous on D ,

then $\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x}$.

This shows that the Hesse matrix is symmetric.



Higher Partial Derivatives: Examples

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Let $D \subseteq \mathbb{R}^n$.

- ε -neighbourhood of $\mathbf{a} \in \mathbb{R}^n$:

$$U_\varepsilon(\mathbf{a}) := \left\{ \mathbf{x} \in \mathbb{R}^n \mid \|\mathbf{x} - \mathbf{a}\|_2 := \sqrt{\sum_{i=1}^n (x_i - a_i)^2} < \varepsilon \right\}$$

- $\mathbf{a} \in D$ is called an **interior point** of $D : \iff$ there exists $U_\varepsilon(\mathbf{a})$ with $U_\varepsilon(\mathbf{a}) \subseteq D$.
- A set D is called **open** : \iff any point of D is an interior point. A set D is called **closed** : $\implies \mathbb{R}^n \setminus D$ is open.
- $\mathbf{b} \in \mathbb{R}^n$ is called a **boundary point** of $D : \implies$ for all $U_\varepsilon(\mathbf{b})$ holds, there exists a $\mathbf{x}, \mathbf{y} \in U_\varepsilon(\mathbf{b})$ with $\mathbf{x} \in D$ and $\mathbf{y} \notin D$. The set ∂D of all boundary points \mathbf{b} is called **boundary** of D .

Examples, see blackboard

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Revision/Definition: Limits and Continuity

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

Let $f : \mathbb{R}^n \subseteq D \rightarrow \mathbb{R}^m$, $\mathbf{a} \in D \cup \partial D$.

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} f(\mathbf{x}) = \mathbf{c} \in \mathbb{R}^m : \Longleftrightarrow$$

for all $\varepsilon > 0$ exists $U_\varepsilon(\mathbf{a}) : \|f(\mathbf{x}) - \mathbf{c}\| \leq \varepsilon$ for all $\mathbf{x} \in D \cap U_\varepsilon(\mathbf{a})$.

f is called **continuous** in $\mathbf{a} \in D : \Longleftrightarrow \lim_{\mathbf{x} \rightarrow \mathbf{a}} f(\mathbf{x}) = f(\mathbf{a})$.

f is called continuous in $D : \Longleftrightarrow f$ is continuous for all $\mathbf{a} \in D$.

Example: For $n = 2, m = 1$, f is continuous means that there are no “jumps” or “ridges”.



Pathologic Example for (Dis)continuity in 2d

Functions of severable variables exhibit phenomena that we do not encounter in 1d:

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f(x, y) := \begin{cases} \frac{2xy}{x^2+y^2}, & \text{if } x \neq 0 \text{ or } y \neq 0, \\ 0, & \text{if } (x, y)^T = (0, 0)^T. \end{cases}$$

A limit in 2d has to hold for any “path” $(x, y) \rightarrow (x_0, y_0)$.
It is not enough to consider each coordinate separately!



In this subsection we consider the total differentiability, a stronger requirement than partial differentiability.

Moreover, we consider expansions of functions of several variables.

Applications:

- Linear approximation of functions, tangent planes
- Taylor series for several variables

Here we consider:

$$f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \mapsto \begin{pmatrix} f_1(\mathbf{x}) \\ \dots \\ f_m(\mathbf{x}) \end{pmatrix} = \begin{pmatrix} f_1(x_1, \dots, x_n) \\ \dots \\ f_m(x_1, \dots, x_n) \end{pmatrix}$$

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Limits and Continuity in \$\mathbb{R}^n\$](#) [Differentiability in \$\mathbb{R}^n\$](#) [Optimization in \$\mathbb{R}^n\$](#) [Introduction into Linear
Regression](#)[Curves](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Definition (Total differentiability)

$f : \mathbb{R}^n \supseteq D \rightarrow \mathbb{R}^m$ is called **totally differentiable** (or linearly approximable) in $\mathbf{x} \in D$,

if there exists a matrix $A \in \mathbb{R}^{m \times n}$,
such that for all \mathbf{x} in a neighbourhood $U(\mathbf{x}_0)$, $\mathbf{x}_0 \in D$,

$$f(\mathbf{x}) = f(\mathbf{x}_0) + A \cdot (\mathbf{x} - \mathbf{x}_0) + R(\mathbf{x} - \mathbf{x}_0),$$

where

$$R : \mathbb{R}^n \supseteq U(\mathbf{x}_0) \rightarrow \mathbb{R}^m \text{ with } \lim_{\|\mathbf{x} - \mathbf{x}_0\| \rightarrow 0} \frac{R(\mathbf{x} - \mathbf{x}_0)}{\|\mathbf{x} - \mathbf{x}_0\|} = 0.$$

f is called totally differentiable in E , if this holds for all $\mathbf{x} \in E \subseteq D$.

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

If f is totally differentiable, then

- f is continuous
- all f_i , $i = 1, \dots, m$, are continuously partially differentiable
- the matrix A is uniquely determined by the so-called **Jacobi matrix**

$$J_f(\mathbf{x}) := \begin{pmatrix} \frac{\partial f_1}{\partial x_1}(\mathbf{x}) & \dots & \dots & \frac{\partial f_1}{\partial x_n}(\mathbf{x}) \\ \vdots & \ddots & \frac{\partial f_i}{\partial x_j}(\mathbf{x}) & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1}(\mathbf{x}) & \dots & \dots & \frac{\partial f_m}{\partial x_n}(\mathbf{x}) \end{pmatrix}$$

Special case $m = 1$: **transposed** gradient

We also write $D_f(\mathbf{x}) = J_f(\mathbf{x})$, indicating this is the general form of the derivative (as a matrix).

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem (Tangent plane)

Let $D \subseteq \mathbb{R}^2$, $f : D \rightarrow \mathbb{R}$ totally differentiable and $(x_0, y_0) \in D$.

Then the points (x, y, z) of the tangent plane in (x_0, y_0) are described by the equation

$$z = f(x_0, y_0) + \nabla f(x_0, y_0)^T \begin{pmatrix} x - x_0 \\ y - y_0 \end{pmatrix}$$

=

The tangent plane is locally the optimal (affine-)linear approximation for f . The plane consists out of all tangents.

In the general case the linear approximation reads


$$f(\mathbf{x}) \approx f(\hat{\mathbf{x}}) + D_f(\hat{\mathbf{x}})(\mathbf{x} - \hat{\mathbf{x}}).$$

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves


Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Tangent Plane as Linear Approximation

Analysis 2

S.-J. Kimmerle

In 2d: tangent plane



Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Tangent Plane: Example

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Chain Rule (Simple Case)

Let $\mathbf{x} : \mathbb{R} \supseteq (a, b) \rightarrow D$ differentiable in $t \in (a, b)$,
 $f : \mathbb{R}^n \supseteq D \rightarrow \mathbb{R}$ (totally) diff.able in $\mathbf{s} := \mathbf{x}(t) \in D$, D open,
then

$$f \circ \mathbf{x} : (a, b) \rightarrow \mathbb{R}, t \mapsto f(\mathbf{s}) = f(\mathbf{x}(t))$$

is totally differentiable in t and

$$\frac{d}{dt}f(\mathbf{x}(t)) = \nabla f(\mathbf{x}(t))^{\top} \dot{\mathbf{x}}(t).$$



Chain Rule (General Case)

Let $g : \mathbb{R}^p \supseteq E \rightarrow D$ (totally) diff.able in $\mathbf{t} \in E$, E open,
 $f : \mathbb{R}^n \supseteq D \rightarrow \mathbb{R}^m$ (totally) diff.able in $\mathbf{s} := g(\mathbf{t}) \in D$, D
open,

then

$$f \circ g : E \rightarrow \mathbb{R}^m, \mathbf{t} \mapsto f(\mathbf{s}) = f(g(\mathbf{t}))$$

is totally differentiable in \mathbf{t} and

$$D_{f \circ g}(\mathbf{t}) = D_f(\mathbf{x}(\mathbf{t})) D_g(\mathbf{t}).$$



Application: Coordinate Transformation I

Analysis 2

S.-J. Kimmerle

Example (Polar Coordinates in \mathbb{R}^2)

Cartesian coordinates $(x, y) \in \mathbb{R}^2$

Polar coordinates $(r, \phi) \in [0, \infty) \times (-\pi, \pi]$

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, $(x, y) \mapsto f(x, y)$ and

$F : [0, \infty) \times (-\pi, \pi], (r, \phi) \mapsto F(r, \phi) := f(r \cos(\phi), r \sin(\phi))$.

How do I transform the derivative when changing coordinates?

Let the change of coordinates be described by

$$g : \mathbb{R}^2 \rightarrow [0, \infty) \times (-\pi, \pi], \begin{pmatrix} r \\ \phi \end{pmatrix} \mapsto \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} r \cos(\phi) \\ r \sin(\phi) \end{pmatrix}.$$

By the chain rule, we compute $D_F = D_f \cdot D_g = \dots$

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n


Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Application: Coordinate Transformation II

$$D_F = D_f \cdot D_g =$$

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n


Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



The **directional derivative** $D_{\mathbf{d}}f(\mathbf{x}) = f'(\mathbf{x}; \mathbf{d})$ of a continuously differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at $\mathbf{x} \in \mathbb{R}^n$ in direction $\mathbf{d} \in \mathbb{R}^n$ is defined as

$$D_{\mathbf{d}}f(\mathbf{x}) := \lim_{t \downarrow 0} \frac{f(\mathbf{x} + t\mathbf{d}) - f(\mathbf{x})}{t}.$$

Properties:

- If $\mathbf{d} = \mathbf{e}_i$, the unit vector of coordinate i , then $D_{\mathbf{d}}f = \frac{\partial f}{\partial x_i}(\mathbf{x})$.
- If f is continuously differentiable, then $D_{\mathbf{d}}f(\mathbf{x}) = \nabla f(\mathbf{x})^\top \mathbf{d}$.
- If $D_{\mathbf{d}}f(\mathbf{x}) = \nabla f(\mathbf{x})^\top \mathbf{d} < 0$, then \mathbf{d} is a **direction of descent**, i.e.

$$f(\mathbf{x} + \alpha \mathbf{d}) < f(\mathbf{x}) \quad \text{for all } \alpha \in (0, \bar{\alpha}].$$

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Taylor expansion

- If f differentiable, then f may be approximated in a neighbourhood of $\hat{\mathbf{x}}$ by an affine-linear function as

$$f(\mathbf{x}) = f(\hat{\mathbf{x}}) + \nabla f(\hat{\mathbf{x}})^\top (\mathbf{x} - \hat{\mathbf{x}}) + R(\|\mathbf{x} - \hat{\mathbf{x}}\|),$$

where

$$\lim_{\mathbf{x} \rightarrow \hat{\mathbf{x}}} \frac{R(\|\mathbf{x} - \hat{\mathbf{x}}\|)}{\|\mathbf{x} - \hat{\mathbf{x}}\|} = 0$$

or with $t \in \mathbb{R}$, $\mathbf{d} \in \mathbb{R}^n$, $\xi_t \in (0, t)$, resp.,

$$f(\hat{\mathbf{x}} + t\mathbf{d}) = f(\hat{\mathbf{x}}) + t \nabla f(\hat{\mathbf{x}} + \xi_t \mathbf{d})^\top \mathbf{d}.$$

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Limits and Continuity in \$\mathbb{R}^n\$](#) [Differentiability in \$\mathbb{R}^n\$](#) [Optimization in \$\mathbb{R}^n\$](#) [Introduction into Linear
Regression](#)[Curves](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

- If f 2times differentiable, then f may be approximated in a neighbourhood of $\hat{\mathbf{x}}$ by a quadratic function as

$$f(\mathbf{x}) = f(\hat{\mathbf{x}}) + \nabla f(\hat{\mathbf{x}})^\top (\mathbf{x} - \hat{\mathbf{x}}) + \frac{1}{2}(\mathbf{x} - \hat{\mathbf{x}})^\top H_f(\hat{\mathbf{x}})(\mathbf{x} - \hat{\mathbf{x}}) + R(\|\mathbf{x} - \hat{\mathbf{x}}\|^2)$$

or, resp.,

$$f(\hat{\mathbf{x}} + t\mathbf{d}) = f(\hat{\mathbf{x}}) + t \nabla f(\hat{\mathbf{x}})^\top \mathbf{d} + \frac{t^2}{2} \mathbf{d}^\top H_f(\hat{\mathbf{x}} + \xi_t \mathbf{d}) \mathbf{d}.$$

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Taylor Expansion - Example

Approximate around $(\hat{x}_1, \hat{x}_2) = (0, 0)$ the function

$$f(x_1, x_2) = \exp(x_1) \ln(1 + x_2).$$

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Instead of the (usually unknown) real values

$$\hat{\mathbf{x}} = (\hat{x}_1, \dots, \hat{x}_n)^\top$$

we measure approximate values

$$\mathbf{x} = (x_1, \dots, x_n)^\top.$$

The measurement errors (uncertainty, observational deviations)

$$\Delta x_i := x_i - \hat{x}_i, 1, \dots, n,$$

yield an error for the value of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$

$$\Delta f(\mathbf{x}) := f(\mathbf{x}) - f(\hat{\mathbf{x}}) \approx \nabla f(\hat{\mathbf{x}})^\top (\mathbf{x} - \hat{\mathbf{x}})$$

or in absolute values

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Suppose

$$|\Delta f(\mathbf{x})| \leq \sum_{i=1}^n \left| \frac{\partial f}{\partial x_i}(\hat{\mathbf{x}}) \right| |\Delta x_i|.$$

If the “scattering” of errors is known as $|x_i| \leq S_i$, then the absolute maximal error S reads

$$|\Delta f(\mathbf{x})| \leq S := \sum_{i=1}^n \left| \frac{\partial f}{\partial x_i}(\hat{\mathbf{x}}) \right| S_i.$$

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Limits and Continuity in \$\mathbb{R}^n\$](#) [Differentiability in \$\mathbb{R}^n\$](#) [Optimization in \$\mathbb{R}^n\$](#) [Introduction into Linear
Regression](#)[Curves](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Propagation of Uncertainty - Example

Given: measurement of 2 time intervals

$$x = (20 \pm 0.1)[ms] \text{ and } y = (15 \pm 0.1)[ms]$$

Searched for: maximal error of time difference $x - y$

Solution: Set $f : \mathbb{R}^2 \rightarrow \mathbb{R}, (x, y) \mapsto f(x, y) = x - y$.

$$\nabla f(x, y) = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

Absolute maximal error:

$$S = \left| \frac{\partial f}{\partial x}(20, 15) \right| \cdot 0.1 + \left| \frac{\partial f}{\partial y}(20, 15) \right| \cdot 0.1 = 0.2 [ms]$$

Relative maximal error:

$$\frac{S}{|f(20, 15)|} = \frac{0.2}{5} = 4\%.$$

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Limits and Continuity in \$\mathbb{R}^n\$](#) [Differentiability in \$\mathbb{R}^n\$](#) [Optimization in \$\mathbb{R}^n\$](#) [Introduction into Linear
Regression](#)[Curves](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Sum or difference:

Absolute maximal error is sum of maximal errors of inputs

Product or quotient:

Relative maximal error is sum of relative errors of inputs

Remark:

More realistic for an error estimation is a stochastic error analysis (Gaussian propagation of uncertainty)

↪ stochastics lecture

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Optimization in Higher Dimensions

Analysis 2

S.-J. Kimmerle

As an application we consider again the problem of minimizing a given function

$$f : \mathbb{R}^n \rightarrow \mathbb{R}$$

but now with n variables $\mathbf{x} = (x_1, x_2, \dots, x_n)^\top \in \mathbb{R}^n$.

For simplicity, we consider the case without restrictions (constraints) on \mathbf{x} .

The (local) conditions derived in this subsection hold in the interior of a domain, but not on boundaries.

Note that maximization problems for $g : \mathbb{R}^n \rightarrow \mathbb{R}$ are equivalent to minimization problems where $f = -g$.

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Unrestricted Optimization for any $n \in \mathbb{N}$

Analysis 2

S.-J. Kimmerle

We consider

Problem (Unrestricted optimization problem (UOP))

Minimize $f(x)$ subject to the constraint $x \in \mathbb{R}^n$.

*Thereby let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ an **at least 1x (better 2x) differentiable** function.*

Idea:

Search at first for local minima (from which one obtains global extrema under certain conditions).

Aims:

Necessary conditions

Sufficient conditions

Wish: necessary & sufficient conditions

Remark:

Numerical methods are mostly based on necessary conditions.

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Matrices do not have signs, but:

Definition (Definiteness of a Matrix)

Let $H \in \mathbb{R}^{n \times n}$ and $v \in \mathbb{R}^n \setminus \mathbf{0}$.

If $v^T H v \begin{cases} > \\ < \\ \geq \\ \leq \end{cases} 0$ for all v , then H is $\begin{cases} \text{positive definite} \\ \text{negative definite} \\ \text{positive semi-definite} \\ \text{negative semi-definite} \end{cases}$.

If $v^T H v > 0$ for some v and $v^T H v < 0$ for another v (i.e. H is neither positive semi-definite nor negative semi-definite), then H is **indefinite**.

If $v^T H v = 0$ for some v , then H is **singular**.

For functions in several variables, the Hessian plays the role of the 2nd derivative.

The definiteness of the Hessian generalizes the sign of the 2nd derivative of a function in 1 variable.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Limits and Continuity in \$\mathbb{R}^n\$](#) [Differentiability in \$\mathbb{R}^n\$](#) [Optimization in \$\mathbb{R}^n\$](#) [Introduction into Linear
Regression](#)[Curves](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Revision/Outlook: Definite Matrices - Examples

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Necessary Condition of 1st Order

Theorem (Necessary Condition of 1st Order)

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable and $\hat{\mathbf{x}} \in \mathbb{R}^n$ a local minimum of f .

Then there holds

$$\nabla f(\hat{\mathbf{x}}) = \mathbf{0}.$$

Definition (Stationary Point)

Any point $\mathbf{x} \in \mathbb{R}^n$ with $\nabla f(\mathbf{x}) = \mathbf{0}$ is called a **stationary point** of $f : \mathbb{R}^n \rightarrow \mathbb{R}$.

Stationary points are not automatically minima, but candidates for (local) minima!

Most numerical methods try to approximate stationary points.

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Theorem (Necessary Condition of 2nd Order)

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be 2x continuously differentiable and $\hat{\mathbf{x}} \in \mathbb{R}^n$ a local minimum of f .

Then the Hessian

$$H_f(\hat{\mathbf{x}})$$

is positive semi-definite.

In case $n = 1$ we have

$$f''(\hat{x}) \geq 0.$$

Likewise, this condition only yields potential candidates for a (local) minimum.

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem (Sufficient Condition of 2nd Order)

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be 2x continuously differentiable and $\hat{\mathbf{x}} \in \mathbb{R}^n$ a stationary point of f with positive definite Hessian.

Then $\hat{\mathbf{x}}$ is a strict local minimum of f .

In case $n = 1$ we assume that

$$f''(\hat{x}) > 0.$$

This allows us to decide whether a candidate $\hat{\mathbf{x}}$ that fulfills necessary conditions (1st or 2nd order) is indeed a local minimum.

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

- The **gradient** of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at a point $\mathbf{x} = (x_1, \dots, x_n)^\top \in \mathbb{R}^n$ is defined as a column vector

$$\nabla f(x_1, \dots, x_n) := \begin{pmatrix} \frac{\partial f}{\partial x_1}(x_1, \dots, x_n) \\ \vdots \\ \frac{\partial f}{\partial x_n}(x_1, \dots, x_n) \end{pmatrix}.$$

In case $n = 1$ the gradient is the 1st derivative of f at point x , shortly $f'(x)$.

Illustratively, the gradient describes the steepest ascent.

The gradient is orthogonal on level curves.



- The **Hessian** of a function f at a point \mathbf{x} is defined as the quadratic matrix

$$H_f(x_1, \dots, x_n) = H_f(\mathbf{x}) := \begin{pmatrix} \frac{\partial^2 f}{\partial x_1 \partial x_1}(\mathbf{x}) & \dots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(\mathbf{x}) \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(\mathbf{x}) & \dots & \frac{\partial^2 f}{\partial x_n \partial x_n}(\mathbf{x}) \end{pmatrix}.$$

In case $n = 1$ the Hessian ist the 2nd derivative of f at a point x , shortly $f''(x)$.

Illustratively, the Hessian describes the local curvature of a function.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Limits and Continuity in \$\mathbb{R}^n\$](#) [Differentiability in \$\mathbb{R}^n\$](#) [Optimization in \$\mathbb{R}^n\$](#) [Introduction into Linear
Regression](#)[Curves](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Hessian in 2d - Example 1

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n


Introduction into Linear
Regression

Curves


Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



$$f(x_1, x_2) = x_1^2 + x_2^2, \quad H_f(\hat{x}_1, \hat{x}_2) = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$$



Hessian in 2d - Example 2

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n


Introduction into Linear
Regression

Curves


Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



$$f(x_1, x_2) = -x_1^2 - x_2^2, \quad H_f(\hat{x}_1, \hat{x}_2) = \begin{pmatrix} -2 & 0 \\ 0 & -2 \end{pmatrix}$$



Hessian in 2d - Example 3

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n


Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



$$f(x_1, x_2) = x_1^2 - x_2^2, \quad H_f(\hat{x}_1, \hat{x}_2) = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix}$$



Hessian in 2d - Examples 4 & 5

Other possibilities for definite matrices in $\mathbb{R}^{2 \times 2}$:

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Points $\hat{\mathbf{x}}$ with $\nabla f(\hat{\mathbf{x}}) = (0, 0, \dots, 0)^\top$ are called **stationary points**.

Depending on the properties of the Hessian we find:

- **positive definite:** local minimum
- **negative definite:** local maximum
- **indefinite:** saddle point
- **singular:** everything is possible (e.g. for positive/negative semi-definite Hessian)

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Optimization in \mathbb{R}^n - Example

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x, y) = \exp(x) \cdot (2x + y^2)$$

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Introduction into Linear Regression

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review


- Theory and experiment:
How to confirm the congruence?

- Example:
Elongation y [cm] of a spring
subject to a force t [N]

Hooke's law:

$$y = y_0 + \frac{1}{D}t$$

(D spring constant,
describes stiffness of spring,
 y_0 pre-elongation of spring
due to own weight)



Source:
[Wikipedia](#)

- Are the free parameters D , y_0 constant over different experiments?



Linear Regression - Intro

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus


Summary -
Outlook and
Review

Theory: $y(t) = x_1 + x_2 t$

Data: (t_i, y_i)

Error (residuum): $r_i = y_i - y(t_i)$

i	t_i	y_i	$y(t_i)$	r_i	r_i^2
1	1	2			
2	2	3			
3	3	7			
4	4	8			
5	5	9			



Linear Regression - 1st Try

Analysis 2


S.-J. Kimmerle

Theory: $y(t) = 1 + t$

Data: (t_i, y_i)

Error (residuum): $r_i = y_i - y(t_i) = y_i - 1 - t_i$

i	t_i	y_i	$y(t_i)$	r_i	r_i^2
1	1	2	2	0	0
2	2	3	3	0	0
3	3	7	4	3	9
4	4	8	5	3	9
5	5	9	6	3	9
					$\sum_{i=1}^5 r_i^2 = 27$



Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review




Linear Regression - 2nd Try

Theory: $y(t) = 4 + t$

Data: (t_i, y_i)

Error (residuum): $r_i = y_i - y(t_i) = y_i - 4 - t_i$

i	t_i	y_i	$y(t_i)$	r_i	r_i^2
1	1	2	5	-3	9
2	2	3	6	-3	9
3	3	7	7	0	0
4	4	8	8	0	0
5	5	9	9	0	0
					$\sum_{i=1}^5 r_i^2 = 18$




Linear Regression - 3rd Try

Theory: $y(t) = 2.5 + t$

Data: (t_i, y_i)

Error (residuum): $r_i = y_i - y(t_i) = y_i - 2.5 - t_i$

i	t_i	y_i	$y(t_i)$	r_i	r_i^2
1	1	2	3.5	-1.5	2.25
2	2	3	4.5	-1.5	2.25
3	3	7	5.5	1.5	2.25
4	4	8	6.5	1.5	2.25
5	5	9	7.5	1.5	2.25
				$\sum_{i=1}^5 r_i^2 = 11.25$	



Linear Regression - Solved

Analysis 2


S.-J. Kimmerle

Theory: $y(t) = 0.1 + 1.9t$

Data: (t_i, y_i)

Error (residuum): $r_i = y_i - y(t_i) = y_i - 0.1 - 1.9t_i$

i	t_i	y_i	$y(t_i)$	r_i	r_i^2
1	1	2	2	0	0
2	2	3	3.9	-0.9	0.81
3	3	7	5.8	1.2	1.44
4	4	8	7.7	0.3	0.09
5	5	9	9.6	-0.6	0.36
					$\sum_{i=1}^5 r_i^2 = 2.7$



Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Linear Regression: Normal Equations

Analysis 2

S.-J. Kimmerle

In general (m arbitrary):

$$A^T A = \begin{pmatrix} m & \sum_{i=1}^m t_i \\ \sum_{i=1}^m t_i & \sum_{i=1}^m t_i^2 \end{pmatrix}, \quad A^T \mathbf{y} = \begin{pmatrix} \sum_{i=1}^m y_i \\ \sum_{i=1}^m t_i y_i \end{pmatrix}$$

$$x_1 = \frac{\sum_{i=1}^m t_i^2 \sum_{j=1}^m y_j - \sum_{i=1}^m t_i \sum_{j=1}^m t_j y_j}{m \sum_{i=1}^m t_i^2 - \left(\sum_{i=1}^m t_i \right)^2}$$

$$x_2 = \frac{m \sum_{i=1}^m t_i y_i - \sum_{i=1}^m t_i \sum_{j=1}^m y_j}{m \sum_{i=1}^m t_i^2 - \left(\sum_{i=1}^m t_i \right)^2}$$

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



We conclude:

- In general more data points y_i (here 5) as parameters x_i (here 2)
- Measurement of process/data collection afflicted with uncertainties (w/o systematical errors)
- Overdetermined LES \rightsquigarrow in general no solution exists

Aims:

- Determine $\mathbf{x} = (x_1, \dots, x_n)^T$ “optimally” from data
- General method to do that

We call this **regression** (in particular in stochastics) or **curve fitting**.

Special case of a mathematical optimization method

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Linear Regression: Stochastic Uncertainties

Analysis 2

S.-J. Kimmerle

Problem (Linear Regression)

Assume: linear relation (e.g. from physics)

$$y(t) = x_1 + x_2 t$$

Given: data points (t_i, y_i) , $i = 1, \dots, m$, afflicted with uncertainties (errors) ε_i .
The uncertainties are random variables and 0 in average.

Searched for: $x_1, x_2 \in \mathbb{R}$, such that

$$\begin{pmatrix} 1 & t_1 \\ \vdots & \vdots \\ 1 & t_i \\ \vdots & \vdots \\ 1 & t_m \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} y_1 \\ \vdots \\ y_i \\ \vdots \\ y_m \end{pmatrix} + \begin{pmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_i \\ \vdots \\ \varepsilon_m \end{pmatrix}$$

We write

$$A\mathbf{x} = \mathbf{y} + \boldsymbol{\varepsilon}$$

with $A \in \mathbb{R}^{m \times 2}$, $\mathbf{x} \in \mathbb{R}^2$, $\mathbf{y}, \boldsymbol{\varepsilon} \in \mathbb{R}^m$.

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Minimization Problem

General situation: $A \in \mathbb{R}^{m \times n}$, $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{y} \in \mathbb{R}^m$

Idea: determine $\mathbf{x} = (x_1, \dots, x_n)^\top$, such that the error in the LES

$$\|\mathbf{Ax} - \mathbf{y}\|_2 = \sqrt{\sum_{i=1}^m \left(\sum_{j=1}^n A_{ij}x_j - y_i \right)^2} \quad \text{bzw.} \quad \frac{1}{2} \|\mathbf{Ax} - \mathbf{y}\|_2^2$$

is minimized.

The minimum is denoted by $\hat{\mathbf{x}}$.

This minimization problem is called a **linear regression problem** or **least-squares problem**.

Method of least squares, better **method of least residual sum of squares**

By **C. F. Gauß** and **A.-M. Legendre** (1805)

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Problem (★) (Lin. regression as minimization problem)

Let be given a matrix $A \in \mathbb{R}^{m \times n}$ and a vector $\mathbf{y} \in \mathbb{R}^m$ with $m, n \in \mathbb{N}$.

We search for the solution $\hat{\mathbf{x}} \in \mathbb{R}^n$ of the minimization problem

$$\frac{1}{2} \|A\hat{\mathbf{x}} - \mathbf{y}\|_2^2 = \min_{\mathbf{x} \in \mathbb{R}^n} \frac{1}{2} \|A\mathbf{x} - \mathbf{y}\|_2^2.$$

- If $m = n$ and A invertible, then $A\mathbf{x} = \mathbf{y}$ has a unique solution.
- The case $m > n$ is of uttermost importance in applications.
- In the case $m < n$ the LES $A\mathbf{x} = \mathbf{y}$ is underdetermined. It may be unsolvable, if $\text{rank}(A) \neq \text{rank}(A \mid \mathbf{y})$.

Moreover, other norms as $\|\cdot\|_2$ could be considered. Then the determination of solutions is harder in general, since differentiability might not be given.

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Theorem (Gaussian normal equations)

$\hat{\mathbf{x}}$ solves Problem (\star) if and only iff the **normal equations**

$$A^T A \hat{\mathbf{x}} = A^T \mathbf{y}.$$

hold true.

$A^T A \in \mathbb{R}^{n \times n}$ is symmetric and positive semi-definite.

Theorem (Uniqueness)

Let $m \geq n$. Suppose $A \in \mathbb{R}^{m \times n}$ has maximal rank,
d.h. $\text{rank}(A) = n$.

Exactly then the minimization problem (\star) or the normal equations, resp., are uniquely solvable.

Then $A^T A \in \mathbb{R}^{n \times n}$ is invertible and positive definite.

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

(Non-)Linear Regression: TTF under Temperature Stress




Figure: Curve fitting over different temperatures T compared with measured data points (squares) with confidence intervals for $q = 90\%$ (triangles). Left-hand side for $\mu_{krit}(T) \pm \sigma_{krit}(T)$, in red the fitted curve, right-hand side $\sigma_{krit}(T)$. [K., Dvorsky, Ließ, Avenhaus 2019]

Model based approach

$$f_{krit}(T) = t_{\Theta} + t_0 \exp\left(\left(\frac{T_a}{T - T_{\infty}}\right)^d\right)$$



Lin. Regression: Transformation

Analysis 2

S.-J. Kimmerle

Example (Measurement of TTF (Time-To-Failure) of Electrical Automotive Components)

T [°C]	115	130	140	155
μ_{krit} [h]	3791.62	987.74	439.66	189.94

Conjecture (model based): Arrhenius law

$$\mu_{krit}(T) = t_0 \exp\left(\frac{T_a}{T - T_\infty}\right)$$

(μ_{krit} TTF in h, t_0 reaction-kinetic period in h, T temperature in K, T_a activation temperature in K, $T_\infty = 173.15$ consolidation temperature in K)

We may transform this into an affine-linear relation

$$\ln(\mu_{krit}(T)) = \ln(t_0) + \frac{T_a}{T - T_\infty} \quad \Longleftrightarrow \quad y(t) = x_1 + x_2 t$$

By insertion of measured data we obtain a linear equation system (LES)

$$y_i = x_1 + x_2 t_i, \quad i = 1, \dots, 4.$$

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Linear Regression: Example - Result

Analysis 2

S.-J. Kimmerle





Figure: Curve fitting over different inverse temperature differences t , in semi-logarithmic representation and compared with measured data points (squares) together with confidence intervals for $q = 90\%$ (triangles). Left-hand side for $t_{krit}(1/t) \pm \sigma_{krit}(1/t)$, in red the fitted curve. Right-hand side $\sigma_{krit}(T)$. [K., Dvorsky, Ließ, Avenhaus 2019]

Introduction

Power series

Differentiation in Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear Regression

Curves

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



Linear Regression: Example - Re-transformed Result




Figure: Curve fitting over different temperatures T , re-transformed in exponential representation and compared with measured data points (squares) together with confidence intervals for $q = 90\%$ (triangles). Left-hand side for $t_{krit}(T) \pm \sigma_{krit}(T)$, in red the fitted curve. Right-hand side $\sigma_{krit}(T)$. [K., Dvorsky, Ließ, Avenhaus 2019]



A set $D \subseteq \mathbb{R}^n$ is called **(path-)connected**, if for any $\mathbf{p}, \mathbf{q} \in D$ there exists an interval $[a, b]$ and a continuous function $r : [a, b] \rightarrow \mathbb{R}^n$, such that:

- $r(t) \in D$ for all $t \in [a, b]$
- $r(a) = \mathbf{p}$ and $r(b) = \mathbf{q}$

The set $r([a, b])$ (the range of r) is called a **curve**.

The mapping r itself is called **parametrization of the curve** or **path** from \mathbf{p} to \mathbf{q} in D .

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

A curve is called **closed**, if $r(a) = r(b)$.

A set D is called **simply connected**, if any closed curve maybe continuously contracted (however this is defined rigorously) to a single point.

A **region** or domain¹ is an open and connected subset of \mathbb{R}^n .

¹not to be confused with a domain of definition

Introduction

Power series

Differentiation in
Higher DimensionsLimits and Continuity in \mathbb{R}^n Differentiability in \mathbb{R}^n Optimization in \mathbb{R}^n Introduction into Linear
Regression

Curves

Integration in
Higher DimensionsFurther Topics in
CalculusSummary -
Outlook and
Review

Examples for Connected Sets

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



We start with 2d:

Consider a parametrization $\mathbf{r}(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$

with x, y differentiable functions.

We call

$$\dot{\mathbf{r}}(t) = \lim_{h \rightarrow 0} \frac{1}{h} (\mathbf{r}(t+h) - \mathbf{r}(t)) = \begin{pmatrix} \dot{x}(t) \\ \dot{y}(t) \end{pmatrix}$$

a tangential vector of the curve.

It may interpreted as the limit vector of the secant vectors.

Analogously this may be generalized to 3d and higher dimensions.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Limits and Continuity in \$\mathbb{R}^n\$](#) [Differentiability in \$\mathbb{R}^n\$](#) [Optimization in \$\mathbb{R}^n\$](#) [Introduction into Linear
Regression](#)[Curves](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Differentiation of Curves - Example

Motion of a mass point in 3d space

Position

Velocity

Acceleration

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Limits and Continuity in \mathbb{R}^n

Differentiability in \mathbb{R}^n

Optimization in \mathbb{R}^n

Introduction into Linear
Regression

Curves

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review



Introduction

Power series

Differentiation in
Higher Dimensions

**Integration in
Higher Dimensions**

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review

- 1 Introduction
- 2 Power series
- 3 Differentiation in Higher Dimensions
- 4 Integration in Higher Dimensions**
 - Curve Integrals and Solids of Revolution
 - Integration of Functions with Several Variables
 - Transformation Formula
- 5 Further Topics in Calculus
- 6 Summary - Outlook and Review



Curve Integrals, Areas of Sectors & Solids of Revolution

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review

Before turning to iterated integrals of functions of more than 1 variable,
we consider integrals of functions in 1 variable:

- Vector-valued functions:
Curve integrals
- Real-valued functions of vectors, but with symmetry:
Areas of sectors,
Solids of revolution



Regular Parametrization of a Curve

Analysis 2

S.-J. Kimmerle

Let $D \subseteq \mathbb{R}^n$ open.

A mapping $\mathbf{r} : [a, b] \rightarrow D : t \mapsto \mathbf{r}(t) = (r_1(t), \dots, r_n(t))^T$ is called **regular parametrization** of a curve in D , if

- \mathbf{r} is continuously differentiable and
- $\mathbf{r}'(t) \neq \mathbf{0} \in \mathbb{R}^n$ for all $t \in [a, b]$.

Different regular parametrizations exist.

The sense of circulation may be different.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Let $D \subseteq \mathbb{R}^n$ open.

For any continuous function $\mathbf{f} : D \rightarrow \mathbb{R}^n$ and any regular parametrization $\mathbf{r} : [a, b] \rightarrow D$ of a curve $K := \mathbf{r}([a, b])$ we call

$$\int_K \mathbf{f}(\mathbf{x}) \bullet d\mathbf{x} := \int_a^b \mathbf{f}(\mathbf{r}(t)) \bullet \mathbf{r}'(t) dt$$

the **curve integral** of \mathbf{f} along the curve K w.r.t. \mathbf{r} .

The curve integral is independent from the parametrization, but its sign depends on the sense of circulation.

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

Arc Length - Parametrized Curve

Analysis 2

S.-J. Kimmerle

Let K be a curve with a regular parametrization $(x(t), y(t))^T$, $t \in [a, b]$,
i.e. with $x(t)$, $y(t)$ cont. diff.able on $[a, b]$ and

$$(\dot{x}(t))^2 + (\dot{y}(t))^2 \neq 0 \quad \text{for all } t \in [a, b].$$

Then the arc length is given by

$$L = \int_a^b \sqrt{\dot{x}(t)^2 + \dot{y}(t)^2} dt.$$

In higher dimensions with $\mathbf{x}(t) \in \mathbb{R}^n$ there holds under analogous assumptions:

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Arc length in 2d - Cartesian Coordinates

Analysis 2

S.-J. Kimmerle

For the graph $y = f(\mathbf{x})$ of a cont. differentiable function
 $f : [a, b] \rightarrow \mathbb{R}$
the arc length is given by

$$L = \int_a^b \sqrt{1 + (y'(x))^2} dx.$$

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions


Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Arc length in 2d - Polar Coordinates

Analysis 2

S.-J. Kimmerle

For a curve with a regular parametrization $r(\phi)$, $\phi \in [\alpha, \beta]$,
i.e. with $r(\phi)$ cont. differentiable on $[\alpha, \beta]$ and

$$r'(\phi) \neq 0 \quad \text{for all } \phi \in [\alpha, \beta].$$

$$L = \int_{\alpha}^{\beta} \sqrt{r(\phi)^2 + (\dot{r}(\phi))^2} d\phi.$$

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Areas of Sectors in Polar Coordinates

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review

Consider $r : [\alpha, \beta] \rightarrow \mathbb{R}_0^+$ continuous, where $\alpha < \beta$.
The sector enclosed by the 3 curves $r(\phi)$, $\phi = \alpha$, and $\phi = \beta$ has the area

$$A = \frac{1}{2} \int_{\alpha}^{\beta} (r(\phi))^2 d\phi.$$

Motivation:

$$\Delta A \approx \pi r^2 \cdot \frac{\Delta \phi}{2\pi}$$
$$A = \lim_{\Delta \phi \rightarrow 0} \sum \frac{1}{2} r^2 \Delta \phi$$



Areas of Sectors in Polar Coordinates - Example

Archimedean spiral (1 turn)

$$r(\phi) = a\phi, \quad a > 0, \quad \text{for } 0 \leq \phi \leq 2\pi$$

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Leibniz sector formula

$$A = \frac{1}{2} \left| \int_{t_1}^{t_2} (x(t)y'(t) - y(t)x'(t)) dt \right|$$

for a parametrized curve $K = \{(x(t), y(t)), t_1 \leq t \leq t_2\}$,
where $x(t)$, $y(t)$ cont. differentiable

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Let $f : [a, b] \rightarrow \mathbb{R}$ continuous.

The solid of revolution generated by rotating the curve $y = f(x)$, $a \leq x \leq b$, around the x -axis has the volume

$$V = \pi \int_a^b (f(x))^2 dx$$

Motivation:

$$\Delta V = A(x) \cdot \Delta x \approx \pi(f(x))^2 \cdot \Delta x$$

$$V = \lim_{\Delta x \rightarrow 0} \sum \pi(f(x))^2 \cdot \Delta x$$

Remark: Rotations around other axes yield analogous formulas.

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

Solids of Revolution - Example

Circular cone

$$f(x) = \frac{r}{h}x$$

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions


Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Up to now we have considered different cases of functions depending in principle on one variable, although the range might be higher dimensional, i.e. in \mathbb{R}^n . We wish to enlarge this to several variables in \mathbb{R}^m .

We consider $f : B \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$.

$$S_n := \sum_{i=1}^n f(x_*^{[i]}, y_*^{[i]}) \cdot \Delta F_i \quad \text{with } (x_*^{[i]}, y_*^{[i]}) \in B$$


Small domains B_i with area $|B_i| = \Delta F_i$ with fineness δ

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

Idea: iterated Riemann integral

But why should we restrict us to an approximation by rectangles?

Like Riemann construction

$$\iint_B f(x, y) dF = \lim_{\delta \rightarrow 0, n \rightarrow \infty} \sum_{i=1}^n f(x_*^{[i]}, y_*^{[i]}) \cdot \Delta F_i$$

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

Area integrals

$$A = \iint_B 1 \, dA = \text{area of } B$$

Volume integrals

$$V = \iiint_C 1 \, dV = \text{volume of } C$$

$$\stackrel{\text{special case}}{=} \iint_B f(x, y) \, dA$$

= volume of solid with base area B
and top area described by $f(B)$

Note $A = |B|$, $V = |C|$

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

For a more formal approach, we need some preparation:

- Recall: A set $D \subseteq \mathbb{R}^n$ is called connected, iff for any 2 points $\mathbf{x}^{[0]}$ and $\mathbf{x}^{[1]}$ there exists a curve $k : [a, b] \rightarrow \mathbb{R}^n$ with $k(a) = \mathbf{x}^{[0]}$ and $k(b) = \mathbf{x}^{[1]}$.

A **region** or domain² is an open and connected subset of \mathbb{R}^n .

- Normal areas (regular domains) \rightsquigarrow see next slide

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

²not to be confused with a domain of definition

Normal areas are in 2D:

- Type I:

$$B_I = \{(x, y) \mid a \leq x \leq b \text{ and } g(x) \leq y \leq h(x)\}$$

with $a, b \in \mathbb{R}$, where $a < b$, and

$g : [a, b] \rightarrow \mathbb{R}$, $h : [a, b] \rightarrow \mathbb{R}$ cont. differentiable.

- Type II:

$$B_{II} = \{(x, y) \mid c \leq y \leq d \text{ and } G(y) \leq x \leq H(y)\}$$

with $c, d \in \mathbb{R}$, where $c < d$, and

$G : [c, d] \rightarrow \mathbb{R}$, $H : [c, d] \rightarrow \mathbb{R}$ cont. differentiable.

The roles of x and y are reversed.

This may be extended to higher dimensions.

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

- We perform the integrations "from inside to outside". The integral and the differential work like a "bracket".
- For practical computations it may be helpful to split the normal area B by cuts (that are parallel to the axes) into smaller (disjunct) normal areas, e.g. B_1 and B_2 . The whole integral is then obtained by the additivity.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Fubini Theorem

The order of integration may not be exchanged in general, but:

Theorem (Fubini Theorem)

Let $R = \{(x, y) \mid a \leq x \leq b \text{ and } c \leq y \leq d\}$, $a, b, c, d \in \mathbb{R}$ a rectangle,
and $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ continuous.

Then there holds

$$\iint_R f \, dA = \int_a^b \int_c^d f(x, y) \, dy \, dx = \int_c^d \int_a^b f(x, y) \, dx \, dy.$$



Fubini Theorem - Example

Consider B bounded by the curves $y = x$, $y = \frac{1}{x}$, and $y = 2$.

$$A = \iint_B dx \, dy = \iint_B dy \, dx$$

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Let $f : B \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$,
 B a normal area.

$$\begin{aligned}\int_B f(\mathbf{x}) \, dF &:= \int \cdots \int_B f(\mathbf{x}) \, d\mathbf{x} \\ &:= \int \cdots \int_B f(\mathbf{x}) \, dx_1 \dots dx_n \\ &= \lim_{\delta \rightarrow 0, n \rightarrow \infty} \sum_{i=1}^n f(\mathbf{x}^{[i]}) \cdot \Delta F_i\end{aligned}$$

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

The following properties transfer to multiple integrals.

Let $f, g : B \rightarrow \mathbb{R}$ integrable functions and $B \subseteq \mathbb{R}^n$ a normal area.

Linearity:

$$\int_B f(\mathbf{x}) + g(\mathbf{x}) d\mathbf{x} = \int_B f(\mathbf{x}) d\mathbf{x} + \int_B g(\mathbf{x}) d\mathbf{x}$$

and (factor rule)
$$\int_B c f(\mathbf{x}) d\mathbf{x} = c \int_B f(\mathbf{x}) d\mathbf{x}$$

Additivity for $B_1 \cup B_2 = B$ with $B_1 \cap B_2 = \emptyset$:

$$\int_{B_1} f(\mathbf{x}) d\mathbf{x} + \int_{B_2} f(\mathbf{x}) d\mathbf{x} = \int_B f(\mathbf{x}) d\mathbf{x}$$

Monotonicity:
$$f \leq g \implies \int_B f(\mathbf{x}) d\mathbf{x} \leq \int_B g(\mathbf{x}) d\mathbf{x}$$

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Inequalities:

“triangle inequality”

$$\left| \int_B f(\mathbf{x}) \, d\mathbf{x} \right| \leq \int_B |f(\mathbf{x})| \, d\mathbf{x}$$

Cauchy-Schwarz

$$\left(\int_B f(\mathbf{x})g(\mathbf{x}) \, d\mathbf{x} \right)^2 \leq \left(\int_B f(\mathbf{x})^2 \, d\mathbf{x} \right) \left(\int_B g(\mathbf{x})^2 \, d\mathbf{x} \right)$$

Integration over a set N of measure “zero”:

$$\int_N f(\mathbf{x}) \, d\mathbf{x} = 0$$

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Sets of Measure Zero

We only give some examples in 2d:

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions


Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Another Application: Barycenter

The **center of mass** \mathbf{s} of a mass distribution (mass density) $\rho : B \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^+$ in space is the unique point where the weighted relative position of the distributed mass sums to zero:

$$\mathbf{s} := \frac{1}{M} \int_B \mathbf{x} \rho(\mathbf{x}) d\mathbf{x}$$

where $M := \int_B \rho(\mathbf{x}) d\mathbf{x}$.

Remarks:

A center of mass may be translated to any distribution ρ or a data set.

If $\rho = 1$, we obtain the so-called centroid (geometric center). The word **barycenter** comprises the terms “center of mass” and “centroid”.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Curve Integrals and Solids
of Revolution](#)[Integration of Functions with
Several Variables](#)[Transformation Formula](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Example: Barycenter of

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Area of a Surface in Space

If $f : B \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ continuously partial differentiable,
 $B \subseteq \mathbb{R}^2$ regular,

then the surface (the graph of f) has the area

$$\int \int_B \sqrt{1 + \left(\frac{\partial f}{\partial x}(x, y)\right)^2 + \left(\frac{\partial f}{\partial y}(x, y)\right)^2} dx dy.$$

Remarks:

- For a $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ this reduces to the length of a curve.
- Surface integrals are a topic of their own.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Curve Integrals and Solids
of Revolution](#)[Integration of Functions with
Several Variables](#)[Transformation Formula](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Transformation Formula: Coordinate Transformation

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review

We consider a domain $S \subseteq \mathbb{R}^2$ generated by a **coordinate transformation**

$$x = x(u, v), y = y(u, v)$$

(being continuously differentiable, bijective, ... see literature) from a domain $B \subseteq \mathbb{R}^2$.

Let

$$g : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \begin{pmatrix} u \\ v \end{pmatrix} \mapsto \begin{pmatrix} x(u, v) \\ y(u, v) \end{pmatrix}.$$

Analogously, we define coordinate transformations for 3d and higher dimensions.



Transformation Formula 2d

Analysis 2

S.-J. Kimmerle

For such a coordinate transformation $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$,
we have for $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ continuous

$$\begin{aligned} & \iint_S f(x, y) \, dx \, dy \\ &= \iint_B f(x(u, v), y(u, v)) \cdot |\det(J_g(u, v))| \, du \, dv. \end{aligned}$$

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Transformation from Polar Coordinates

For polar coordinates there holds

$$x = r \cos(\phi),$$

$$y = r \sin(\phi),$$


thus

$$g : \mathbb{R}^+ \times [0, 2\pi) \rightarrow \mathbb{R}^2, \begin{pmatrix} r \\ \phi \end{pmatrix} \mapsto \begin{pmatrix} x(r, \phi) \\ y(r, \phi) \end{pmatrix} = \begin{pmatrix} r \cos(\phi) \\ r \sin(\phi) \end{pmatrix}.$$

Hence the Jacobian determinant (functional determinant) is:

$$\det \begin{pmatrix} \cos(\phi) & -r \sin(\phi) \\ \sin(\phi) & r \cos(\phi) \end{pmatrix} = r \cos(\phi)^2 + r \sin(\phi)^2 = r$$

Since $r > 0$, we replace the “area element” $dx dy$ by $r dr d\phi$.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Curve Integrals and Solids
of Revolution](#)[Integration of Functions with
Several Variables](#)[Transformation Formula](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Example: Polar Coordinates

Let $S = \{x^2 + y^2 \leq 1\}$.

We apply the transformation formula to compute

$$\iint_S \rho x^2 dx dy$$

(the moment of inertia of the full circular disc w.r.t. the y -axis).

We assume $\rho = \text{const.}$



Transformation Formula 3d

For a suitable (see 2d case) coordinate transformation

$$g : \mathbb{R}^3 \rightarrow \mathbb{R}^3,$$

we have for $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$ continuous

$$\begin{aligned} & \iiint_S f(x, y, z) \, dx \, dy \, dz \\ &= \iiint_B f(x(u, v, w), y(u, v, w)) \cdot |\det(J_g(u, v, w))| \, du \, dv \, dw. \end{aligned}$$

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Curve Integrals and Solids
of Revolution](#)[Integration of Functions with
Several Variables](#)[Transformation Formula](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Transformation from Cylinder Coordinates

Analysis 2

S.-J. Kimmerle

For cylinder coordinates there holds

$$x = r \cos(\phi),$$

$$y = r \sin(\phi),$$

$$z = z$$

thus

$$g : \mathbb{R}^+ \times (-\pi, \pi] \times \mathbb{R} \rightarrow \mathbb{R}^3, \begin{pmatrix} r \\ \phi \\ z \end{pmatrix} \mapsto \begin{pmatrix} x(r, \phi, z) \\ y(r, \phi, z) \\ z(r, \phi, z) \end{pmatrix} = \begin{pmatrix} r \cos(\phi) \\ r \sin(\phi) \\ z \end{pmatrix}.$$

Hence the Jacobian determinant (functional determinant) is:

$$\det \begin{pmatrix} \cos(\phi) & -r \sin(\phi) & 0 \\ \sin(\phi) & r \cos(\phi) & 0 \\ 0 & 0 & 1 \end{pmatrix} = (r \cos(\phi)^2 + r \sin(\phi)^2) \cdot 1 = r$$

Since $r > 0$, we replace the “volume element” $dx dy dz$ by $r dr d\phi dz$.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula


Further Topics in
Calculus

Summary -
Outlook and
Review



Example: Cylinder Coordinates

We compute the volume of a cylinder with radius $R > 0$ and height $h > 0$:



Transformation from Spherical Coordinates

Analysis 2

S.-J. Kimmerle

For spherical coordinates there holds

$$x = r \cos(\phi) \sin(\theta),$$

$$y = r \sin(\phi) \sin(\theta),$$

$$z = r \cos(\theta)$$

thus

$$g : \mathbb{R}^+ \times (-\pi, \pi] \times [0, \pi] \rightarrow \mathbb{R}^3, \begin{pmatrix} r \\ \phi \\ \theta \end{pmatrix} \mapsto \begin{pmatrix} x(r, \phi, \theta) \\ y(r, \phi, \theta) \\ z(r, \phi, \theta) \end{pmatrix} = \begin{pmatrix} r \cos(\phi) \sin(\theta) \\ r \sin(\phi) \sin(\theta) \\ r \cos(\theta) \end{pmatrix}.$$

Hence the Jacobian determinant (functional determinant) is:

$$\det \begin{pmatrix} \cos(\phi) \sin(\theta) & -r \sin(\phi) \sin(\theta) & r \cos(\phi) \cos(\theta) \\ \sin(\phi) \sin(\theta) & r \cos(\phi) \sin(\theta) & r \sin(\phi) \cos(\theta) \\ \cos(\theta) & 0 & -r \sin(\theta) \end{pmatrix} = -r^2 \sin(\theta)$$

Since $\sin(\theta) > 0$, we replace the “volume element” $dx \, dy \, dz$ by $r^2 \sin(\theta) \, dr \, d\phi \, d\theta$.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus


Summary -
Outlook and
Review



Example: Spherical Coordinates

We compute the mass of a homogeneous sphere with radius $R > 0$.

Let $\rho(x, y, z) = \rho_0 = \text{const}$ be the constant mass density.



Bell Curve, Continued I

An improper integral important for stochastics is the integral over the Gauss bell curve:

$$f(x) = \frac{1}{\sqrt{2\pi} \cdot \sigma} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right), \quad \mu \in \mathbb{R}, \sigma > 0.$$

For a probability we have to check:

$$\int_{-\infty}^{\infty} f(x) dx = 1.$$

However, for $f(x)$ there exists no closed form for a primitive (cf. Analysis 1):

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_{-\infty}^x \exp\left(-\frac{\tilde{x}^2}{2}\right) d\tilde{x}.$$

By now, we are able to check the normalization factor $\frac{1}{\sqrt{2\pi}}$ for the standard normal distribution ($\mu = 0, \sigma = 1$).

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Curve Integrals and Solids
of Revolution](#)[Integration of Functions with
Several Variables](#)[Transformation Formula](#)[Further Topics in
Calculus](#)[Summary -
Outlook and
Review](#)

Bell Curve, Continued II

We consider the square of the sought-after integral and apply Fubini:

$$\begin{aligned} & \int_{\mathbb{R}} \exp\left(-\frac{1}{2}x_1^2\right) dx_1 \cdot \int_{\mathbb{R}} \exp\left(-\frac{1}{2}x_2^2\right) dx_2 \\ &= \int \int_{\mathbb{R}^2} \exp\left(-\frac{1}{2}(x_1^2 + x_2^2)\right) dx_1 dx_2 \\ &= \int_0^{2\pi} \int_0^\infty \exp\left(-\frac{1}{2}r^2\right) r dr d\phi \\ &= \int_0^{2\pi} \int_0^\infty -\exp\left(-\frac{1}{2}r^2\right) \frac{d}{dr} \left(-\frac{1}{2}r^2\right) dr d\phi \\ &= 2\pi \cdot \lim_{b \rightarrow \infty} \left[-\exp\left(-\frac{1}{2}r^2\right) \right]_0^b \\ &= 2\pi \cdot (0 - (-1)) \\ &= 2\pi \end{aligned}$$

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

Theorem (Mean Value Theorem for Definite Double Integrals)

*Let $B \subseteq \mathbb{R}^2$ connected and closed,
then there exists for any continuous function $f : B \rightarrow \mathbb{R}$ a
point $(x^*, y^*) \in B$ such that*

$$\iint_B f \, dA = f(x^*, y^*) A \quad \text{with } A := \iint_B 1 \, dA$$

We call $\frac{1}{A} \iint f \, dA$ the **integral mean value**.

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

Mean Value Theorem for Integration II

Analysis 2

S.-J. Kimmerle

Note that:

- The mean value theorem holds for higher dimensional domains $B \subseteq \mathbb{R}^n$ as well.
- But this theorem fails for vector-valued functions.
- We do not consider indefinite integrals here.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



A **parameter integral** is an integral which integrand depends on parameters:

$$\int f(x_1, x_2, \dots, x_n) dx_i, \quad i \in \{1, \dots, n\}$$

Here x_i is considered as variable and all other $x_j, j \neq i$, are held fixed.

The x_j enter as parameters only.

In 2d we often write x for the variable and t for the parameter.

Introduction

Power series

Differentiation in
Higher DimensionsIntegration in
Higher DimensionsCurve Integrals and Solids
of RevolutionIntegration of Functions with
Several Variables

Transformation Formula

Further Topics in
CalculusSummary -
Outlook and
Review

Parameter Integrals - Fundamental Theorem

Analysis 2

S.-J. Kimmerle

A parameter integral is the inverse operation w.r.t. to partial differentiation:

For instance, let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ continuous and $s \in [c, d]$

$$F(x, s) := \int_c^s f(x, t) dt,$$

then

$$\frac{\partial}{\partial s} F(x, s) = \frac{\partial}{\partial s} \int_c^s f(x, t) dt = f(x, s).$$

Note that improper parameter integrals are more subtle.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



- Gamma function:

$$\Gamma(x) = \int_0^{\infty} t^{x-1} \exp(-t) dt, \quad t > 0$$

- Bessel functions:


$$J_n(x) := \frac{1}{\pi} \int_0^{\pi} \cos(x \sin(t) - nt) dt, \quad n \in \mathbb{Z}$$

- Laplace transform of a given function f

$$F(s) = \int_0^{\infty} f(t) \exp(-st) dt$$

- Gravitational potential V generated by a solid $M \subset \mathbb{R}^3$ with variable density ρ in a point $x \in \mathbb{R}^3 \subset M$ (G the gravitational constant)

$$V(\mathbf{x}) = -G \iiint_M \frac{\rho(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|} d\mathbf{y}$$



Parameter Integrals - Exchanging Integration & Differentiation

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review

Theorem (Leibniz Rule for Parameter Integrals)

Let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ on $[a, b]$ continuously partial differentiable w.r.t. x ,
then $F(x) := \int_c^d f(x, t) dt$ is differentiable (w.r.t. x) and

$$F'(x) = \frac{d}{dx} \int_c^d f(x, t) dt = \int_c^d \frac{\partial}{\partial x} f(x, t) dt.$$

We may exchange integration and differentiation in this situation.

In general the latter is not true.

The Fubini theorem may be demonstrated by considering in both orders one integral as a parameter integral.



- Using primitives
See integral tables
- By transformation formula
- Numerical methods
Relying on the Riemann approach, on interpolation,
...
- Special integrals
Tabulated, e.g., the error function
- Using series
- ...

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



- **Riemann integral**

(mainly the approach of our lectures)

↪ advantage w.r.t. improper integrals

- Cauchy integral

- **Lebesgue integral**

↪ important for stochastics

- Stieltjes integral

- Bochner integral

- Itô integral (stochastic integral)

- ... any many more ...

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Curve Integrals and Solids
of Revolution

Integration of Functions with
Several Variables

Transformation Formula

Further Topics in
Calculus

Summary -
Outlook and
Review



Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

**Further Topics in
Calculus**

Vector Analysis
Integral Transforms

Summary -
Outlook and
Review

- 1 Introduction
- 2 Power series
- 3 Differentiation in Higher Dimensions
- 4 Integration in Higher Dimensions
- 5 Further Topics in Calculus**
 - Vector Analysis
 - Integral Transforms
- 6 Summary - Outlook and Review



Let $D \subseteq \mathbb{R}^n$.

A function $\mathbf{F} : D \rightarrow \mathbb{R}^n$ with $n \geq 2$ is also called a **vector field**.

Example:

Let $f : D \rightarrow \mathbb{R}$, then ∇f is a vector field.

A vector field F is called **conservative** or **gradient field**, if there exists a function ϕ s.t.:

$$\mathbf{F} = \nabla \phi.$$

Then we call ϕ a (scalar) **potential (function)**.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Vector Analysis](#)[Integral Transforms](#)[Summary -
Outlook and
Review](#)

Path independence:

Let D be a simply connected domain.

$$\int_a^b \mathbf{F} \cdot d\mathbf{s} = \phi(\mathbf{b}) - \phi(\mathbf{a})$$

for a sufficiently smooth curve from \mathbf{a} to \mathbf{b} .

Note that the integral does not depend on the curve, but only on the start and end points, \mathbf{a} and \mathbf{b} .

How can we decide whether a vector field is conservative?

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Vector Analysis](#)[Integral Transforms](#)[Summary -
Outlook and
Review](#)

Characterization of Potential Functions

Consider a simply connected domain D .

The following statements are equivalent:

- $\int_a^b \mathbf{F} \cdot d\mathbf{s} = \phi(\mathbf{b}) - \phi(\mathbf{a})$ for a curve from \mathbf{a} to \mathbf{b} .
- The curve integral over \mathbf{F} along any closed curve is zero.
- $\nabla \times \mathbf{F} = 0$

We suppose that the involved curves are sufficiently smooth.

Note that ϕ is unique up to an additive constant.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



Examples: Potential

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



The fundamental theorem of Analysis 1

$$F(x) := \int_a^x F'(\tilde{x}) d\tilde{x}$$

or

$$\int_a^b F'(x) dx = F(b) - F(a)$$

cannot be easily generalized to several dimensions.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



Some generalizations are:

- Gradient theorem (“1d”)

$$\int_{\gamma} \nabla f(\mathbf{r}) \cdot d\mathbf{r} = f(\mathbf{r}(t_b)) - f(\mathbf{r}(t_a))$$

(γ a curve from $\mathbf{a} = \mathbf{r}(t_a)$ to $\mathbf{b} = \mathbf{r}(t_b)$ in \mathbb{R}^n with parametrization \mathbf{r})

- Stokes theorem (2d)

$$\iint_M (\nabla \times \mathbf{F}) \cdot d\mathbf{A} = \int_{\partial M} \mathbf{F} \cdot d\boldsymbol{\gamma}$$

(M a 2d submanifold of \mathbb{R}^3 , bounded by a curve γ representing ∂M)

- Gauss divergence theorem in 3d

$$\iiint_K \nabla \cdot \mathbf{F} dV = \iint_{\partial K} \mathbf{F} \cdot d\mathbf{A}$$

($K \subseteq \mathbb{R}^3$ compact. May be generalized to \mathbb{R}^n .)

Introduction

Power series

Differentiation in
Higher Dimensions


Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



Recall: gradient

$$\text{grad } f(\mathbf{x}) = \nabla f(\mathbf{x}) = \left(\frac{\partial}{\partial x_1} f(\mathbf{x}), \dots, \frac{\partial}{\partial x_n} f(\mathbf{x}) \right)^\top$$

Divergence (sources or sinks)

$$\text{div } \mathbf{F}(\mathbf{x}) = \nabla \cdot \mathbf{F}(\mathbf{x}) = \sum_{i=1}^n \frac{\partial}{\partial x_i} F_i(\mathbf{x})$$

Curl or rot(ation), in \mathbb{R}^3 (\mathbb{R}^2 also possible):

$$\text{curl } \mathbf{F}(\mathbf{x}) = \nabla \times \mathbf{F}(\mathbf{x}) = \begin{pmatrix} \frac{\partial}{\partial x_2} F_3(\mathbf{x}) - \frac{\partial}{\partial x_3} F_2(\mathbf{x}) \\ \frac{\partial}{\partial x_3} F_1(\mathbf{x}) - \frac{\partial}{\partial x_1} F_3(\mathbf{x}) \\ \frac{\partial}{\partial x_1} F_2(\mathbf{x}) - \frac{\partial}{\partial x_2} F_1(\mathbf{x}) \end{pmatrix}$$

Laplace operator

$$\text{div grad } f(\mathbf{x}) = \nabla \cdot \nabla f(\mathbf{x}) = \Delta f(\mathbf{x}) = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} f(\mathbf{x})$$

Introduction

Power series

Differentiation in
Higher Dimensions


Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



Integration by parts

$$\iiint_V g \nabla \cdot \mathbf{f} \, dV = \iint_{\partial V} g \mathbf{f} \cdot d\mathbf{A} - \iiint_V \nabla g \cdot \mathbf{f} \, dV$$

(May be generalized to \mathbb{R}^n .)

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Vector Analysis](#)[Integral Transforms](#)[Summary -
Outlook and
Review](#)

Fourier and Laplace transforms are useful tools for solving (partial) differential equations

Applications of the Fourier transform

- Information technology/electrical engineering:
Low-pass filter
- Spectroscopy (NMR, FTIR, ...)
- Acoustics
- Quantum mechanics
- ...
- Generalizes Fourier series from periodic to (some) non-periodic functions

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



Suppose

$$f(x) = \sum_{k=-\infty}^{\infty} c_k \exp(ik(\Delta\omega)x), \quad \Delta\omega = \frac{2\pi}{T}$$

with complex Fourier coefficients

$$\begin{aligned} c_k &= \frac{1}{T} \int_0^T f(x) \exp(-ik(\Delta\omega)x) dx \\ &= \frac{1}{T} \int_{-T/2}^{T/2} f(x) \overline{\exp(ik(\Delta\omega)x)} dx, \quad k \in \mathbb{Z} \end{aligned}$$

Introduce $\omega_k := k\Delta\omega$ and insert c_k into the series:

$$f(x) = \sum_{k=-\infty}^{\infty} \left(\frac{1}{T} \int_{-T/2}^{T/2} f(x) \overline{\exp(i\omega_k x)} dx \right) \exp(i\omega_k x) = \dots$$

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Vector Analysis](#)[Integral Transforms](#)[Summary -
Outlook and
Review](#)

$$f(x) = \sum_{k=-\infty}^{\infty} \left(\frac{1}{T} \int_{-T/2}^{T/2} f(\xi) \overline{\exp(i\omega_k \xi)} d\xi \right) \exp(i\omega_k x) \\ = \dots$$

$$= \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \left(\int_{-T/2}^{T/2} f(\xi) \overline{\exp(i\omega_k \xi)} d\xi \right) \exp(i\omega_k t) \Delta\omega$$

The series may be interpreted as a limit of Riemann sums that approximate an integral on $[-\infty, \infty]$ with refinement $\Delta\omega$ and grid points ω_k , $k \in \mathbb{Z}$.



Fourier Transform - Definition

For $T \rightarrow \infty$ (and $\Delta\omega \rightarrow 0$) we may expect convergence to an improper integral:

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) \exp(i\omega x) d\omega$$

with

$$F(\omega) = \int_{-\infty}^{\infty} f(x) \overline{\exp(i\omega x)} dx$$

Definition:

$$F(\omega) := \mathcal{F}\{f(x)\} := \int_{-\infty}^{\infty} f(x) \exp(-i\omega x) dx$$

is called **Fourier transform** (or spectral function) of $f : \mathbb{R} \rightarrow \mathbb{C}$, if the integral exists for all $\omega \in \mathbb{R}$.

$$\mathcal{F}^{-1}\{F(\omega)\} := f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) \exp(i\omega x) d\omega$$

is called **inverse Fourier transform** of $F : \mathbb{R} \rightarrow \mathbb{C}$, if the integral exists for all $x \in \mathbb{R}$.

[Introduction](#)[Power series](#)[Differentiation in
Higher Dimensions](#)[Integration in
Higher Dimensions](#)[Further Topics in
Calculus](#)[Vector Analysis](#)[Integral Transforms](#)[Summary -
Outlook and
Review](#)

Fourier Transform - Properties

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions


Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review

- Linearity
- Displacement
- Similarity
- Derivatives
- Modulation



Laplace transform

$$F(s) := \mathcal{L}\{f(t)\} := \int_0^{\infty} f(t) \exp(-st) dt,$$

is called **Laplace transform** of $f : \mathbb{R}^+ \rightarrow \mathbb{R}$,
if the integral exists for **some** $s \in \mathbb{R}$.

The inverse Laplace transform is hard to calculate.
Usually, reference tables are used.

The Laplace transform has similar properties as the
Fourier transform,
in particular

$$\mathcal{L}\{y'(t)\} = s\mathcal{L}\{y(t)\} - y(0) = sY(s) - y(0)$$

transforming differential equations (incl. initial values) into
algebraic equations

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Vector Analysis

Integral Transforms

Summary -
Outlook and
Review



- 1 Introduction
- 2 Power series
- 3 Differentiation in Higher Dimensions
- 4 Integration in Higher Dimensions
- 5 Further Topics in Calculus
- 6 **Summary - Outlook and Review**
 - Insight and outlook
 - References

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

**Summary -
Outlook and
Review**

Insight and outlook
References



Outlook and insight: What we did in Analysis 2

- **Differential calculus in higher dimensions (mainly 2d, 3d)**
- **Integral calculus in higher dimensions (mainly 2d, 3d)**
- Sequences of functions
- **Power series,**
Taylor series/Taylor expansions
Fourier series
- ...

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References



What we did in Analysis 2 - More Details

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References

- Sequences of functions, uniform convergence
- Power series, especially Taylor series, Fourier series, discrete Fourier transform
- Limits and continuity (for any d), curves (application: ECC)
- Differential calculus (for any d), especially optimization, linear regression
- Integrals along curves, on sectors; solids of revolution
- Integral calculus (for any d)
- Vector analysis (brief insight)
- Fourier transform (very brief insight)



What we have not discussed here

- Total differential, implicit functions, implicit differentiation
- Optimization with constraints, Lagrange multipliers
- Curvature
- **Differential equations**
- Integrals over curved surfaces, ...
- Laplace transform
- Fundamental theorem of Algebra to be proved by analytical methods ;)
- ...

... and all topics of this lecture may be discussed with all proofs and more details and extensions



References: recommended literature


 James Stewart: *Calculus*. Brooks/Cole, 6th edition, 2009.


 J. Hass, C. Heil, M.D. Weir: *Thomas' Calculus: Early transcendentals*. Pearsons, 14th edition, 1999.

In German only:

 Forster, O.: *Analysis 1*. Springer-Spektrum, 11. Auflage, 2013.

 Forster, O.: *Analysis 2*. Springer-Spektrum, 11. Auflage, 2017.

 Forster, O.: *Analysis 3*. Springer-Spektrum, 8. Auflage, 2017.


 Bronstein, Semendjajew, Musiol, Mühlig: *Taschenbuch der Mathematik*, Verlag Harri Deutsch, Frankfurt a. M./Thun, 4., überarb. u. erw. Aufl. der Neubearb., 1999.

Formula collections are also available in English.



Further sources (for sake of completeness)

-  Kimmerle, S.-J., Dvorsky, K., Ließ, H.-D., Avenhaus, R.: Time-to-Failure under Varying Thermal Stresses, Preprint, 2022.
-  Gerds, M.: *Mathematik II*, Vorlesungsskript, Bachelorstudiengänge Luft- und Raumfahrttechnik, Elektrotechnik und Bau- und Umweltingenieurwesen, Universität der Bundeswehr München, Herbsttrimester 2014.
-  Fetzer, A., Fränkel, H.: *Mathematik 1*, Springer, Berlin/Heidelberg, 6. Aufl., 2000.
-  Jäger W.: *Analysis I-III*, Vorlesungsskripte, Diplom-Studiengänge Mathematik und Physik, Universität Heidelberg, Wintersemester 1998/99 – Wintersemester 1999/2000.
-  Kimmerle, W., Stoppel, M.: *Analysis für Ingenieure, Mathematiker und Physiker*, edition Delkhofen, 4. Aufl., 1., korr. Nachdruck, 2015.
-  Wellisch, U.: Mathematik 2 Maschinenbau, lecture notes, TH Rosenheim, 2021.
-  Meyberg, K., Vachenaue, P.: Höhere Mathematik 1, Springer, 6. korr. Aufl., 1. korr. Nachdruck, 2003.
-  Papula, L.: *Mathematik für Ingenieure und Naturwissenschaftler, Band 1*, Springer Vieweg, Wiesbaden, 14. Auflage, 2014.
-  Papula, L.: *Mathematische Formelsammlung*, Springer Vieweg, Wiesbaden, 11. Auflage, 2014.
-  Tapken, G., Weiß, M.: *Mathematik für Informatiker*, Vorlesungsskript, BSc.-Studiengänge "Informatik" und "Wirtschaftsinformatik", OTH Regensburg, 2020.
-  Schneeberger, S.: *Analysis 1 & 2*, Vorlesungsskripte, BSc.-Studiengang WMA, TH Rosenheim, 2020/21.
-  Stingl, P.: *Mathematik für Fachhochschulen, Technik und Informatik*, Hanser, München, 8., aktual. Auflage, 2009.



General mathematical software:

- SCILAB, free software, scilab.org)
- MATLAB, campus licence
- Wolfram Mathematica
- ...

Computer algebra, i.e. “symbolic calculators”:

- MAPLE
- MATLAB, symbolic toolbox, campus licence ???
- Wolfram Alpha
- ...

For optimization, finite element methods, machine learning,
... specialized software packages exist.

Programming languages as Python, Java, C, ... are always
useful.

Introduction

Power series

Differentiation in
Higher Dimensions

Integration in
Higher Dimensions

Further Topics in
Calculus

Summary -
Outlook and
Review

Insight and outlook

References

