

- 1 Introduction
- 2 **Power series**
  - Sequences of Functions
  - Uniform Convergence
  - Continuity and Uniform Convergence
  - Power Series
  - Taylor Series
  - Fourier Series
- 3 Differentiation in Higher Dimensions
- 4 Integration in Higher Dimensions
- 5 Further Topics in Calculus
- 6 Summary - Outlook and Review


We know the concepts of  
functions,  
sequences, and  
series.

We are interested in series representing a function  $f(x)$  at  
every  $x$ :

$$f(x) = \sum_{j=0}^{\infty} a_j (x - x_0)^j$$

For this purpose we need the concept of a sequence of  
functions at first.

Sequence of partial sums :  $\left\{ \sum_{j=0}^n a_j (x - x_0)^j \right\}_{n \in \mathbb{N}_0}$



A sequence of elements of  $\mathbb{R}$  (cf. Analysis 1):

$$g : \mathbb{N} \rightarrow \mathbb{R}, n \mapsto a_n =: g(n)$$
$$\{a_n\}_{n \in \mathbb{N}} = \{a_n\}_{n \geq 1} = \{a_1, a_2, \dots, a_k, \dots\}$$

## Definition (Sequences of Functions)

Let  $D$  be a set. A mapping


$$g : D \times \mathbb{N} \rightarrow \mathbb{R}, (x, n) \mapsto f_n = f_n(x) =: g(x, n)$$

is called a **sequence** of functions  $f_n : D \rightarrow \mathbb{R}, n \in \mathbb{N}$ .

Other notations by writing the functions, e.g., are:

$$\{f_n\}_{n \in \mathbb{N}} = \{f_n\}_{n \geq 1} = \{f_1, f_2, f_3, \dots, f_n, \dots\}$$


The domain of definition  $D$  and the target area, here  $\mathbb{R}$ , have to be identical for all functions  $f_n$ .



# Example (Pointwise Convergence)


Consider  $\{f_n\}_{n \in \mathbb{N}}$ ,  $f_n : [0, 2] \rightarrow \mathbb{R}$  with

$$f_n(x) = \begin{cases} n^2 x, & 0 \leq x \leq \frac{1}{n}, \\ 2n - n^2 x, & \frac{1}{n} \leq x \leq \frac{2}{n}, \\ 0, & \frac{2}{n} \leq x \leq 2. \end{cases}$$



$$\tilde{n} > n > 1$$

This example exhibits that we may not swap the limit and the integral (another limit process) in general!



## Definition (Pointwise Convergence)


Let  $D$  be a set. A sequence  $\{f_n\}_{n \geq n_0}$  of functions  $f_n : D \rightarrow \mathbb{R}$  is called **pointwise convergent** to a function  $f : D \rightarrow \mathbb{R}$ , if and only if

$$\lim_{n \rightarrow \infty} f_n(x) = f(x) \quad \text{for any } x \text{ in } D.$$

Equivalently,

For any  $x \in D$  and  $\varepsilon > 0$   
there exists a  $N = N(x, \varepsilon) \geq n_0$  s.t.:

$$|f_n(x) - f(x)| < \varepsilon \quad \text{for any } x \text{ in } D \text{ and all } n \geq N.$$



## Definition (Uniform Convergence)

Let  $D$  be a set. A sequence  $\{f_n\}_{n \geq n_0}$  of functions  $f_n : D \rightarrow \mathbb{R}$  is called **uniformly convergent** to a function  $f : D \rightarrow \mathbb{R}$ , if and only if

For any  $x \in D$  and  $\varepsilon > 0$   
there exists a  $N = N(\varepsilon) \geq n_0$  s.t.:

$$|f_n(x) - f(x)| < \varepsilon \quad \text{for any } x \text{ in } D \text{ and all } n \geq N.$$

Notice that  $N$  may depend only on  $\varepsilon$  but not on the point  $x$ .

**Uniform** convergence always implies **pointwise** convergence, the opposite is not true (see last example).

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series


Fourier Series

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



# Example Uniform Convergence


Let  $x \in D = [0, 1)$ . The sequence

$$\left\{ \frac{1}{2^{x+n}} \right\}_{n \in \mathbb{N}}$$

converges uniformly:


Assume  $\varepsilon = \frac{1}{4}$  :

$$f_n(x) = \frac{1}{2^{x+n}} = \frac{1}{2^n} \cdot \frac{1}{2^x} < \frac{1}{4} \text{ for all } n \geq 2 \text{ for any } x \in D$$



$$f_n \xrightarrow{n \rightarrow \infty} f_\infty \equiv 0$$


as  $x \in D$  ( $x \neq 0$ !)



## Theorem (Uniform convergence preserves continuity)

*Let  $D \subseteq \mathbb{C}$  and  $f_n : D \rightarrow \mathbb{C}$ ,  $n \in \mathbb{N}$ , a sequence of continuous functions, that uniformly converge to a function  $f : D \rightarrow \mathbb{C}$ , then  $f$  is continuous.*

The limit of a uniformly convergent sequence of continuous functions, is again continuous.





# Example: Saw-tooth function

Analysis 2

S.-J. Kimmerle

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series

Fourier Series

Differentiation in Higher Dimensions


Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review

Let  $\sigma: \mathbb{R} \rightarrow \mathbb{R}$ ,


$$x \mapsto \begin{cases} 0 & ; x = 0 \\ \frac{\pi - x}{2} & ; 0 < x < 2\pi \\ \dots & \text{repeat periodically with period } 2\pi \end{cases}$$



Spoiler:  $\sigma(x) = \sum_{k=1}^{\infty} \frac{\sin(kx)}{x}$

↑  
whatever "equal" means here

$\sigma$  is discontinuous  $\Rightarrow$  this Fourier series is not uniformly convergent



## Definition (Uniform norm or sup norm)

Let  $D$  be a set and  $f : D \rightarrow \mathbb{C}$ .


We set

$$\|f\|_D := \sup_{x \in D} |f(x)|.$$

$\|\cdot\|_D$  defines a norm on  $D$ .

A function  $f$  is bounded iff  $\|f\| < \infty$ .

When misunderstandings are excluded, we just write  $\|f\|$  instead of  $\|f\|_D$ .



# Uniform Norm and Uniform Convergence

Analysis 2

S.-J. Kimmerle


By this notation we may reformulate the uniform convergence:

$\{f_n\}_{n \in \mathbb{N}}$  converges uniformly on  $D$

$$\iff \lim_{n \rightarrow \infty} \|f_n - f\|_D = 0$$

$$\|f_n - f\|_D \leq \varepsilon$$

$$\iff |f_n(x) - f(x)| \leq \varepsilon \quad \text{for all } x \in D$$



Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series


Fourier Series

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review



## Theorem (Weierstrass Convergence Criterion)

Let  $f_n : D \rightarrow \mathbb{C}$ ,  $n \in \mathbb{N}$ .

If


$$\sum_{n=0}^{\infty} \|f_n\|_D < \infty$$

then the series

$$\sum_{n=0}^{\infty} f_n = F$$

converges absolutely and uniformly on  $D$  to a function  $F : D \rightarrow \mathbb{C}$ .

For an example, see next slide



# Example: Convergence of a Power Series

Analysis 2

S.-J. Kimmerle

The series

$$\sum_{n=1}^{\infty} \frac{\cos(nx)}{n^2}$$

converges uniformly on  $\mathbb{R}$ .

$$f_n(x) = \frac{\cos(nx)}{n^2}, \quad n \in \mathbb{N} \quad ; \quad f_0 = 0$$

We know  $\sum_{n=1}^{\infty} \frac{1}{n^2} < \infty \quad \Rightarrow \quad \left\{ \begin{array}{l} \sum_{n=1}^{\infty} \frac{\cos(nx)}{n^2} \\ \text{converges} \\ \text{absolutely} \\ \text{\& uniformly on } D = \mathbb{R} \end{array} \right. \quad \left. \begin{array}{l} \|\cos(nx)\| \leq 1 \end{array} \right\}$

Introduction

Power series

Sequences of Functions

Uniform Convergence

Continuity and Uniform Convergence

Power Series

Taylor Series

Fourier Series

Differentiation in Higher Dimensions

Integration in Higher Dimensions

Further Topics in Calculus

Summary - Outlook and Review

